

P0001

241-145

Jacket 1 of 18

RESTRICTED USE CLASSIFICATION

Due to acute oral and dermal toxicity and avian hazards

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

THIMET® 15-G

soil and systemic insecticide

Active Ingredient: Phorate: O,O-diethyl S-[(ethylthio) methyl] phosphorodithioate.....	15.0
Inert Ingredients.....	85.0%
Total.....	100.0%

EPA Reg. No. 241-145

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO!



POISON



PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

FIRST AID


If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get medical attention.

If on skin, wash thoroughly with soap and water. Remove contaminated clothing and shoes. Wash clothing and decontaminate shoes before reuse.

If in eyes, immediately flush eyes with plenty of water. Get medical attention.

Antidote: Atropine is an antidote.



3818

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, nausea, vomiting, diarrhea, and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine.
Use according to label directions.

See Back Panel For Additional Precautionary Statements

IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

SOLD BY
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1990

**Net Weight: 50 lbs.
22.68 kg.**

©Registered Trademark of American Cyanamid Company

2-18

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER!**

Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

Do Not Breathe Dust

Wear a face mask or other respiratory equipment while emptying bags of product into a hopper. While emptying bags into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

Do Not Contaminate Food or Feed Products

Once a bag has been opened, use it completely. Make sure the hoppers are emptied while still in the field. Refer to STORAGE AND DISPOSAL statement for further instructions.

Keep all unprotected persons out of operating areas.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Keep Out of Reach of Domestic Animals.

Not for Use or Storage in or Around the Home.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife. Birds feeding in treated areas may be killed.

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Do not contaminate water when disposing of equipment washwaters.

1 of 18

REENTRY

After dusts have settled, do not enter or allow entry into the treated area until the 48 hour reentry interval has expired, unless the person entering the treated area is wearing the personal protective equipment listed below for early reentry.

For early reentry into treated areas wear a protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical-resistant gloves and chemical resistant shoes, coverings, or boots.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER. Area treated with THIMET 15-G on (date of application). Do not enter without protective clothing.

PERSONAL PROTECTIVE EQUIPMENT

WEAR THE FOLLOWING PROTECTIVE CLOTHING DURING LOADING, APPLICATION, EQUIPMENT REPAIR/CLEANING, AND DISPOSAL OF THE PESTICIDE: A protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical resistant gloves; chemical resistant shoes, shoe covering, or boots.

Unless a chemical resistant protective suit is otherwise required, mixers/loaders must wear a chemical resistant apron and goggles or a face shield.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATE: Long-sleeved shirt and long-legged pants; shoes and socks. Chemical resistant gloves must be available in the cab or cockpit and must be worn when exiting. This clothing is inadequate to protect you during equipment repair or cleaning, reentry, or pesticide disposal work. Do not allow contaminated clothing to come in contact with the insides of the cab.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or using the toilet.

5.8/18

AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear clean clothes. Do not reuse contaminated clothing. Personal clothing worn during work must be laundered separately from household articles. Store protective clothing separately from personal clothing. Clean or launder protective clothing after each use. Clothing and protection equipment heavily contaminated must be destroyed according to state and local regulations. **HEAVILY CONTAMINATED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.**

WORK SAFETY RULE

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposures should be allowed until it has been determined by means of blood tests that the cholinesterase level has returned to normal. Before using this product consult the National Pesticide Telecommunications Network for recommendations regarding such blood tests, poisoning management, emergency treatment, and other information regarding the toxicity of phorate. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7378.

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away, see First Aid section.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

6 of 18

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

STORAGE AND DISPOSAL

Storage

Store pesticide products in a secure locked area where children, unauthorized persons and animals cannot enter. Do not store in the same area with food or feed. Do not store opened bags.

Prohibitions

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Cover or incorporate sprays.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General

Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

- 218

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risk referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FROM A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID COMPANY'S LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THIMET 15-G. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

BEFORE USING, READ PRECAUTIONARY STATEMENTS.

THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

2/18

CROPS

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
BEANS At Planting	Mexican bean beetles	6.5-12.5 ozs. per 1,000 ft. of row for any row (minimum 30-inch spacing)	Drill granules to the side of the seed	Do not place THIMET 15-G granules in direct contact with seed. Do not graze or feed treated hay or forage to livestock. Do not apply later than 60 days before harvest.
	Leafhoppers Aphids Lygus bugs Thrips Mites Seedcorn maggots	6.0-9.3 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)		
	Seedcorn maggots		Apply granules in a band over the row	

**FIELD CORN
AND SWEET
CORN
(excluding
popcorn)**

At Planting

Corn
rootworms
Wireworms
White grubs
Seedcorn
maggots
Seedcorn
beetles
Flea beetles
Mites

8 ozs. per 1,000
ft. of row for any
row spacing
(minimum 30-inch
row spacing)

Place granules
in a 7-inch
band over the
row directly
behind the
planter shoe in
front of or
behind the
press wheel and
lightly
incorporate.

Do not use in-
furrow
application.
One additional
application can
be applied at
cultivation.

For field corn:
Accent*
herbicide and
Beacon**
herbicide may
be applied
after BANDED
applications of
THIMET 15-G.

Applications of
Accent
herbicide or
Beacon
herbicide
following
applications of
THIMET 15-G may
result in
temporary crop
injury.

**FIELD CORN
AND SWEET
CORN
(excluding
popcorn)**

**At
Cultivation**

Corn
rootworms

Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil.

Do not make any
applications of
THIMET 15-G
after
cultivation
treatment.

Do not harvest
within 30 days
of last
treatment.

**FIELD CORN
(excluding
sweet and
popcorn)**

**Chinch bug
nymphs
(Colorado,
Kansas, and
Nebraska)**

**At
Cultivation**

**Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil.
Apply as soon
as chinch bug
nymphs are
observed at
base of plants.
Granules must
be placed at
the infested
region of the
plants and
adequately
covered with
soil.**

**Later
infestations of
chinch bugs may
not be
controlled.**

**Do not make any
applications of
THIMET 15-G
after
cultivation
treatment.**

**Do not harvest
within 30 days
of last
treatment.**

**Do not graze or
cut for forage
within 30 days
of last
treatment.**

**Do not apply
more than 2
applications
per season.**

COTTON At Planting	Early season control of: Thrips Mites Leaf miners	4.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature moisture and tilth should be favorable for good germination and emergence.	Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 2.0 lbs. THIMET 15-G per acre for hill-dropped cotton. Do not feed treated hay or forage to livestock.
	Mites Thrips Aphids Black cutworms Leafhoppers White flies Leaf miners	8.0-12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing) 8.0-10 ozs per 1,000 ft. of row for 30-inch row spacing	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	Recommended for areas requiring and extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils. Do not feed treated hay or forage to livestock.

Crop injury may occur if applications of diuron containing herbicides such as *Karmex or ***Dixex are made in conjunction with soil-applied organo-phosphate pesticides.

Crop injury may occur if applications of diuron containing herbicides such as Karmex or Direx are made in conjunction with soil-applied organo-phosphate pesticides.

COTTON
Side-Dressing
(Irrigated Cotton Only)

Mites from spring to early summer

12.0-16.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)

12.0-13.3 ozs. per 1,000 ft. of row for 30 inch row spacing

Incorporate granules into the soil as a side-dress application deep enough to avoid disturbance by future cultivations.

Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest. Do not feed treated hay or forage to livestock.

Do not apply more than 2 applications per season

Crop injury may occur if applications of diuron containing herbicides such as Karmex or Direx are made in conjunction with soil-applied organo-phosphate pesticides.

PEANUTS At Planting	Thrips Leafhoppers	7.3 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage to livestock. Wait 90 days after treatment before harvesting peanuts.
-------------------------------	-----------------------	---	---	--

PEANUTS At Pegging	Southern corn rootworms and Leafhoppers	14.6 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules as a band over the fruiting zone at pegging time. Work into the top few inches of soil immediately.	Do not graze or feed treated hay or forage to livestock. Wait 90 days after treatment before harvesting peanuts. Do not apply more than 2 applications per season.
------------------------------	---	---	--	--

POTATOES At Planting	Aphids Leafhoppers Leaf miners Psyllids Wireworms Flea beetle larvae and reduction of flea beetle adults.	Light or Sandy Soils 15.0 ozs. per 1,000 ft. of row for any spacing (minimum 32-inch spacing) Heavy or Clay Soils 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)	Distribute the granules evenly in the furrow or band on each side of the row.	Wait 90 days after treatment before harvesting potatoes. Do not use for Colorado potato beetle control in the Long Island area of New York and the Del Marva Peninsula.
	Colorado potato beetle (early season control)	Heavy or Clay Soils: 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)		

14818

<p>SORGHUM At Planting</p>	<p>Greenbugs</p>	<p>8.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)</p>	<p>Place granules in a 7-inch band over the row OR drill 1 inch below and 2 inches to the side of the seed.</p>	<p>Do not place THIMET 15-G granules in direct contact with seed. A second application of granules may be applied at cultivation later in the season.</p>
---	------------------	--	---	---

<p>SORGHUM At Cultivation</p>	<p>Chinch bug nymphs (Colorado, Kansas, Nebraska)</p>	<p>Apply granules to base of plants just ahead of cultivation shovels so as to cover granules with soil. Apply as soon as chinch bug nymphs are observed at base of plants. Granules must be placed at the infested region of the plants and adequately covered with soil.</p>	<p>Later infestation of chinch bugs may not be controlled.</p> <p>Do not make any application of THIMET 15-G after cultivation treatment.</p> <p>Do not feed foliage before grain harvest.</p> <p>Do not apply within 30 days of harvest.</p> <p>Do not apply more than 2 applications per season.</p>
--	---	--	--

SOYBEANS At Planting	For the early season control of: Mexican bean beetles Leafhoppers Lygus bugs Seedcorn maggots Thrips Aphids Mites	12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch row spacing)	Drill granules to the side of the seed.	Do not place THIMET 15-G granules in direct contact with seed. Do not feed the foliage of treated soybeans. Crop injury may result when preplant incorporated or preemergence applications of metribuzin herbicides are used in conjunction with this product or other soil-applied organo-phosphate pesticides.
	<hr/> Seedcorn maggots		<hr/> Place granules in a 7-inch band over the row directly behind the planter shoe and in front of the press wheel.	
SUGAR BEETS	Aphids Beets root maggots Leafhoppers Mites Leaf miners	6.0 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch spacing)	Drill granules to the side of the seed OR place granules in a band over the row.	Do not place THIMET 15-G granules in direct contact with the seed. Do not apply within 30 days of harvest. Do not feed sugar beet tops or silage to dairy cattle. Consult your state experiment station or state extension service for proper timing of application in your area.

SUGAR BEETS Post-emergence	Mites Aphids	6.5-10.0 lbs. per acre	Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.	Do not apply by air. Do not apply more than 2 applications per season.
SUGAR CANE (Florida)	Wireworms	26.0 lbs. per acre	Apply in a 10-12 inch band directly on the seed piece and surrounding soil in the open furrow immediately before covering.	Do not graze or feed treated hay or forage to livestock.
WHEAT At Planting	Grasshoppers ¹ Aphids (Western and North Central States only) Hessian fly	1.6 ozs. per 1,000 ft. of row for any row spacing (minimum 8-inch spacing)	Apply the granules in the seed furrow.	Do not feed or graze foliage within 70 days of treatment. Do not make any later applications after planting time treatment. ¹ Grasshoppers controlled in winter wheat only. Do not harvest within 70 days of application.

*Accent and Karmex are registered trademarks of E.I. duPont deNemours and Company
**Beacon is a registered trademark of Ciba Geigy Corporation.
***Direx is a registered trademark of Griffin Ag Product Co. Inc.

17-812

TYPICAL APPLICATION RATES FOR VARIOUS ROW SPACINGS

	Rate Oz./1,000 Ft. of Row												
	<u>1.6 oz</u>	<u>2 oz</u>	<u>4 oz</u>	<u>6 oz</u>	<u>8 oz</u>	<u>10 oz</u>	<u>12 oz</u>	<u>14 oz</u>	<u>16 oz</u>	<u>18 oz</u>	<u>20 oz</u>	<u>22 oz</u>	<u>24</u>
	EQUIVALENT POUNDS PER ACRE												
40"				4.9	6.5	8.2	9.8	11.4	13.1	14.7	16.3	18.0	19
38"				5.2	6.9	8.6	10.3	12.0	13.7	15.5	17.2	18.9	20
36"				5.4	7.3	9.1	10.9	12.7	14.5	16.3	18.2	20.0	21
34"				5.6	7.7	9.7	11.6	13.5	15.4	17.4	19.3	21.3	23
32"				6.1	8.2	10.2	12.3	14.3	16.3	18.3	20.5	22.4	24
30"				6.5	8.7	10.9	13.1	15.2	17.4	19.6	21.8	24.0	26
26"				7.1	9.5	11.8	14.2	16.6	18.9	21.3	23.7	26.0	28
24"			5.4	8.2	10.9	13.6	16.3	19.1	21.8	24.5	27.2	30.0	32
22"			5.9	8.9	11.9	14.9	17.9	21.0	24.0	26.9	29.9	33.0	35
20"			6.5	9.8	13.1	16.3	19.6	22.8	26.1	29.3	32.6	35.9	39
16"			8.2	12.3	16.3	20.5	24.6	28.7	32.7	36.8	40.9		
12"		5.4	10.9	16.3	21.8	27.2	32.7	38.1	43.6				
8"	6.5	8.2	16.3	24.6	32.7	40.9							
6"	8.7	10.9	21.8	32.7	43.6								

18 of 18

CALIBRATION INFORMATION

FIRST READ THE LABEL.

It is important that applicator equipment be properly set to deliver the labeled rate. The table below contains suggested starting gauge settings for calibration at planting speeds of 5 or 7 mph and application rates of 4.5, 6.0 or 9.0 oz. 1,000 ft. of row.

NOTE: The settings in this table SHOULD ONLY BE USED AS STARTING POINTS. Continually check the amount of THIMET 15-G used against a known length of row and make further adjustments accordingly. Also check calibration occasionally to make sure equipment wear, changing moisture conditions, etc. have not caused a change in flow rate.

SUGGESTED GRANULAR APPLICATOR SETTING¹

Applicator	Planter Speed 5 mph			Planter Speed 7 mph		
	4.5 oz.	6.0 oz.	9.0 oz.	4.5 oz.	6.0 oz.	9.0 oz.
John Deere MaxEmerge 2	14	17	23	17	21	33
John Deere Max-Emerge (Odd Notches 5-15-25-35)	9	11	14	11	13	17
John Deere Max-Emerge (Even Notches 0-10-20-30)	9	13	20	13	19	26
John Deere 71 Flexi-Planter (John Deere Metal Hoppers)	1/23	1/26	2/3	1/26	1/32	2/14
New International Harvester (2 gauges hopper)	1/9	2/2	2/6	2/2	2/5	3/3
Old International Harvester (1 gauge hopper)	1/6	1/8	2/3	1/9	2/2	2/9
New Gandy	15	18	22	18	21	25
New Noble	6	8	11	8	11	14
Old Noble	8	11	16	12	15	22
Allis Chalmers (78 and 79 Series)	4	5	7	4	5	7
Allis Chalmers #385	10	11	14	11	13	18

¹Suggested starting point only.

PM14

241-145

Yellowe 1 of 73

JUN 22 1992

~~CONFIDENTIAL~~

5414579

American Cyanamid Company
Agricultural Research Division
P.O. Box 400
Princeton, NJ 08543-0400

Gentlemen:

Subject: Thimet 15G Soil and Systemic Insecticide
EPA Registration No. 241-145 ✓
Thimet 20G Soil and Systemic Insecticide
EPA Registration No. 241-257
Revised Bulk Container Labeling Submitted
February 28, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the revised labeling:
 - a. Add a statement on the dispensing tube labeling prohibiting use of the tube for other purposes.
 - b. On your Thimet 20G Soil and Insecticide product, reinstate the following restriction for sugar beets which appears on your labeling accepted November 21, 1991:

Do not broadcast

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

63607:I:WP50:Mautz:WP14-5:KEVRIC:06/11/92:07/10/92:DD

BEST AVAILABLE COPY

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

RESTRICTED USE CLASSIFICATION

Due to acute oral and dermal toxicity and avian hazards.

retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

THIMET® 15-G soil and systemic insecticide

Active Ingredient: Phorate: O,O-diethyl S-[(ethylthio) methyl] phosphorodithioate.....	15.0%
Inert Ingredients.....	85.0%
Total.....	100.0%

EPA Reg. No. 241-145

KEEP OUT OF REACH OF CHILDREN



DANGER/PELIGRO!



POISON

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

FIRST AID

If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get medical attention.

If on skin, wash thoroughly with soap and water. Remove contaminated clothing and shoes. Wash clothing and decontaminate shoes before reuse.

If in eyes, immediately flush eyes with plenty of water. Get medical attention.

Antidote: Atropine is an antidote.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUN 22 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-145

4423
11

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, nausea, vomiting, diarrhea, and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine.

Use according to label directions.

See Back Panel For Additional Precautionary Statements

SOLD BY
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1991

Net Weight: 1500 lbs.
681 kg.

Registered Trademark of American Cyanamid Company

5573
0

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER!**

Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

Do Not Breathe Dust

Wear a face mask or other respiratory equipment and fill downwind while emptying container into equipment. Do not allow dust to reach the breathing zone.

Do Not Contaminate Food or Feed Products

When container is not in use, make sure the discharge ends of the dispensing tube and container are capped or plugged (caps and plugs are provided). Make sure the hoppers on the planter are emptied while still in the field. Refer to STORAGE AND DISPOSAL statement for further instructions.

Keep all unprotected persons out of operating areas.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Keep Out of Reach of Domestic Animals.

Not for Use or Storage in or Around the Home.

REENTRY

After dusts have settled, do not enter or allow entry into the treated area until the 48 hour reentry interval has expired, unless the person entering the treated area is wearing the personal protective equipment listed below for early reentry.

For early reentry into treated areas wear a protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical-resistant gloves and chemical resistant shoes, coverings, or boots.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER. Area treated with THIMET 15-G on (date of application). Do not enter without protective clothing.

10, 23

PERSONAL PROTECTIVE EQUIPMENT

WEAR THE FOLLOWING PROTECTIVE CLOTHING DURING LOADING, APPLICATION, EQUIPMENT REPAIR/CLEANING, AND DISPOSAL OF THE PESTICIDE: A protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical resistant gloves; chemical resistant shoes, shoe covering, or boots.

Unless a chemical resistant protective suit is otherwise required, mixers/loaders must wear a chemical resistant apron and goggles or a face shield.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATE: Long-sleeved shirt and long-legged pants; shoes and socks. Chemical resistant gloves must be available in the cab or cockpit and must be worn when exiting. This clothing is inadequate to protect you during equipment repair or cleaning, reentry, or pesticide disposal work. Do not allow contaminated clothing to come in contact with the insides of the cab.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or using the toilet.

AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear clean clothes. Do not reuse contaminated clothing. Personal clothing worn during work must be stored separately from household articles. Store protective clothing separately from personal clothing. Clean or launder protective clothing after each use. Clothing and protection equipment heavily contaminated must be destroyed according to state and local regulations. **HEAVILY CONTAMINATED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.**

WORK SAFETY RULE

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposures should be allowed until it has been determined by means of blood tests that the cholinesterase level has returned to normal. Before using this product consult the National Pesticide Telecommunications Network for recommendations regarding such blood tests, poisoning management, emergency treatment, and other information regarding the toxicity of phorate. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7378.

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away, see First Aid section.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife. Birds feeding in treated areas may be killed.

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

Bulk Handling

See BULK HANDLING INSTRUCTIONS section on the container for details.

STORAGE AND DISPOSAL

Storage

Store pesticide products in a secure locked area where children, unauthorized persons and animals cannot enter. Do not store in the same area with food or feed.

Prohibitions

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Cover or incorporate spills.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Return dispensing tube with empty bulk container to the point of purchase. DO NOT DISCONNECT DISPENSING TUBE FROM CONTAINER UNTIL CONTAINER IS EMPTY AND THE DISPENSING TUBE HAS BEEN CLEANED (See Clean-out procedure under the BULK HANDLING INSTRUCTIONS section for details). If disconnecting the dispensing tube, BE SURE TO PLUG THE DISCHARGE END OF THE CONTAINER WITH THE PLUG PROVIDED AND CAP THE FEED END OF THE DISPENSING TUBE.

2/153
1)

DISCLAIMER

Label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risk referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO PAYMENT OF THE PURCHASE PRICE OF THIMET 15-G. In no case shall Cyanamid or its seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

BEFORE USING, READ PRECAUTIONARY STATEMENTS.

THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

1,225
U

CROPS

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
BEANS At Planting	Mexican bean beetles	6.5-12.5 ozs. per 1,000 ft. of row for any row (minimum 30-inch spacing)	Drill granules to the side of the seed	Do not place THIMET 15-G granules in direct contact with seed. Do not graze or feed treated hay or forage to livestock. Do not apply later than 60 days before harvest.
	Leafhoppers Aphids Lygus bugs Thrips Mites Seedcorn maggots	6.0-9.3 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)		
	Seedcorn maggots		Apply granules in a band over the row	

FIELD CORN
AND SWEET
CORN
(excluding
popcorn)

At Planting

Corn
rootworms
Wireworms
White grubs
Seedcorn
maggots
Seedcorn
beetles
Flea beetles
Mites

8 ozs. per 1,000
ft. of row for any
row spacing
(minimum 30-inch
row spacing)

Place granules
in a 7-inch
band over the
row directly
behind the
planter shoe in
front of or
behind the
press wheel and
lightly
incorporate.

Do not use in-
furrow
application.
One additional
application can
be applied at
cultivation.

For field corn:
Accent*
herbicide and
Beacon**
herbicide may
be applied
after BANDED
applications of
THIMET 15-G.

Applications of
Accent
herbicide or
Beacon
herbicide
following
applications of
THIMET 15-G may
result in
temporary crop
injury.

FIELD CORN
AND SWEET
CORN
(excluding
popcorn)

At
Cultivation

Corn
rootworms

Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil.

Do not make any
applications of
THIMET 15-G
after
cultivation
treatment.

Do not harvest
within 30 days
of last
treatment.

BEST AVAILABLE COPY

11 of 23

ELD CORN
(including
sweet and
popcorn)

Chinch bug
nymphs
(Colorado,
Kansas, and
Nebraska)

At
Cultivation

Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil.
Apply as soon
as chinch bug
nymphs are
observed at
base of plants.
Granules must
be placed at
the infested
region of the
plants and
adequately
covered with
soil.

Later
infestations of
chinch bugs may
not be
controlled.

Do not make any
applications of
THIMET 15-G
after
cultivation
treatment.

Do not harvest
within 30 days
of last
treatment.

Do not graze or
cut for forage
within 30 days
of last
treatment.

Do not apply
more than 2
applications
per season.

COTTON At Planting	Early season control of: Thrips Mites Leaf miners	4.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature moisture and tilth should be favorable for good germination and emergence.	Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 2.0 lbs. THIMET 15-G per acre for hill- dropped cotton. Do not feed treated hay or forage to livestock.
	Mites Thrips Aphids Black cutworms Leafhoppers White flies Leaf miners	8.0-12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)	Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.	Recommended for areas requiring an extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils. Do not feed treated hay or forage to livestock.
COTTON Side- Dressing (Irrigated Cotton Only)	Mites from spring to early summer	12.0-16.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)	Incorporate granules into the soil as a side-dress application deep enough to avoid disturbance by future cultivations.	Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest. Do not feed treated hay or forage to livestock.

BEST AVAILABLE COPY

Do not apply
more than 2
applications
per season

13823

PEANUTS At Planting	Thrips Leafhoppers	7.3 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage to livestock. Wait 90 days after treatment before harvesting peanuts.
PEANUTS At Pegging	Southern corn rootworms and Leafhoppers	14.6 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)	Distribute the granules as a band over the fruiting zone at pegging time. Work into the top few inches of soil immediately.	Do not graze or feed treated hay or forage to livestock. Wait 90 days after treatment before harvesting peanuts. Do not apply more than 2 applications per season.
POTATOES At Planting	Aphids Leafhoppers Leaf miners Psyllids Wireworms Flea beetle larvae and reduction of flea beetle adults.	Light or Sandy Soils 15.0 ozs. per 1,000 ft. of row for any spacing (minimum 32-inch spacing)	Distribute the granules evenly in the furrow or band on each side of the row.	Wait 90 days after treatment before harvesting potatoes.
	Colorado potato beetle (early season control)	Heavy or Clay Soils: 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)		Do not use for Colorado potato beetle control in the Long Island area of New York and the Del Marva Peninsula.

11823

<p>SORGHUM At Planting</p>	<p>Greenbugs</p>	<p>8.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)</p>	<p>Place granules in a 7-inch band over the row OR drill 1 inch below and 2 inches to the side of the seed.</p>	<p>Do not place THIMET 15-G granules in direct contact with seed. A second application of granules may be applied at cultivation.</p>
<p>SORGHUM At Cultivation</p>	<p>Chinch bug nymphs (Colorado, Kansas, Nebraska)</p>		<p>Apply granules to base of plants just ahead of cultivation shovels so as to cover granules with soil. Apply as soon as chinch bug nymphs are observed at base of plants. Granules must be placed at the infested region of the plants and adequately covered with soil.</p>	<p>Later infestation of chinch bugs may not be controlled.</p> <p>Do not make any application of THIMET 15-G after cultivation treatment.</p> <p>Do not feed foliage before grain harvest.</p> <p>Do not apply within 30 days of harvest.</p> <p>Do not apply more than 2 applications per season.</p>

11-23

<p>BEANS</p> <p>At Planting</p>	<p>For the early season control of:</p> <p>Mexican bean beetles Leafhoppers Lygus bugs Seedcorn maggots Thrips Aphids Mites</p> <hr/> <p>Seedcorn maggots</p>	<p>12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch row spacing)</p>	<p>Drill granules to the side of the seed.</p> <hr/> <p>Place granules in a 7-inch band over the row directly behind the planter shoe and in front of the press wheel.</p>	<p>Do not place THIMET 15-G granules in direct contact with seed. Do not feed the foliage of treated soybeans. Crop injury may result when preplant incorporated or preemergence applications of metribuzin herbicides are used in conjunction with this product or other soil-applied organo-phosphate pesticides.</p>
<p>SUGAR BEETS</p> <p>At Planting</p>	<p>Aphids Beet root maggots Leafhoppers Mites Leaf miners</p>	<p>6.0 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch spacing)</p>	<p>Drill granules to the side of the seed OR place granules in a band over the row.</p>	<p>Do not place THIMET 15-G granules in direct contact with the seed.</p> <p>Do not apply within 30 days of harvest.</p> <p>Do not feed sugar beet tops or silage to dairy cattle.</p> <p>Consult your state experiment station or state extension service for proper timing of application in your area.</p>

<p>SUGAR BEETS</p> <p>Post-emergence</p>	<p>Mites Aphids</p>	<p>6.5-10.0 lbs. per acre</p>	<p>Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.</p>	<p>Do not apply by air.</p> <p>Do not apply more than 2 applications per season.</p>
<p>SUGAR CANE (Florida)</p>	<p>Wireworms</p>	<p>26.0 lbs. per acre</p>	<p>Apply in a 10-12 inch band directly on the seed piece and surrounding soil in the open furrow immediately before covering.</p>	<p>Do not graze or feed treated hay or forage to livestock.</p>
<p>WHEAT</p> <p>At Planting</p>	<p>Grasshoppers¹ Aphids (Western and North Central States only) Hessian fly</p>	<p>1.6 ozs. per 1,000 ft. of row for any row spacing (minimum 8-inch spacing)</p>	<p>Apply the granules in the seed furrow.</p>	<p>Do not feed or graze foliage within 70 days of treatment. Do not make any later applications after planting time treatment.</p> <p>¹Grasshoppers controlled in winter wheat only.</p> <p>Do not harvest within 70 days of application.</p>

*Accent is a registered trademark of E.I. duPont de Nemours and Company.

**Beacon is a registered trademark of Ciba Geigy Corporation.

17423
✓

USE RATES AS POUNDS PER ACRE
(based on various row spacings and application rates)

Rate Oz./1,000 Ft. of Row

	<u>1.6 oz</u>	<u>2 oz</u>	<u>4 oz</u>	<u>6 oz</u>	<u>8 oz</u>	<u>10 oz</u>	<u>12 oz</u>	<u>14 oz</u>	<u>16 oz</u>	<u>18 oz</u>	<u>20 oz</u>	<u>22 oz</u>	<u>24 oz</u>
40"				4.9	6.5	8.2	9.8	11.4	13.1	14.7	16.3	18.0	19.6
38"				5.2	6.9	8.6	10.3	12.0	13.7	15.5	17.2	18.9	20.6
36"				5.4	7.3	9.1	10.9	12.7	14.5	16.3	18.2	20.0	21.8
34"				5.6	7.7	9.7	11.6	13.5	15.4	17.4	19.3	21.3	23.2
32"				6.1	8.2	10.2	12.3	14.3	16.3	18.3	20.5	22.4	24.6
30"				6.5	8.7	10.9	13.1	15.2	17.4	19.6	21.8	24.0	26.1
26"				7.1	9.5	11.8	14.2	16.6	18.9	21.3	23.7	26.0	28.4
24"			5.4	8.2	10.9	13.6	16.3	19.1	21.8	24.5	27.2	30.0	32.7
22"			5.9	8.9	11.9	14.9	17.9	21.0	24.0	26.9	29.9	33.0	35.9
20"			6.5	9.8	13.1	16.3	19.6	22.8	26.1	29.3	32.6	35.9	39.1
16"			8.2	12.3	16.3	20.5	24.6	28.7	32.7	36.8	40.9		
12"		5.4	10.9	16.3	21.8	27.2	32.7	38.1	43.6				
8"	6.5	8.2	16.3	24.6	32.7	40.9							
6"	8.7	10.9	21.8	32.7	43.6								

CALIBRATION INFORMATION

FIRST READ THE LABEL

It is important that applicator equipment be properly set to deliver the labeled rate. The table below contains suggested starting gauge settings for calibration at planting speeds of 5 or 7 mph and application rates of 4.5, 6.0 or 9.0 oz. 1,000 ft. of row.

NOTE: The settings in this table SHOULD ONLY BE USED AS STARTING POINTS. Continually check the amount of THIMET 15-G used against a known length of row and make further adjustments accordingly. Also check calibration occasionally to make sure equipment wear, changing moisture conditions, etc. have not caused a change in flow rate.

SUGGESTED GRANULAR APPLICATOR SETTING¹

Applicator	Planter Speed 5 mph			Planter Speed 7 mph		
	4.5 oz.	6.0 oz.	9.0 oz.	4.5 oz.	6.0 oz.	9.0 oz.
John Deere MaxEmerge 2	14	17	23	17	21	33
John Deere Max-Emerge (Odd Notches 5-15-25-35)	9	11	14	11	13	17
John Deere Max-Emerge (Even Notches 0-10-20-30)	9	13	20	13	19	26
John Deere 71 Flexi-Planter (John Deere Metal Hoppers)	1/23	1/26	2/3	1/26	1/32	2/14
New International Harvester (2 gauges hopper)	1/9	2/2	2/6	2/2	2/5	3/3
Old International Harvester (1 gauge hopper)	1/6	1/8	2/3	1/9	2/2	2/9
New Gandy	15	18	22	18	21	25
New Noble	6	8	11	8	11	14
Old Noble	8	11	16	12	15	22
Allis Chalmers (78 and 79 Series)	4	5	7	4	5	7
Allis Chalmers #385	10	11	14	11	13	18

¹Suggested starting point only.

1905-2

BULK HANDLING INSTRUCTIONS

(USE ONLY WITH LOCK'N LOAD COMPATIBLE SYSTEMS)

Start-up Procedure

1. Position the truck carrying the mini bulk container perpendicular to the insecticide hoppers on the planter. The clearance between the truck and hoppers should be 3-4 feet.
2. Connect drive motor to motor mount located in the rear of the mini bulk container. This is accomplished by inserting the motor pivot rods into the motor mount and matching mating couplings between the motor and container.
3. Connect the electrical control wire from the motor to the mini bulk container's electrical socket. The container socket is located up and to the right of the motor mount.
4. Connect dispensing tube to mini bulk container and lock into position. The system will not operate unless the dispensing tube is in a locked position.
5. **WARNING: BE SURE TO OBSERVE ALL SAFETY PRECAUTIONS AND USE PROPER SAFETY EQUIPMENT WHEN CONNECTING BATTERY CABLES. KEEP SPARKS, FLAMES, AND CIGARETTES AWAY FROM BATTERY. WEAR EYE PROTECTION. DO NOT LEAN OVER BATTERY WHILE DRIVE MOTOR IS IN OPERATION. BE SURE VENT CAPS ARE TIGHT AND LEVEL. FOR GREATER SAFETY, PLACE DAMP CLOTH OVER VENT CAPS. THE MINI BULK HAS A 12 VOLT SYSTEM. USE A 12 VOLT BATTERY ONLY.**

Connect power cables from drive motor to the proper battery terminals in the following order:

- a. Clamp the red cable to the positive (+) terminal
 - b. The black cable to the negative (-) terminal
 - c. Stand back from battery
 - d. Start vehicle containing the battery being used and keep it running while operating the drive motor on the mini bulk
 - e. Remove cables in reverse order
6. Connect discharge end of the dispensing tube to the insecticide hopper (LOCK'N LOAD only). Be sure valves are completely engaged.
 7. Pull the slide gate handle out to allow free flow of the granular insecticide. The handle is located above the container auger on the dispensing tube side of the mini bulk container.
 8. To operate the auger system, press the green start button; to stop press the red button. These buttons are located on the discharge end of the dispensing tube. A second stop switch is located on the dispensing tube side of the mini bulk container.
 9. The auger will shut down automatically once the insecticide hopper is filled.
 10. Before disconnecting dispensing tube from insecticide hopper, be sure auger is not operating. When not in use, make sure the dispensing tube remains connected to the container and the discharge end is capped and placed on the transporting vehicle bed.

CAUTION: IN CASE OF SYSTEM FAILURE, DO NOT ATTEMPT REPAIR. CALL YOUR LOCAL AGRICENTER FOR HELP. IF FILLED WITH PRODUCT, DO NOT DISCONNECT DISPENSING TUBE FROM MINI BULK CONTAINER.

Clean-out Procedure

1. Close slide gate on the mini bulk container. Be sure the dispensing tube is connected to the mini bulk container and the insecticide hopper.
2. Run the auger for approximately 1 1/2 minutes. During this operation hold the dispensing tube level with the discharge end. This will facilitate emptying the remaining product in the dispensing tube.
3. Disconnect the dispensing tube from the mini bulk container. Immediately plug this end of the dispensing tube with plug provided, and lock it into position.
4. Cap and lock discharge end of auger on the mini bulk container.
5. Disconnect the dispensing tube from the insecticide hopper and cap the end.

Transporting

When transporting the mini bulk system, be sure it is properly secured.

THIMET® soil and systemic insecticide

DANGER/PROLIGRO!



POISON



DISPENSING TUBE LABEL INSTRUCTIONS

1. Refer to the container label for Precautionary Statements
2. Refer to the container label for Bulk Handling Instructions (start-up and clean-out procedures).
3. Container Disposal: Return dispensing tube with empty mini bulk container to the point of purchase. DO NOT DISCONNECT DISPENSING TUBE FROM CONTAINER UNTIL CONTAINER IS EMPTY AND THE DISPENSING TUBE HAS BEEN CLEANED (See Clean-out procedure under the BULK HANDLING INSTRUCTIONS section for details). If disconnecting the dispensing tube, BE SURE TO PLUG THE DISCHARGE END OF THE CONTAINER WITH THE PLUG PROVIDED AND CAP THE FEED END OF THE DISPENSING TUBE.

CAUTION: IN CASE OF SYSTEM FAILURE, DO NOT ATTEMPT REPAIR. CALL YOUR LOCAL AGRICENTER FOR HELP. IF FILLED WITH PRODUCT, DO NOT DISCONNECT DISPENSING TUBE FROM MINI BULK CONTAINER.

21 of 23

5415682 $\frac{300}{17}$

5415686 $\frac{300}{17}$

JUN 22 1992

~~JUN 22 1991~~

American Cyanamid Company
Agricultural Research Division
P.O. Box 400
Princeton, NJ 08543-0400

Gentlemen:

Subject: Sugar Beet Supplemental Labeling for Beet
Root Maggots
Thimet 15G Soil and Systemic Insecticide
EPA Registration No. 241-145 ✓
Thimet 20G Soil and Systemic Insecticide
EPA Registration No. 241-257
Your Amendment Submitted March 18, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the labeling changes listed below before you release the Thimet 20 Soil and Systemic Insecticide product for shipment bearing the revised labeling:
 - Reinstate the following restriction for sugar beets which appear on your labeling accepted November 21, 1991:

Do not broadcast.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

63607:I:WP50:Mautz:WP14-5:KEVRIC:06/11/92:07/10/92:DD

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

22 of 23

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

with COMMENTS
to EPA Letter Dated:

JUN 22 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-145

RESTRICTED USE PESTICIDE

Due to acute oral and dermal toxicity and bird toxicity.

For retail sale to, and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

EPA REG. 241-145

FOR USE IN SUGAR BEETS FOR CONTROL OF BEET ROOT MAGGOTS

BEFORE USING, READ PRECAUTIONARY
STATEMENTS ON BAG LABEL

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the end of treated rows, turns and loading areas by deep discing immediately after treating fields.

CROP	PESTS CONTROLLED	RATE OF THIMET 15-G	APPLICATION	REMARKS
Sugar Beets	Beet root maggots	6.5 - 10.0 lbs. per acre.	Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.	Do not apply within 30 days of harvest.
Post-emergence				Do not apply in air. Do not apply more than 2 applications per season. Do not feed sugar beet tops or silage to dairy cattle.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

*Registered Trademark of American Cyanamid

PE

2/11/92 SD



CYANAMID

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470