

Reg # 241-145

PM- 14

NOV 19 1992

1720

5424183

$\frac{655}{17}$

American Cyanamid Company  
Agricultural Research Division  
P.O. Box 400  
Princeton, NJ 08543-0400

Gentlemen:

Subject: Grazing Restriction Revisions for Cotton and Sugar Beets  
Thimet 10G Soil and Systemic Insecticide  
EPA Registration No. 241-53  
Thimet 15G Soil and Systemic Insecticide  
EPA Registration No. 241-145  
Thimet 20G Soil and Systemic Insecticide  
EPA Registration No. 241-257  
Your Labeling Submitted June 17, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you:

1. Make the labeling change listed below before you release the Thimet 20G Soil and Systemic Insecticide product for shipment bearing the revised labeling.
  - Reinstate the following restriction for sugar beets which appears on your labeling accepted November 21, 1991 and February 22, 1992.

Do not broadcast.

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

63621:I:WP5.1-01:Forrest:01:11/02/92:11/29:92

2720

-2-

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

3 20

**RESTRICTED USE CLASSIFICATION**

Due to acute oral and dermal toxicity and avian hazards

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

**THIMET® 15-G**

soil and systemic insecticide

Active Ingredient: Phorate: O,O-diethyl S-[(ethylthio) methyl] phosphorodithioate.....	15.0
Inert Ingredients.....	85.0%
Total.....	100.0%

EPA Reg. No. 241-145

**KEEP OUT OF REACH OF CHILDREN**

**DANGER/PELIGRO!**

**POISON**

**PRECAUCION AL USUARIO:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**FIRST AID**

If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get medical attention.

If on skin, wash thoroughly with soap and water. Remove contaminated clothing and shoes. Wash clothing and decontaminate shoes before reuse.

If in eyes, immediately flush eyes with plenty of water. Get medical attention.

Antidote: Atropine is an antidote.

**ACCEPTED**  
with **COMMENTS**  
in EPA Letter Dated:

**NOV 19 1992**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

241-145

4920

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, nausea, vomiting, diarrhea, and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine.

Use according to label directions.

See Back Panel For Additional Precautionary Statements

IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

SOLD BY  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1990

Net Weight: 50 lbs.  
22.68 kg.

®Registered Trademark of American Cyanamid Company

5720

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER!**

Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

**Do Not Breathe Dust**

Wear a face mask or other respiratory equipment while emptying bags of product into a hopper. While emptying bags into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

**Do Not Contaminate Food or Feed Products**

Once a bag has been opened, use it completely. Make sure the hoppers are emptied while still in the field. Refer to STORAGE AND DISPOSAL statement for further instructions.

Keep all unprotected persons out of operating areas.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Keep Out of Reach of Domestic Animals.

Not for Use or Storage in or Around the Home.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and wildlife. Birds feeding in treated areas may be killed.

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Do not contaminate water when disposing of equipment washwaters.

6920

REENTRY

After dusts have settled, do not enter or allow entry into the treated area until the 48 hour reentry interval has expired, unless the person entering the treated area is wearing the personal protective equipment listed below for early reentry.

For early reentry into treated areas wear a protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical-resistant gloves and chemical resistant shoes, coverings, or boots.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER. Area treated with THIMET 15-G on (date of application). Do not enter without protective clothing.

PERSONAL PROTECTIVE EQUIPMENT

WEAR THE FOLLOWING PROTECTIVE CLOTHING DURING LOADING, APPLICATION, EQUIPMENT REPAIR/CLEANING, AND DISPOSAL OF THE PESTICIDE: A protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical resistant gloves; chemical resistant shoes, shoe covering, or boots.

Unless a chemical resistant protective suit is otherwise required, mixers/loaders must wear a chemical resistant apron and goggles or a face shield.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATE: Long-sleeved shirt and long-legged pants; shoes and socks. Chemical resistant gloves must be available in the cab or cockpit and must be worn when exiting. This clothing is inadequate to protect you during equipment repair or cleaning, reentry, or pesticide disposal work. Do not allow contaminated clothing to come in contact with the insides of the cab.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or using the toilet.

7 9 20

AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear clean clothes. Do not reuse contaminated clothing. Personal clothing worn during work must be laundered separately from household articles. Store protective clothing separately from personal clothing. Clean or launder protective clothing after each use. Clothing and protection equipment heavily contaminated must be destroyed according to state and local regulations. HEAVILY CONTAMINATED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

WORK SAFETY RULE

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposures should be allowed until it has been determined by means of blood tests that the cholinesterase level has returned to normal. Before using this product consult the National Pesticide Telecommunications Network for recommendations regarding such blood tests, poisoning management, emergency treatment, and other information regarding the toxicity of phorate. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7378.

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away, see First Aid section.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

8 9 20

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

#### STORAGE AND DISPOSAL

##### Storage

Store pesticide products in a secure locked area where children, unauthorized persons and animals cannot enter. Do not store in the same area with food or feed. Do not store opened bags.

##### Prohibitions

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Cover or incorporate spills.

##### Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

##### Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

##### General

Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.



9 7 20

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risk referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FROM A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID COMPANY'S LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THIMET 15-G. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

BEFORE USING, READ PRECAUTIONARY STATEMENTS.

THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

10 7 20

CROPS

Crop	Pests Controlled	Rate of THIMET 15-G	Application	Remarks
BEANS At Planting	Mexican bean beetles	6.5-12.5 ozs. per 1,000 ft. of row for any row (minimum 30-inch spacing) or no more than 13.6 lb/A	Drill granules to the side of the seed	Do not place THIMET 15-G granules in direct contact with seed.  Do not graze or feed treated hay or forage to livestock.  Do not apply later than 60 days before harvest.
	Leafhoppers Aphids Lygus bugs Thrips Mites Seedcorn maggots	6.0-9.3 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing) or no more than 10.1 lb/A		
	Seedcorn maggots		Apply granules in a band over the row	

11/9/20

FIELD CORN  
AND SWEET  
CORN  
(excluding  
popcorn)

At Planting

Corn  
rootworms  
Wireworms  
White grubs  
Seedcorn  
maggots  
Seedcorn  
beetles  
Flea beetles  
Mites

8 ozs. per 1,000  
ft. of row for any  
row spacing  
(minimum 30-inch  
row spacing) or no  
more than 8.7 lb/A

Place granules  
in a 7-inch  
band over the  
row directly  
behind the  
planter shoe in  
front of or  
behind the  
press wheel and  
lightly  
incorporate.

Do not use in-  
furrow  
application.  
One additional  
application can  
be applied at  
cultivation.

For field corn:  
Accent\*  
herbicide and  
Beacon\*\*  
herbicide may  
be applied  
after BANDED  
applications of  
THIMET 15-G.

Applications of  
Accent  
herbicide or  
Beacon  
herbicide  
following  
applications of  
THIMET 15-G may  
result in  
temporary crop  
injury.

FIELD CORN  
AND SWEET  
CORN  
(excluding  
popcorn)

At  
Cultivation

Corn  
rootworms

Apply granules  
to base of  
plants just  
ahead of  
cultivation  
shovels so as  
to cover  
granules with  
soil.

Do not make any  
applications of  
THIMET 15-G  
after  
cultivation  
treatment.

Do not harvest  
within 30 days  
of last  
treatment.

---

FIELD CORN  
(excluding  
sweet and  
popcorn)

---

Chinch bug  
nymphs  
(Colorado,  
Kansas, and  
Nebraska)

At  
Cultivation

---

Apply granules  
to base of  
plants just  
ahead of  
cultivation  
shovels so as  
to cover  
granules with  
soil.  
Apply as soon  
as chinch bug  
nymphs are  
observed at  
base of plants.  
Granules must  
be placed at  
the infested  
region of the  
plants and  
adequately  
covered with  
soil.

---

Later  
infestations of  
chinch bugs may  
not be  
controlled.

Do not make any  
applications of  
THIMET 15-G  
after  
cultivation  
treatment.

Do not harvest  
within 30 days  
of last  
treatment.

Do not graze or  
cut for forage  
within 30 days  
of last  
treatment.

Do not apply  
more than 2  
applications  
per season.

COTTON

At Planting

Early season control of:  
Thrips  
Mites  
Leaf miners

4.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing) or no more than 4.4 lb/A

Distribute the granules evenly in the furrow at planting time. Soil temperature moisture and tilth should be favorable for germination and emergence.

Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 2.0 lbs. THIMET 15-G per acre for hill-dropped cotton. Do not graze or feed treated hay or forage to livestock.

Crop injury may occur if applications of diuron containing herbicides such as \*Karmex or \*\*\*Direx are made in conjunction with soil-applied organo-phosphate pesticides.

Mites  
Thrips  
Aphids  
Black cutworms  
Leafhoppers  
White flies  
Leaf miners

8.0-12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing) or no more than 10.9 lb/A  
8.0-10 ozs per 1,000 ft. of row for 30-inch row spacing or no more than 10.9 lb/A

Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.

Recommended for areas requiring an extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils. Do not graze or feed treated hay or forage to livestock.

Crop injury may occur if applications of diuron containing herbicides such as Karmex or Direx are made in conjunction with soil-applied organo-phosphate pesticides.

COTTON	Mites from spring to early summer	12.0-16.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing) or no more than 14.5 lb/A	Incorporate granules into the soil as a side-dress application deep enough to avoid disturbance by future cultivations.	Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest. Do not graze or feed treated hay or forage to livestock.
Side-Dressing (Irrigated Cotton Only)		12.0-13.3 ozs. per 1,000 ft. of row for 30 inch row spacing or no more than 14.5 lb/A		Do not apply more than 2 applications per season
				Crop injury may occur if applications of diuron containing herbicides such as Karmex or Direx are made in conjunction with soil-applied organo-phosphate pesticides.

PEANUTS At Planting	Thrips Leafhoppers	7.3 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing) or no more than 9.9 lb/A	Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.	Do not graze or feed treated hay or forage to livestock.  Wait 90 days after treatment before harvesting peanuts.
PEANUTS At Pegging	Southern corn rootworms and Leafhoppers	14.6 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing) or no more than 19.9 lb/A	Distribute the granules as a band over the fruiting zone at pegging time. Work into the top few inches of soil immediately.	Do not graze or feed treated hay or forage to livestock.  Wait 90 days after treatment before harvesting peanuts.  Do not apply more than 2 applications per season.
POTATOES At Planting	Aphids Leafhoppers Leaf miners Psyllids Wireworms Flea beetle larvae and reduction of flea beetle adults.	Light or Sandy Soils 15.0 ozs. per 1,000 ft. of row for any spacing (minimum 32-inch spacing) or no more than 15.3 lb/A  Heavy or Clay Soils 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing) or no more than 23.5 lb/A	Distribute the granules evenly in the furrow or band on each side of the row.	Wait 90 days after treatment before harvesting potatoes.

	Colorado potato beetle (early season control)	Heavy or Clay Soils: 23 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing) or no more than 23.5 lb/A		Do not use for Colorado potato beetle control in the Long Island area of New York and the Del Marva Peninsula.
SORGHUM At Planting	Greenbugs	8.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing) or no more than 8.7 lb/A	Place granules in a 7-inch band over the row OR drill 1 inch below and 2 inches to the side of the seed.	Do not place THIMET 15-G granules in direct contact with seed. A second application of granules may be applied at cultivation later in the season.
SORGHUM At Cultivation	Chinch bug nymphs (Colorado, Kansas, Nebraska)		Apply granules to base of plants just ahead of cultivation shovels so as to cover granules with soil. Apply as soon as chinch bug nymphs are observed at base of plants. Granules must be placed at the infested region of the plants and adequately covered with soil.	Later infestation of chinch bugs may not be controlled.  Do not make any application of THIMET 15-G after cultivation treatment.  Do not feed foliage before grain harvest.  Do not apply within 30 days of harvest.  Do not apply more than 2 applications per season.



**SOYBEANS**

**At Planting**

For the early season control of:  
 Mexican bean beetles  
 Leafhoppers  
 Lygus bugs  
 Seedcorn maggots  
 Thrips  
 Aphids  
 Mites

Seedcorn maggots

12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing) or no more than 13.1 lb/A

Drill granules to the side of the seed.

Place granules in a 7-inch band over the row directly behind the planter shoe and in front of the press wheel.

Do not place THIMET 15-G granules in direct contact with seed. Do not feed the foliage of treated soybeans. Crop injury may result when preplant incorporated or preemergence applications of metribuzin herbicides are used in conjunction with this product or other soil-applied organo-phosphate pesticides.

**SUGAR BEETS**

**At Planting**

Aphids  
 Beet root maggots  
 Leafhoppers  
 Mites  
 Leaf miners

6.0 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch spacing) or no more than 9.8 lb/A

Drill granules to the side of the seed OR place granules in a band over the row.

Do not place THIMET 15-G granules in direct contact with the seed.  
  
 Do not apply within 30 days of harvest.  
  
 Consult your state experiment station or state extension service for proper timing of application in your area.

18 9 20

<p><b>SUGAR BEETS</b>  Post-emergence</p>	<p>Mites Aphids Beet root maggots</p>	<p>6.5-10.0 lbs. per acre</p>	<p>Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.</p>	<p>Do not apply by air.  Do not apply more than 2 applications per season.</p>
<p><b>SUGAR CANE (Florida)</b></p>	<p>Wireworms</p>	<p>26.0 lbs. per acre</p>	<p>Apply in a 10-12 inch band directly on the seed piece and surrounding soil in the open furrow immediately before covering.</p>	<p>Do not graze or feed treated hay or forage to livestock.</p>
<p><b>WHEAT</b>  At Planting</p>	<p>Grasshoppers<sup>1</sup> Aphids (Western and North Central States only) Hessian fly</p>	<p>1.6 ozs. per 1,000 ft. of row for any row spacing (minimum 8-inch spacing) or no more than 6.5 lb/A</p>	<p>Apply the granules in the seed furrow.</p>	<p>Do not feed or graze foliage within 70 days of treatment. Do not make any later applications after planting time treatment. <sup>1</sup>Grasshoppers controlled in winter wheat only. Do not harvest within 70 days of application.</p>

\*Accent and Karmex are registered trademarks of E.I. duPont deNemours and Company

\*\*Beacon is a registered trademark of Ciba Geigy Corporation.

\*\*\*Direx is a registered trademark of Griffin Ag Product Co. Inc.

19 9 20

TYPICAL APPLICATION RATES FOR VARIOUS ROW SPACINGS

Rate Oz./1,000 Ft. of Row

1.6 oz 2 oz 4 oz 6 oz 8 oz 10 oz 12 oz 14 oz 16 oz 18 oz 20 oz 22 oz 24

EQUIVALENT POUNDS PER ACRE

40"			3.3	4.9	6.5	8.2	9.8	11.4	13.1	14.7	16.3	18.0	19
38"			3.4	5.2	6.9	8.6	10.3	12.0	13.7	15.5	17.2	18.9	20
36"			3.6	5.4	7.3	9.1	10.9	12.7	14.5	16.3	18.2	20.0	21
34"			3.8	5.6	7.7	9.7	11.6	13.5	15.4	17.4	19.3	21.3	23
32"			4.1	6.1	8.2	10.2	12.3	14.3	16.3	18.3	20.5	22.4	24
30"			4.4	6.5	8.7	10.9	13.1	15.2	17.4	19.6	21.8	24.0	26
26"			5.0	7.1	9.5	11.8	14.2	16.6	18.9	21.3	23.7	26.0	28
24"			5.4	8.2	10.9	13.6	16.3	19.1	21.8	24.5	27.2	30.0	32
22"			5.9	8.9	11.9	14.9	17.9	21.0	24.0	26.9	29.9	33.0	35
20"			6.5	9.8	13.1	16.3	19.6	22.8	26.1	29.3	32.6	35.9	39
16"			8.2	12.3	16.3	20.5	24.6	28.7	32.7	36.8	40.9		
12"	4.4	5.4	10.9	16.3	21.8	27.2	32.7	38.1	43.6				
8"	6.5	8.2	16.3	24.6	32.7	40.9							
6"	8.7	10.9	21.8	32.7	43.6								

20 9 20

CALIBRATION INFORMATION

FIRST READ THE LABEL

It is important that applicator equipment be properly set to deliver the labeled rate. The table below contains suggested starting gauge settings for calibration at planting speeds of 5 or 7 mph and application rates of 4.5, 6.0 or 9.0 oz. 1,000 ft. of row.

NOTE: The settings in this table SHOULD ONLY BE USED AS STARTING POINTS. Continually check the amount of THIMET 15-G used against a known length of row and make further adjustments accordingly. Also check calibration occasionally to make sure equipment wear, changing moisture conditions, etc. have not caused a change in flow rate.

SUGGESTED GRANULAR APPLICATOR SETTING<sup>1</sup>

Applicator	Planter Speed 5 mph			Planter Speed 7 mph		
	4.5 oz.	6.0 oz.	9.0 oz.	4.5 oz.	6.0 oz.	9.0 oz.
John Deere MaxEmerge 2	14	17	23	17	21	33
John Deere Max-Emerge (Odd Notches 5-15-25-35)	9	11	14	11	13	17
John Deere Max-Emerge (Even Notches 0-10-20-30)	9	13	20	13	19	26
John Deere 71 Flexi-Planter (John Deere Metal Hoppers)	1/23	1/26	2/3	1/26	1/32	2/14
New International Harvester (2 gauges hopper)	1/9	2/2	2/6	2/2	2/5	3/3
Old International Harvester (1 gauge hopper)	1/6	1/8	2/3	1/9	2/2	2/9
New Gandy	15	18	22	18	21	25
New Noble	6	8	11	8	11	14
Old Noble	8	11	16	12	15	22
Allis Chalmers (78 and 79 Series)	4	5	7	4	5	7
Allis Chalmers #385	10	11	14	11	13	18

<sup>1</sup>Suggested starting point only.