


RESTRICTED USE PESTICIDE

Due to acute oral and dermal toxicity and avian hazards

For retail sale to, and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

EPA Reg. No. 241-145



POSTEMERGENCE SIDE DRESSING AT HILLING IN POTATOES

DIRECTIONS FOR USE

BEFORE USING, READ PRECAUTIONARY STATEMENTS ON BAG/CONTAINER LABEL

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. THIMET 15-G should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows, turns and loading areas by deep discing immediately after treating fields.

| Crop | Pests Controlled | Rate of THIMET 15-G | Application | Remarks |
|---------------------------|--|--|---|---|
| POTATOES Postemergence | Aphids Leafhoppers Wireworms Colorado potato beetle (early season control) | Light or Sandy Soils: 15 oz. per 1,000 ft. of row for any row spacing (32-inch minimum) Do not use on heavy or clay soils. | For postemergence treatment: Place granules on each side of the row at the level of the seed pieces before covering at hilling. Apply within 4 to 6 weeks after planting. | Wait 90 days after treatment before harvesting potatoes. DO NOT apply THIMET as a side-dress treatment if THIMET has been applied at planting time. Do not use for Colorado potato beetle control on the Del Marva Peninsula. |

THIS LABEL SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION

