


Product:

AMINO TRIAZOLE
Weedkiller 90



(red)

To be used only on weeds. Do not use on crops.
Apply in the morning, early afternoon, or at night,
when the temperature is above 50°F.


Active Ingredient:

Amitrole.....90.0%

Inert Ingredients.....10.0%

*9-amino-1,2,4-triazole

EPA Reg. No. 241-168



(19 pt. type)
(12 pt. type)

CAUTION
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed.
Avoid contact with skin, eyes or clothing.

REMARKS

AL. B. 10016: Si usted no lee inglés, no use este producto hasta que le
ofrezcamos la ayuda explicada anteriormente.
(10 pt. type) If you cannot read English, do not use this product until
the label has been fully explained to you.)

(Illustration)

Net weight: 100 lbs.

24703-02

D44

AMINO TRIAZOLE
EPA Reg. No. 241-168

10/12/71

AMINO TRIAZOLE WEEDKILLER 90 is not effective when applied directly to
the foliage of actively growing plants. It is absorbed by the leaves and
travels through the entire plant - down to the deepest roots. Leaves wilt
within 10 days to 3 weeks, and the plants subsequently brown and die out.

AMINO TRIAZOLE WEEDKILLER 90 should be applied in sufficient water to
thoroughly cover the foliage. Treatment may be necessary to control
plants not thoroughly sprayed or plants arising from dormant seeds.

HOW TO MIX AND APPLY

AMINO TRIAZOLE WEEDKILLER 90 is not a wettable powder; it is soluble in
water. This unique feature allows a proper flow through the spray
system without clogging nozzles or corroding spray equipment. When mixing
AMINO TRIAZOLE WEEDKILLER 90 with chlorine, first dilute the AMINO TRIAZOLE
Weedkiller 90 and then add chlorine wettable powder with care during
mixing and spraying operations.

Do not use on crops, lawns, or ornamentals. DRAINAGE DITCHBANKS,
ROADS, AND - DO NOT

For best results, use a backpack sprayer application of AMINO TRIAZOLE
Weedkiller 90 for control of annual and perennial weeds. Mix 1 cup
in the table below, plus 1 cup of AMINO TRIAZOLE WEEDKILLER 90 in
1 gallon of water (1 cup equals 3 level tablespoons). Use the higher
rate for control of perennial weeds. One gallon of solution generally
covers 1,000 sq. ft. of actively growing weeds. In
intermediate situations, or on dryland crops, mix 1/2 cup of this
product in 1 gallon of water to cover 1,000 sq. ft. Apply when weeds are
growing actively. Let all foliage thoroughly. Repeat applications may
be necessary to control dormant perennial weeds.

WOODY PLANTS AND VINES

Weeds	Amount Per Acre*	When to Apply
Poison Ivy Poison Oak	2 lbs. in 100 gals. water	Spray when plants are in full leaf. Spray until plants are thoroughly wet to ground level.
Honeysuckle Kudzu Vine	5 1/2 lbs. in 125 gals. water	Spray in June when leaves are fully developed. If regrowth appears, repeat application in August or following spring. Wet plants thoroughly.

*A strip 8 feet wide and 1 mile long equals approximately 1 acre.

Weeds Amount Per Acre* When to Apply

For control of ANNUAL GRASSES and BROADLEAF WEEDS such as: <u>Foxtail</u> , <u>Rye Grass</u> , <u>Cheatgrass</u> , <u>Barley Grass</u> , <u>Annual Bluegrass</u> , <u>Pigweed</u> , <u>Puncture Vine</u> , <u>Kochia</u> , <u>Sunflower</u>	2 lbs. in 30-60 gals. water	Spray when weeds are 3-6 inches tall. KEEP L ^{IVESTOCK} OFF TREATED AREAS.
For control of PERENNIAL BROADLEAF WEEDS such as: <u>White Top</u> , <u>Canada Thistle</u> , <u>Milkweed</u> , <u>Sow Thistle</u> For suppression of PERENNIAL GRASSES such as: <u>Quack Grass</u> , <u>Rip-gut Grass</u> , <u>Bermuda Grass</u> , <u>Reed Canary Grass</u> , <u>Wild Barley</u>	4 lbs. in 30-60 gals. water	When most weeds are 6-10 inches tall. Repeat if needed. KEEP LIVESTOCK OFF TREATED AREAS.
	7 to 11 lbs. in 300 to 400 gals. water for ground spray; 15 gals. water for aerial spray.	Spray when catkins are fully formed and up to 3 weeks before expected frost. Do not disturb plants after treatment. Spray until plants are wet. Do not concentrate water near top of plants.

Side 2

HARDWOOD NURSERIES

Weeds	Amount per Acre	When to Apply
Chrysanthemum	9 lbs. in 100 gals. water	Apply as a directed spray on young weed growth 4 to 6 inches high making sure to keep the spray off the foliage of hardwood nursery stock.

FOR GENERAL WEED CONTROL IN NURSERY STOCK*, ORNAMENTAL
PLANTINGS* AND FOREST PLANTATIONS

AERIO TRIANGLE		Gals.	Area
<u>Weedkiller 90</u>	<u>Simazine (80-1)</u>	<u>Water</u>	
¼ lb.	1 lb.	20	110 x 100
½ lb.	2 lbs.	40	220 x 100
1 lb.	4 lbs.	80	1 acre

Apply mixture to small weeds in the spring or early summer. Weeds should be wet thoroughly. Cut or mow tall weeds to ground line, remove cuttings and spray regrowth 1 or 2 weeks later. Where Johnson grass, Bermuda grass or other weeds are not controlled by the first application, treat again with the same rate when regrowth is 6 to 12 inches tall (about 2 months).

*The following plant species may be treated. Use a directed spray.
Balsam fir, Douglas fir, Fraser fir, Norway spruce, Red spruce, White spruce, Red pine, Scotch pine, Arborvitae (Thuja), Juniper, Yew (Taxus), Oleander, Pyracantha and Viburnum.

Camera's Note

Poor Copy

CAUTION: Lilac, privet, honeysuckle and barberry are susceptible to this mixture and may be injured if contacted by spray.

DISCLAIMER

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable by American Cyanamid Company.

DISPOSAL OF DRUMS: Destroy empty drums by burning. Stay away from smoke and fumes.

Lot No.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N.J. 08540