

Face

(Illustration)



6-24-24 Fertilizer + THIME[®] Soil and Systemic Insecticide

(red)

(Contains 0.50 Lb. of phorate per 100 lbs.)

FOR CONTROL OF CORN FOOTWORMS ON CORN



Active Ingredient:

Phorate (O,O-diethyl S-[[(ethylthio)methyl]phosphorodithioate)..... 90.50%

Inert Ingredients..... 9.50%

USDA Reg. No. 141-XXX

5-4-1977

ACCEPTED WITH COMMENTS

(18-pt. type)

DANGER!

(12-pt. type)

KEEP OUT OF REACH OF CHILDREN



(red)

See Side Panel For Antidote and Other Warnings

Net Weight:

Am:gc

2/1/68

5

DANGER!

TOXIC BY SKIN CONTACT, INHALATION, OR SWALLOWING

RAPIDLY ABSORBED THROUGH SKIN

REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS,

PROGRESSIVELY INCREASE SUSCEPTIBILITY TO

POISONING

Do Not Get In Eyes, On Skin, Or Clothing

Wear freshly-laundered, long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. Launder clothing before re-use. Wash thoroughly with soap and water before eating or drinking. Bathe at the end of the work day and change outer clothing.

Do Not Breathe Dust

While emptying cans into equipment, wear downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

Do Not Contaminate Food or Feed Products

Sweep up and bury spillage, whether it occurs in camp or in the field. Once a spill has been cleaned, wash it completely or bury the residue. Be sure that the hoppers of equipment are emptied while in the field.

Keep all unrotected persons out of handling areas.

Destroy empty cans and cartons by burning. Stay away from open fires.

Keep Out of Reach of Children

Do not take to field or wilderness areas any containers or water for supply to any animal or human.

Gusnot



Remove patient from exposure area immediately and wash the skin clean with copious amounts of water. Remove traces of the pesticide. If swallowed, induce vomiting with 1-2 glasses of salty or soapy water. Treat with vomit fluid. Patient should lie down and keep quiet.

Call a doctor immediately for further instructions.

Remove patient from exposure area immediately and wash the skin clean with copious amounts of water. Remove traces of the pesticide. If swallowed, induce vomiting with 1-2 glasses of salty or soapy water. Treat with vomit fluid. Patient should lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO A PERSON WHO IS UNCONSCIOUS.

NOTE TO PHYSICIANS: Warning symptoms include weakness, dizziness, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine intramuscularly or intravenously, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) at once and every hour as required until pupils dilate. Up to 30 milligrams, or more, may be required during the first 24 hours. Never give morphine or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 24 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause increasing susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until cholinesterase regeneration has taken place as determined by blood tests.

ANALYZED FERTILIZER ANALYSIS

Total Nitrogen (N)	6.0%
Available Phosphoric Acid (P ₂ O ₅)	24.0%
Soluble Potash (K ₂ O).....	24.0%

IMPORTANT! Read precautionary statements before using this product.

DISCLAIMER

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid Company.

DIRECTIONS FOR USE

Do not use this product for any uses other than those specified on this label.

TRIFL is an effective soil and systemic insecticide for control of corn rootworms on corn which has been combined with fertilizer for efficient and uniform application. Apply specified dosage rate at planting time in the starter fertilizer.

CCRN: For control of corn rootworms, apply at the rate of 2.0 lbs. (1 lb. actual) per acre in a band on each side of the row. Do not place in direct contact with the seed.

© Trademark

Made and printed in U.S.A.

Lot No.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N. J.



WARNING
KEEP AWAY from FEED or FOOD PRODUCTS

POISON
CAUTION--DO NOT DROP

IF **DON'T** BREATHE FUMES
TOUCH CONTENTS
SWALLOW

This is to certify that the contents of this package are properly described by name and are packed and marked and are in proper condition for transportation according to the Regulations prescribed by the Interstate Commerce Commission.

Shipper's name required hereon for shipments by EXPRESS