

**DOSAGES**

POUR **WARBEX** AT THE RATE OF 1 OUNCE PER 200 POUNDS BODY WEIGHT FROM THE SHOULDER TO THE TAIL HEAD. DO NOT REPEAT TREATMENT. FOLLOW THE SUGGESTED SCHEDULE BELOW:

AV. CATTLE WEIGHT	DOSAGE	ONE GALLON TREATS
200 LBS.	1 OZ.	128 HEAD
400 LBS.	2 OZS.	64 HEAD
600 LBS.	3 OZS.	42 HEAD
800 LBS. & OVER	4 OZS.	32 HEAD


**PRECAUTION**

DO NOT USE ON LACTATING DAIRY COWS OR DRY DAIRY COWS WITHIN 21 DAYS OF FRESHENING; CALVES LESS THAN 3 MONTHS OLD; ANIMALS STRESSED FROM CASTRATION, OVEREXCITEMENT

OR DEHORNING; SICK OR CONVALESCENT ANIMALS. ANIMALS MAY BECOME DEHYDRATED AND UNDER STRESS FOLLOWING SHIPMENT. DO NOT TREAT UNTIL THEY ARE IN GOOD CONDITION.

**WARNING**

ON OCCASION, CATTLE TREATED WITH **WARBEX** MAY SHOW SIGNS OF ORGANOPHOSPHATE POISONING SUCH AS EXCESSIVE SALIVATION, STIFFNESS OF LIMBS AND DYSPNEA. ATROPINE IS ANTIDOTAL. A VETERINARIAN SHOULD BE CONSULTED. **WARBEX** IS A CHOLINESTERASE INHIBITOR. DO NOT USE ANY DRUG, PESTICIDE, INSECTICIDE, OR CHEMICALS HAVING CHOLINESTERASE-INHIBITING ACTIVITY EITHER SIMULTANEOUSLY OR WITHIN A FEW DAYS BEFORE OR AFTER TREATMENT WITH **WARBEX**.



**DO NOT SLAUGHTER WITHIN 35 DAYS AFTER TREATMENT.**

**AGRICULTURAL DIVISION • AMERICAN CYANAMID COMPANY • PRINCETON, NEW JERSEY 08540**

CONTROL NO.

**WARNING  
HARMFUL OR  
OR ABSORBED  
COMBUST**

AVOID CONTACT  
CLOTHING  
WASH THOROUGHLY  
DO NOT CONTACT  
PRODUCTS  
KEEP AWAY FROM  
OPEN FLAME

**FIRST AID  
IF POISONING  
CALL A PHYSICIAN**

**NOTE TO PHYSICIAN  
THIS PRODUCT  
PROLONGED USE  
CAUSE CHOLINESTERASE  
ATROPINE IS ANTIDOTAL**

® TRADEMARK

**WARNING!**

**HARMFUL OR FATAL IF SWALLOWED  
OR ABSORBED THROUGH SKIN**

**COMBUSTIBLE**

AVOID CONTACT WITH SKIN, EYES AND  
CLOTHING

WASH THOROUGHLY AFTER HANDLING  
DO NOT CONTAMINATE FOOD OR FEED  
PRODUCTS

KEEP AWAY FROM HEAT, SPARKS AND  
OPEN FLAME

**FIRST AID:**

**IF POISONING SHOULD OCCUR,  
CALL A PHYSICIAN.**

**NOTE TO PHYSICIANS:**

THIS PRODUCT, UPON REPEATED,  
PROLONGED OR CARELESS USE, MAY  
CAUSE CHOLINESTERASE INHIBITION.  
ATROPINE IS ANTIDOTAL.

\* TRADEMARK

**WARBEX<sup>®</sup>**

**FAMPHUR**

**POUR-ON CATTLE INSECTICIDE 13.2%**

**FOR CONTROL OF CATTLE GRUBS AND LICE INFESTATIONS**

**CYANAMID**

**DIRECTIONS FOR USE**

ONE TREATMENT WITH EASY, READY-TO-USE  
**WARBEX** WILL EFFECTIVELY CONTROL CATTLE  
GRUBS AND REDUCE CATTLE LICE INFESTATIONS.  
FOR GRUB CONTROL, APPLY AS SOON AS POSSIBLE  
AFTER HEEL FLY ACTIVITY CEASES. CATTLE FROM  
OR IN TEXAS, OKLAHOMA, KANSAS, LOUISIANA,  
MISSISSIPPI, ARKANSAS, EASTERN 1/3 OF NEW  
MEXICO, COLORADO AND SOUTHWESTERN 1/2 OF  
MISSOURI SHOULD NOT BE TREATED AFTER  
OCTOBER 1; CATTLE FROM REST OF U.S. AFTER  
NOVEMBER 1. ANIMALS TREATED AFTER THESE  
DATES MAY SUFFER TOXIC SIDE EFFECTS DUE TO  
KILLING OF MIGRATING LARVAE IN TISSUES. THE  
PREPARATION SHOULD NOT BE USED IF THE  
WEATHER CONDITIONS ARE SUCH THAT IF THE  
ANIMALS GET WET, THE EFFECTIVENESS OF THE  
PREPARATION WILL DIMINISH. SWINE SHOULD BE  
ELIMINATED FROM AREA WHERE RUN-OFF OCCURS.

TO PROTECT FISH AND WILDLIFE, DO NOT  
CONTAMINATE STREAMS, LAKES, OR PONDS WITH  
THIS MATERIAL.

PERFORATE OR CRUSH, AND BURY CONTAINER  
AWAY FROM WATER SUPPLIES WHEN EMPTY. NEVER  
RE-USE.

**BUYER ASSUMES THE RISK OF ANY USE CONTRARY  
TO LABEL INSTRUCTIONS, OR UNDER ABNORMAL  
CONDITIONS, OR UNDER CONDITIONS NOT  
REASONABLY FORESEEABLE BY AMERICAN  
CYANAMID COMPANY. ANY DAMAGES ARISING FROM  
THE USE OF THIS MATERIAL SHALL BE LIMITED TO  
DIRECT DAMAGES, AND SHALL NOT INCLUDE  
CONSEQUENTIAL COMMERCIAL DAMAGES SUCH AS  
LOSS OF PROFITS OR VALUES, ETC.**