

50 lb.

XXXXX-XX

FAMIX*

FAMPHUR

CATTLE PREMIX

Medicated

For Use in the Manufacture of Feeds

To Control Cattle Grubs

and Aid in Control of Sucking Lice

USDA Reg. No. 241-X

Trademark

Control No.

D1

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, NEW JERSEY

ACCEPTED
NOV 7 - 1968
UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 241-205 SUBJECT
TO ATTACHED COMMENTS

Pesticidally Active Drug Ingredient:

Famphur (0,0 dimethyl 0-[p-(dimethylsulfamoyl)phenyl] phosphorothioate

33.3%

Inert Ingredient: phosphate, defluorinated

66.7%

Each pound contains 151 grams of famphur

DIRECTIONS

For the control of cattle grubs, feed FAMIX Cattle Premix at a rate to provide 2.3 mg. famphur per pound body weight daily for 10 days in feed. For control of grubs and aid in the control of sucking lice, feed FAMIX Cattle Premix at a rate to provide 1 mg. famphur per pound body weight daily for 30 days in feed.

Mix sufficient FAMIX Cattle Premix thoroughly in feed to supply the appropriate dosage for either 10 or 30 days. Medication should be started as soon as possible after heel fly activity ceases. Cattle from or in Texas, Oklahoma, Kansas, Louisiana, Mississippi, Arkansas, eastern 1/3 of New Mexico, Colorado and southwestern 1/3 of Missouri should not be treated after October 1; cattle from rest of U.S. after November 1. Animals treated after these dates may suffer toxic side effects due to killing of migrating larvae in tissues.

RESTRICTIONS

Do not feed FAMIX to lactating dairy cow or dry dairy cows and heifers within 21 days of freshening. Do not administer to sick or convalescent cattle; or within 4 days of slaughter. Do not treat animals under stress from confinement, overexertion or de-horning. Follow directions carefully. Famphur is a cholinesterase inhibitor. Do not use any drug, insecticide, pesticide or chemical having cholinesterase inhibiting activity either simultaneously or within a few days before or after treatment with FAMIX. Overdosing or improper dosing may result in unusual or abnormal behavior (such as excessive salivation, stiffness of limbs and dyspnea) and may cause marked cholinesterase inhibition. The antidote for FAMIX is atropine.

This preparation and dosage form must be the sole source of famphur.

WARNING:

Hazardous by inhalation, skin contact or swallowing

Repeated inhalation or skin contact may, without symptoms,

progressively increase susceptibility to poisoning.

Do not get in eyes, on skin, on clothing.

Do not breathe dust.

Destroy empty containers by burning. Stay away from smoke and fumes.

FIRST AID:

If poisoning should occur, "call a physician".

Note to Physician:

This product, upon prolonged, repeated, or excessive use, may cause cholinesterase inhibition. The antidote is atropine.