


241-238

12/11/1974

1/5



ACCEPTED
11 DEC 1974

241-238

SOIL INSECTICIDE

FOR USE ON FIELD CORN

CYANAMID

Active Ingredient
 S-(methylthio)methyl 10-dithyl phosphorodithioate 15.0%
 Inert Ingredients 85.0%

EPA Est. No.
 1145-IL-1

DANGER!

KEEP OUT OF REACH OF CHILDREN



EPA Reg. No.
 241-298-AA

See Back Panel For Antidote And Other Warnings
PELIGRO

Atusuario: Si usted no habla inglés, no use este producto hasta que la etiqueta
 le haya sido explicada completamente.
**(TO THE USER: If you cannot read English, do not use this product until the label
 has been fully explained to you.)**

Net Contents 15 Lbs.

24696-01

D40

IF LEAFLET IS NOT ATTACHED, CALL COLLECT 201-835-3100.

CROP	PESTS CONTROLLED	RATE COUNTER 15G	APPLICATION	REMARKS
Field Corn: Application at Planting Time	Corn rootworms	5.0-6.5 lbs. per acre (40-inch row spacing) or 6-8 ozs. per 1000 ft. of row for any row spacing (mini- mum 20-inch row spacing).	Place granules in a 7" band over the row directly behind the planter shoe in front of the press wheel. Or, place directly in the seed furrow behind the planter shoe.	When applied in the seed furrow do not use green fodder or cut for silage prior to 130 days after planting. Do not rotate treated area for 1 year following application except for field corn and soybeans. Cover crops may be planted in treated area if plowed under and not grazed.

TEAR ON DOTTED LINE

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N.J. 08540

9/5

DANGER!

**POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING
RAPIDLY ABSORBED THROUGH SKIN
REPEATED INHALATION OR SKIN CONTACT, MAY, WITHOUT SYMPTOMS,
PROGRESSIVELY INCREASE SUSCEPTIBILITY TO POISONING**

DO NOT GET IN EYES, ON SKIN, ON CLOTHING. Wear freshly laundered, long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. Launder clothing before reuse. Wash thoroughly with soap and water before eating or smoking. Bathe at the end of the work day and change outer clothing.

DO NOT BREATHE DUST. While emptying bags into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone. For further handling instructions see leaflet attached below.

DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. Sweep up and bury spillage, whether it occurs indoors or in the field. Once a bag has been opened, use it completely or bury the remainder. Make sure that the hoppers of equipment are emptied while still in the field. Keep all unprotected persons out of operating areas.

KEEP OUT OF REACH OF DOMESTIC ANIMALS.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

Destroy empty bags and cartons by burning. Stay away from smoke and fumes.

This product is toxic to fish, birds and other wildlife. Treated granules exposed on soil surface may be hazardous to birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DISCLAIMER

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

USER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid Company.

4/5



POISON



IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain (0.6 milligram) tablets for emergency use. If symptoms include blurred vision, stomach cramps or tightness in chest, don't wait for a physician but take two tablets at once. Do not take atropine unless symptoms of poisoning have occurred. Anyone who has been sick enough to have taken atropine must be seen by a physician as soon as possible.

FIRST AID

IF SWALLOWED, induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person.

IF INHALED, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen.

IN CASE OF CONTACT, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.


CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) every 15 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests. Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

© Trademark
Lot No.

Made and Printed in U.S.A.

Sold By
AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N.J. 08540

 ©103
SIFT-GARD®

5/5