


241-238

07/29/1975

1/5




SOIL INSECTICIDE

FOR USE ON FIELD CORN

CYANAMID

DANGER!

KEEP OUT OF REACH OF CHILDREN



POISON

DANGER!

**POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING
RAPIDLY ABSORBED THROUGH SKIN
REPEATED INHALATION OR SKIN CONTACT, MAY, WITHOUT SYMPTOMS,
PROGRESSIVELY INCREASE SUSCEPTIBILITY TO POISONING**

DO NOT GET IN EYES, ON SKIN, ON CLOTHING. Wear freshly laundered, long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. Launder clothing before reuse. Wash thoroughly with soap and water before eating or smoking. Bathe at the end of the work day and change outer clothing.

DO NOT BREATHE DUST. While emptying bags into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone. For further handling instructions see leaflet attached below.


DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. Sweep up and bury spillage, whether it occurs indoors or in the field. Once a bag has been opened, use it completely or bury the remainder. Make sure that the hoppers of equipment are emptied while still in the field. Keep all unprotected persons out of operating areas.

KEEP OUT OF REACH OF DOMESTIC ANIMALS.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

Destroy empty bags and cartons by burning. Stay away from smoke and fumes.

This product is toxic to fish, birds and other wildlife. Treated granules exposed on soil surface may be hazardous to birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.





POISON



IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

ANTIDOTE: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain (0.6 milligram) tablets for emergency use. If symptoms include blurred vision, stomach cramps or tightness in chest, don't wait for a physician but take two tablets at once. Do not take atropine unless symptoms of poisoning have occurred. Anyone who has been sick enough to have taken atropine must be seen by a physician as soon as possible.

FIRST AID

IF SWALLOWED, induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Never give anything by mouth to an unconscious person.

IF INHALED, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen.

IN CASE OF CONTACT, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING


NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests. Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

© Trademark
Lot No.

Sold By

Made and Printed in U.S.A.

**AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N.J. 08540**



475

DISCLAIMER

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.


American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

USER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by American Cyanamid Company.

DIRECTIONS FOR USE

COUNTER 15G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated row and turns and loading areas by deep discing immediately after treating fields.

Crops	Pests Controlled	Rate—COUNTER 15G	Application	Remarks
Field Corn At Planting	Corn rootworms Seed corn maggots	Banded or In-Furrow 6.5 lbs. per acre (40-inch row spacing) or 8 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing).	Banded For corn rootworms place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel. In-Furrow For seed corn maggots and/or corn rootworms place granules directly in the seed furrow behind the planter shoe.	In-Furrow Applications Only Do not feed green fodder or use for silage prior to 130 days after planting. Dry fodder may be fed except where corn has been furrow irrigated. <i>These restrictions do not apply to band applications.</i> All Applications Treated area may be rotated to corn or soybeans without restriction. Do not rotate treated area for 365 days to any other crops. Cover crops may be planted in treated area if not grazed and plowed under.
	Early season control of adult southern corn billbugs (Southeastern states)	Banded 6.5-13.0 lbs. per acre (40-inch row spacing) or 8-16 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing).	Banded Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel.	
	Reduction of corn wireworms	In-Furrow 6.5 lbs. per acre (40-inch row spacing) or 8 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing).	In-Furrow Place granules directly in the seed furrow behind the planter shoe.	



5/5