



241-238

COUNTER® 15-G Soil Insecticide

EPA Reg. No. 241-238-AA

DIRECTIONS FOR USE

BEFORE USING, READ PRECAUTIONARY STATEMENTS ON BAG
 COUNTER 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated row and turns and loading areas by deep discing immediately after treating fields.

Crops	Pests Controlled	Rates COUNTER 15-G	Application
FIELD CORN, POPCORN AND SWEET CORN	Corn rootworms Seedcorn maggots	Banded or In-Furrow 6.5 lbs. per acre (40-inch row spacing) or 8 ozs. per 1000 ft. of row for any row spacing (minimum 20-inch row spacing).	Banded For corn rootworms place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel. In-Furrow For seedcorn maggots and/or corn root- worms place granules directly in the seed furrow behind the planter shoe.
		Banded 6.5-13.0 lbs. per acre (40-inch row spacing) or 8-16 ozs. per 1000 ft. or row for any row spacing (minimum 20-inch row spacing).	Banded Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel.
At Planting	Early season control of adult southern corn billbugs* Reduction of corn wireworms *(Southeastern States)	In-Furrow 6.5 lbs. per acre (40-inch row spacing) or 8 ozs. per 1000 ft. of row for any row spacing (minimum 20-inch row spacing).	In-Furrow Place granules directly in the seed furrow behind the planter shoe.

05/04/1976

APRIL 1976

D41

1/1