

4-26-1979
McLendon 1/4

CONFIDENTIAL - SECURITY INFORMATION

FOR USE ON FIELD CORN, POPCORN, SWEET CORN AND SUGAR BEETS



ORGANIC PHOSPHATE COMPOUND MIXTURE DRY

KEEP OUT OF REACH OF CHILDREN



040

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-238

2/4

DANGER!

MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN
RAPIDLY ABSORBED THROUGH SKIN
REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS,
PROGRESSIVELY INCREASE SUSCEPTIBILITY TO POISONING

DO NOT GET IN EYES, ON SKIN, ON CLOTHING. Wear freshly-laundered, long sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently.
In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. Launder clothing before reuse. Wash thoroughly with soap and water before eating or smoking. Bathe at the end of the work day and change outer clothing.
DO NOT BREATHE DUST. While emptying bags into equipment, pour downward and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.
DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. Sweep up and bury spillage, whether it occurs indoors or in the field. Once a bag has been opened, use it completely or bury the remainder. Make sure that the hoppers of equipment are emptied while still in the field.
Keep all unprotected persons out of operating areas.
KEEP OUT OF REACH OF DOMESTIC ANIMALS.
Destroy empty bags and cartons by burning. Stay away from smoke and fumes.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

This product is toxic to fish, birds and other wildlife. Treated granules exposed on soil surface may be hazardous to birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

DIRECTIONS FOR USE

COUNTER 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated row and turns and leading areas by deep disking immediately after treating fields.

Crops	Pests Controlled	Rate of COUNTER 15-G (See chart below for rate/acre conversion)	Application
FIELD CORN, POPCORN AND SWEET CORN	Corn rootworms Seed corn maggots	Banded or In-Furrow 8 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing).	Banded For corn rootworms place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel. In-Furrow For seed corn maggots and/or corn rootworms place granules directly in the seed furrow behind the planter shoe.
	At Planting Early season control of adult southern corn billbugs* Reduction of corn wireworms *(Southeastern States)	Banded 8-16 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing). In-Furrow 8 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing).	Banded Place granules in a 7-inch band over the row directly behind the planter shoe in front of the press wheel. In-Furrow Place granules directly in the seed furrow behind the planter shoe.
SUGAR BEETS	Sugar beet root maggot	4.5 to 9.0 ozs. per 1,000 feet of row for any row spacing (minimum 20-inch row spacing).	At Planting Apply in a 5 to 7-inch band over the row and lightly incorporate into the soil. Do not place granules in direct contact with the seed. Power incorporation if it is used. Do not incorporate deeper than 2 inches. Post Emergence Apply in a 5 to 7-inch band over the row at the time of first fly emergence and lightly incorporate into the soil.

Typical Application Rates for Various Row Spacings

Rate — OZ./1,000 FT. of Row

	4 in.	6 in.	8 in.	10 in.	12 in.	14 in.	16 in.
40"		4.9	6.5	8.2	9.8	11.4	13.1
38"		5.2	6.9	8.6	10.3	12.0	13.7
36"		5.4	7.3	9.1	10.9	12.7	14.5
34"		5.6	7.7	9.7	11.6	13.5	15.4
32"		6.1	8.2	10.2	12.3	14.3	16.3
30"		6.5	8.7	10.9	13.1	15.2	17.4
28"		7.1	9.5	11.8	14.2	16.6	18.9
26"	8.4	9.2	10.9	13.6	16.3	19.1	21.8
24"	8.9	9.9	11.9	14.9	17.9	21.0	24.0
20"	9.5	10.6	13.1	16.3	19.6	22.8	26.1

EQUIVALENT POUNDS PER ACRE

COUNTER 15-G
SOIL INSECTICIDE

3/4 C



POISON



In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201 - 835-3100.

Antidote: Atropine is an antidote. Consult your physician about obtaining a supply of 1/100 grain (0.6 milligram) tablets for emergency use. If symptoms include blurred vision, stomach cramps or tightness in chest, don't wait for a physician but take two tablets at once. Do not take atropine unless symptoms of poisoning have occurred. Anyone who has been sick enough to have taken atropine must be seen by a physician as soon as possible.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING

FIRST AID

If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams (3 to 6 tablets, 1/100 grain each) every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests. Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine. Use according to label directions.

© Trademark


Made and Printed in U.S.A.

LOT NO: IMPRINTED ON BAG

Sold By
AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, N. J. 08540

SO

FOR USE ON FIELD



Active Ingredient:
Terbufos (S)
Inert Ingredients

EPA Reg. No.
241-238-AA

See B

AL USU
le hoyo
(TO THE
box bag

Net Weight: 20
9.0

4/4

COUNTER® 15-G Soil Insecticide

EPA Reg. No. 241-238-AA

DIRECTIONS FOR USE IN FIELD CORN

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BEFORE USING, READ THE PRECAUTIONARY STATEMENTS ON BAG.

COUNTER 15-G should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage.

Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep disking immediately after treating fields.

Crops	Pests Controlled	Rates COUNTER 15-G	Application
FIELD CORN At Planting	Nematodes:	BANDED 8-16 oz. per 1000 ft. of row for any row spacing (minimum 20-inch row spacing).	BANDED Place granules in a 7-inch band directly behind the planter shoe in front of the press wheel.
	Lesion Spiral Stunt Sting Stubby-root Dagger	IN-FURROW 8 oz. per 1000 ft. of row for any row spacing (minimum 20-inch row spacing.)	IN-FURROW Place granules directly into the seed furrow behind the planter shoe.

AMERICAN CYANAMID COMPANY
AGRICULTURAL DIVISION
PRINCETON, NEW JERSEY 08540

4/79