

COUNTER[®] 15-G

soil insecticide-nematicide

Supplemental Labeling

(EPA Reg. No. 241-238)

DIRECTIONS FOR USE

BEFORE USING, READ PRECAUTIONARY
STATEMENTS ON BAG

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. COUNTER 15-G should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

Crop	Pest Controlled	Rates	Application	Remarks
Grain Sorghum At bedding or At planting	Corn leaf aphid	Knifed-in 8-16 oz. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing) or no more than 26 lbs. per acre.	Knifed-in Drill granules 1-4 inches directly below the seed OR 1-4 inches below the seed and up to 5 inches to the side of the seed.	Do not place granules in direct contact with seed as crop injury may occur. Only one application per year may be used.
Grain Sorghum At planting	Corn leaf aphid	Banded 8-16 oz. per 1,000 ft. of row for any row spacing (minimum 20-inch row spacing)	Banded Place granules in a 5-7 inch band directly behind the planter shoe in front of the press wheel.	Do not use band applications in New Mexico, West Texas and the Panhandle of Oklahoma.

*Registered Trademark of American Cyanamid Company



American Cyanamid Company
Agricultural Division
Wayne, N.J. 07470

