

(EPA Reg. No. 241-238)

ACCEPTED

## DIRECTIONS FOR USE

BEFORE USING, READ PRECAUTIONARY  
STATEMENTS ON BAG

MAK'S 1057  
This product has been found to be  
effective for the control of  
insects registered under  
EPA Reg. No. 241-238

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. COUNTER should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

|                                      |                                                                                     |                                                                                                                                          |                                                                                                                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|--------------------------------------|-------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| SUGAR<br>BEETS<br>At Planting        | Sugar beet<br>root maggots (1)<br>Wireworms (1)<br>White grubs (1)                  | Modified in-furrow or<br>Banded<br>4-8 oz. per 1,000 ft. of row<br>for any row spacing<br>(minimum 20-inch row<br>spacing).              | Modified in-furrow<br>Apply in-furrow at<br>planting time 2-3 inches<br>behind the seed drop<br>zone after some soil has<br>covered the seed. | Only one application per<br>year may be made.<br><br>At Planting<br>Do not place granules in<br>direct contact with the seed<br>as crop injury may occur.<br>Power incorporation may be<br>used. Do not incorporate<br>deeper than 2 inches.<br><br>Postemergence<br>Apply at the time of first fly<br>emergence. Lightly<br>incorporate into the soil.<br><br>If especially heavy<br>infestations are expected<br>use the 8 oz. rate. |
| SUGAR<br>BEETS<br>Post-<br>emergence | Sugar beet<br>root maggots                                                          | Banded<br>4-8 oz. per 1,000 ft. of row<br>for any row spacing<br>(minimum 20-inch row<br>spacing).                                       | Banded<br>Apply in a 5 to 7-inch<br>band over the row and<br>lightly incorporate into<br>the soil.                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| SUGAR<br>BEETS<br>At Planting        | Suppression of:<br>Cutworms                                                         | Banded<br>8 oz. per 1,000 ft. of row for<br>any row spacing (minimum<br>20-inch row spacing).                                            |                                                                                                                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|                                      | Suppression of:<br>Sugar Beet<br>cyst nematode<br>Control of:<br>Beet<br>Leafhopper | Knifed-in<br>18 oz. per 1,000 ft. of row<br>for any row spacing<br>(minimum 20-inch row<br>spacing) or no more than<br>29 lbs. per acre. | Knifed-in<br>Omit granules 2 inches to<br>the side of the seed and<br>2-4 inches below the<br>seed.                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                        |

THIS LABEL SHOULD BE IN THE POSSESSION OF THE  
USER AT THE TIME OF PESTICIDE APPLICATION.



**CYANAMID**

Agricultural Division  
Crop Protection Chemicals Department  
Wayne, New Jersey 07470

(EPA Reg. No. 241-238)

**ACCEPTED**

MAR 3 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-238

## DIRECTIONS FOR USE

BEFORE USING, READ PRECAUTIONARY  
STATEMENTS ON BAG

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. COUNTER should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.

| Crops                               | Pests Controlled                                                                                                                                                                   | Rates of COUNTER                                                                                           | Application                                                                                                                                                                                                   | Remarks                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|-------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| FIELD CORN, POPCORN, AND SWEET CORN | Corn rootworms<br>Wireworms<br>White grubs<br>Seedcorn maggot<br>Seedcorn beetles<br>Corn flea beetles<br>Maize billbugs<br>Southern corn billbugs                                 | Banded or In-Furrow<br>8 oz. per 1,000 ft. of row for any row spacing. Do Not exceed 17.4 pounds per acre. | Banded<br>Place granules in a 7-inch band over the row, in front of or behind the presswheel and lightly incorporate.<br><br>In-Furrow<br>Place granules directly in the seed furrow behind the planter shoe. | For use on conventional and conservation tillage corn. In situations where crop debris or winds can prevent proper placement of granules, in-furrow applications are recommended. In-furrow applications reduce the potential for insecticide runoff resulting from excessive rain.<br><br>Under dry soil conditions or heavy infestations, it may be necessary to apply an insecticide rescue treatment with another registered insecticide after corn emergence to control surviving cutworm larvae.<br><br>If especially heavy infestations are expected, banded applications of up to 16 oz. per 1,000 ft. of row may be used. |
| At Planting                         | Thrips<br>Symphylans<br>Nematodes:<br>Lance<br>Lesion<br>Root knot<br>Spiral<br>Stunt<br>Sting<br>Stubby root<br>Dagger<br>Suppression of:<br>Cutworms<br>Lesser corn-stalk borers |                                                                                                            |                                                                                                                                                                                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |

 **CYANAMID**  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, New Jersey 07470