

100-25

241-243

use & directions

106

STOMP® herbicide

Active Ingredient:

pendimethalin (N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine) 42.3%

Inert Ingredients 57.7%

(1 gal. contains 4 lbs. of pendimethalin)

**FOR CONTROL OF MOST ANNUAL GRASSES
AND CERTAIN BROADLEAF WEEDS IN NONCROPLAND AREAS**

EPA Reg. No. 241

AVISO!

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(12 pt. type)


KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

(18 pt. type)

WARNING!



CAUSES EYE IRRITATION

HARMFUL IF SWALLOWED, ABSORBED THROUGH THE SKIN, OR INHALED.

Avoid contact with eyes, skin, and clothing. Wear goggles or face shield. Avoid breathing spray mist. Wash skin thoroughly with soap and water after handling. Use with adequate ventilation. Keep container closed. Remove contaminated clothing and wash before reuse.

FIRST AID: If swallowed, DO NOT induce vomiting. Call a physician or Poison Control Center immediately. In case of eye contact, immediately flush eyes with plenty of water. For skin contact, wash thoroughly with soap and water. Call a physician if eye or skin irritation persists.

This product contains monochlorobenzene.

NOTE TO PHYSICIAN: Because of increased risk of chemical pneumonia or pulmonary edema caused by aspiration of the hydrocarbon solvent (monochlorobenzene), vomiting should be induced only under professional supervision.

(2/6/85)

•Registered trademark of American Cyanamid Company

6

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, and ponds. DO NOT apply when weather conditions favor drift from target area. DO NOT contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL HAZARDS

FLAMMABLE LIQUID AND VAPOR

DO NOT use, pour, spill, or store near heat or open flame. DO NOT use in the vicinity of pilot lights.


DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY.



3/16

(12 pt. type)

STORAGE AND DISPOSAL

STORAGE: DO NOT STORE BELOW 40°F. Extended storage at temperatures below 40°F can result in the formation of crystals on the bottom of the container. If crystallization does occur, store the container on its side at room temperature (70°F) and rock occasionally until crystals redissolve.


DO NOT contaminate water, food, or feed by storage or disposal.

DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER

DISPOSAL: Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.



DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of pesticide application.

DO NOT use in California.

Observe all cautions and limitations in this leaflet and the labels of products used in combination with STOMP. The use of STOMP not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

General Worker Protection Statements

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **WARNING.** Area treated with STOMP on (date of application). Do not enter without appropriate protective clothing.

No. 1000
10/1/68



GENERAL INFORMATION

STOMP is recommended for preemergence control of most annual grasses and certain broadleaf weeds as they germinate on noncropland areas such as railroad, utility and pipeline rights-of-way, utility substations, petroleum tank farms, pumping installations, fence rows, storage areas, and other similar areas. STOMP treatments are most effective in controlling weeds when adequate rainfall is received within 7 days after application.

STOMP will not control established weeds. Therefore, areas to be treated should be free of established weeds at the time of treatment, or STOMP may be used in conjunction with herbicides registered for postemergence use in noncropland areas. Consult the labels of those herbicides for suggested treatments, rates to be used and precautions or restrictions for use in noncropland areas.

MIXING INSTRUCTIONS

1. Fill tank one-half to three-quarters full with clean water.
2. Add STOMP to the partially filled tank while agitating and then fill the remainder of the tank with water.
3. MAINTAIN CONTINUOUS AGITATION WHILE ADDING STOMP AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.
4. If STOMP is to be used in tank mixtures with other registered herbicides, then follow directions on the labels of those products which recommend tank mixing.

SPRAYING INSTRUCTIONS

Uniformly apply with properly calibrated ground equipment in sufficient water per acre to uniformly treat the area with a spray pressure of 25 to 50 psi. Maintain continuous agitation during spraying with good mechanical or bypass agitation.

WEED SPECIES CONTROLLED

The following grass and broadleaf weed species are controlled by preemergence treatments of STOMP at the rates recommended below:

GRASSES CONTROLLED

- Barnyardgrass (Echinochloa crus-galli)
- Bluegrass (annual) (Poa annua)
- Browntop panicum (Panicum fasciculation)
- Crabgrass (Digitaria spp.)
- Fall panicum (Panicum dichotomiflorum)
- Foxtail (giant) (Setaria faberi)
- Foxtail (green) (Setaria viridis)
- Foxtail (yellow) (Setaria glauca)
- Goosegrass (Eleusine indica)
- Itchgrass (Rottboellia exaltata)
- Johnsongrass (from seed) (Sorghum halepense)
- Lovegrass (Eragrostis spp.)
- Signalgrass (Brachiaria platyphylla)
- Texas panicum (Panicum texanum)
- Witchgrass (Panicum capillare)
- Woolly cupgrass (Eriochloa villosa)

BROADLEAF WEEDS CONTROLLED

- Carpetweed (Mollugo verticillata)
- Chickweed (common) (Stellaria media)
- Fiddleneck (Armsinckia intermedia)
- Filaree (Erodium spp.)
- Florida pusley (Richardia scabra)
- Henbit (Lamium amplexicaule)
- Knotweed (prostrate) (Polygonum aviculare)
- Lambsquarters (Chenopodium album)
- London rocket (Sisymbrium irio)
- Pigweed (Amaranthus spp.)
- Purslane (Portulaca oleracea)
- Sheperdspurse (Capsella bursa-pastoris)
- Smartweed (Polygonum pensylvanicum)
- Spurge, annual (Euphorbia spp.)
- Velvetleaf (Buttonweed) (Abutilon theophrasti)

RECOMMENDED RATES

Short-term control (4 months) : 2 quarts STOMP per acre
 Long-term control (6-8 months): 4 quarts STOMP per acre

American Cyanamid Company
 Agricultural Division
 Wayne, NJ 07470

