


EPA Reg. No. 241-243

FOR USE IN SUNFLOWERS

FALL APPLICATION OF PROWL¹ OR PROWL PLUS EPTAM² TANK MIXTURE



OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL LEAFLET
LABEL BEFORE USING

GENERAL INFORMATION

PROWL or PROWL plus Eptam 7E tank mix may be applied and incorporated in late fall prior to planting sunflowers the following spring.

Destroy existing weeds before applying PROWL or PROWL plus Eptam 7E. The soil should be dry enough to permit good incorporation.

DIRECTIONS FOR USE

Uniformly apply PROWL or PROWL plus Eptam 7E in 10 or more gallons of water or in 20 or more gallons of liquid fertilizer per acre by ground equipment. PROWL alone may be applied in 5 or more gallons of water per acre by aircraft, but PROWL plus Eptam tank mixtures may only be applied by ground equipment.

Application should be made in the late fall when soil temperature are 45°F or below but before the ground freezes. DO NOT apply PROWL or PROWL plus Eptam 7E tank mixes when the air temperature is below 45°F.

¹Registered Trademark of American Cyanamid Company

²Eptam is a trademark of Stauffer Chemical Company

PROWL or PROWL plus Eptam 7E tank mixture should be incorporated immediately after application, following the instructions for incorporation given in the PROWL and Eptam leaflet labels.

Prior to sunflower planting in the spring, fields treated with PROWL or PROWL plus Eptam 7E should receive at least one shallow additional incorporation. Spring incorporation should be at an angle to the last tillage operation.

Refer to the PROWL leaflet label for specific directions regarding mixing, spraying, and liquid fertilizer compatibility determinations.

PROWL PREPLANT INCORPORATED APPLICATION

WEED SPECIES CONTROLLED


When applied as directed, PROWL will control the following weed species:

GRASSES:

Barnyardgrass (Echinochloa crus-galli)
Crabgrass (Digitaria spp.)
Fall panicum (Panicum dichotomiflorum)
Field sandbur (Cenchrus incertus)
Foxtail (giant) (Setaria faberi)
Foxtail (green) (Setaria viridis)
Foxtail (yellow) (Setaria glauca)
Goosegrass (Eleusine indica)
Witchgrass (Panicum capillare)

BROADLEAF WEEDS:

Carpetweed (Mollugo verticillata)
Florida pusley (Richardia scabra)
Kochia (Kochia scoparia)
Lambsquarters (Chenopodium album)
Pigweed (Amaranthus spp.)
Purslane (Portulaca oleracea)
Spurge, annual (Euphorbia spp.)



PROWL will aid in the control of and reduce competition from:

Pennsylvania smartweed (Polygonum pensylvanicum)

BROADCAST RATES PER ACRE FOR PROWL
PREPLANT INCORPORATED APPLICATION¹


	PROWL (pints)	PROWL (pints)
SOIL TEXTURE	UP TO 3% ORGANIC MATTER	MORE THAN 3% ORGANIC MATTER
COARSE	1 1/2 to 2 1/2	2 1/2
MEDIUM	2 to 3	3 to 3 1/2
FINE	2 1/2 to 3 1/2	3 1/2

¹For use in North Dakota, South Dakota, and Minnesota only.

DO NOT USE on peat or muck soils.

Use the high rate for each soil texture if heavy weed pressure is anticipated.

NOTE: DO NOT feed forage or graze livestock in treated sunflower fields. If sunflower crop loss occurs due to weather conditions, sunflowers or any crop registered for PROWL preplant incorporated use can be replanted. If replanting is necessary, DO NOT rework the soil deeper than the treated zone.



PROWL PLUS EPTAM PREPLANT INCORPORATED TANK MIXTURES

WEEDS CONTROLLED

Tank mixtures of PROWL plus Eptam will control the following weeds in addition to those controlled by applications of PROWL alone:

GRASSES:

Annual bluegrass (Poa annua)
Lovegrass (Stinkgrass) (Eragrostis cilianensis)
Wild oats (Avena fatua)

BROADLEAF WEEDS:

Annual Morningglory (Ipomoea purpurea)
Black Nightshade (Solanum nigrum)
Corn spurry (Spergula arvensis)
Hairy Nightshade (Solanum villosum)
Nettleleaf goosefoot (Chenopodium murale)

545

**BROADCAST RATES PER ACRE FOR PROWL PLUS EPTAM 7E PREPLANT
INCORPORATED TANK MIXTURES¹**

	PROWL + EPTAM 7E (pints)	PROWL + EPTAM 7E (pints)
SOIL TEXTURE	UP TO 3% ORGANIC MATTER	MORE THAN 3% ORGANIC MATTER
COARSE	1 to 2 + 2 1/4	2 + 2 1/4
MEDIUM	1 1/2 to 2 1/2 + 2 1/4	2 1/2 to 3 + 2 1/4
FINE	2 to 3 + 2 1/4	3 + 2 1/4

¹For use in North Dakota or Minnesota only.

DO NOT USE on peat or muck soils.

Use the high rate for each soil texture if heavy weed pressure is anticipated.

NOTE: DO NOT feed forage or graze livestock in treated sunflower fields. If sunflower crop loss occurs due to weather conditions, sunflowers or any crop registered for PROWL + Eptam preplant incorporated use can be replanted. If replanting is necessary, DO NOT rework the soil deeper than the treated zone. Observe all precautions and limitations on the Eptam 7E label.

