

## FOR USE IN FIELD CORN AND SWEET CORN

PROWL PLUS BICEP<sup>1</sup> TANK MIXTURE

PROWL may be applied in sweet corn (processing varieties) in Illinois, Minnesota, New York and Wisconsin ONLY.

Observe all Precautionary Statements in the PROWL and Bicep leaflet labels before using.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Uniformly apply PROWL plus Bicep with properly calibrated ground equipment in 10 to 40 gallons of water or 20 or more gallons of liquid fertilizer per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. Preemergence applications in field corn may be made through center pivot irrigation systems. CHEMIGATION: Refer to supplemental labeling entitled, PROWL and Bicep leaflet labels for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

See the PROWL leaflet label for Mixing Instructions and Liquid Fertilizer Compatibility Determinations. Use the teaspoons of herbicide specified for Cotoran<sup>1</sup> when determining liquid fertilizer compatibility with Bicep. Add PROWL to the spray tank prior to the addition of Bicep.

DO NOT apply on peat or muck soils.

DO NOT use in California.

## PREEMERGENCE APPLICATIONS

Apply preemergence (surface treatment) after planting, but before weeds and crop emerge. Use PROWL and Bicep rates listed in the table on page 2.

DO NOT INCORPORATE or serious corn injury can result.

## EARLY POSTEMERGENCE APPLICATIONS

Apply after corn emergence but no later than when corn is in the 4-leaf stage of growth and when weeds are no larger than the 1 - 2 leaf growth stage. Use PROWL rates listed in the following table with the Bicep rates listed under the "More than 3% Organic Matter" section of the table. EMERGED WEEDS WILL NOT BE CONTROLLED BY THIS TANK MIX. Destroy emerged weeds by cultivation prior to application.

DO NOT INCORPORATE.

<sup>1</sup> Bicep and Cotoran are trademarks of Ciba-Geigy Corp.

© Registered Trademark of American Cyanamid Company

DO NOT apply postemergence in liquid fertilizer.

DO NOT apply following PROWL preemergence treatments.

## WEED SPECIES CONTROLLED

When applied as directed, PROWL plus Bicep tank mix will control the following grass and broadleaf weed species:

## GRASSES CONTROLLED

Barnyardgrass (*Echinochloa crus-galli*)

Browntop panicum (*Panicum fasciculatum*)

Crabgrass (*Digitaria* spp.)

Crowfootgrass (*Dactyloctenium aegyptium*)

Field sandbur (*Cenchrus incertus*)

Foxtails

giant (*Setaria faberi*)

green (*Setaria viridis*)

yellow (*Setaria glauca*)

Goosegrass (*Eleusine indica*)

Johnsongrass (from seed) (*Sorghum halepense*)

Red rice (*Oryza sativa*)

Panicums

fall (*Panicum dichotomiflorum*)

Texas (*Panicum texanum*)

Signalgrass (*Brachiaria platyphylla*)

Southwestern cupgrass (*Brachiaria platyphylla*)

Witchgrass (*Panicum capillare*)

Woolly cupgrass (*Eriochloa villosa*)

Yellow nutsedge (*Cyperus esculentus*)

## BROADLEAF WEEDS CONTROLLED

Black nightshade (*Solanum nigrum*)

Carpenterweed (*Mollugo verticillata*)

Cocklebur (*Xanthium strumarium*)

Florida pusley (*Richardia scabra*)

Galinsoga (*Galinsoga* spp.)

Hairy nightshade (*Solanum sarrachoides*)

Jimsonweed (*Datura stramonium*)

Kochia (*Kochia scoparia*)

Lambsquarters (*Chenopodium album*)

Morningglory (*Ipomea* spp.)

Pigweed (*Amaranthus* spp.)

Purslane (*Portulaca oleracea*)

in the control of and reduce competition from:  
 Shattercane (*Sorghum bicolor*)  
 Sicklepod (*Cassia obtusifolia*)  
 Volunteer sorghum (*Sorghum vulgare*)  
 Wild proso millet (*Panicum miliaceum*)

When applied early postemergence as directed, PROWL plus Bicep tank mix will control the following grass and broadleaf weed species:

**GRASSES CONTROLLED**

Barnyardgrass (*Echinochloa crus-galli*)  
 Crabgrass (*Digitaria* spp.)  
 Crowfootgrass (*Dactyloctenium aegyptium*)  
 Field sandbur (*Cenchrus incertus*)  
 Foxtails  
 giant (*Setaria faberi*)  
 green (*Setaria viridis*)  
 yellow (*Setaria glauca*)  
 Goosegrass (*Eleusine indica*)  
 Johnsongrass (from seed) (*Sorghum halepense*)

Carpetweed (*Mollugo verticillata*)  
 Florida pusley (*Richardia scabra*)  
 Jimsonweed (*Datura stramonium*)  
 Kochia (*Kochia scoparia*)  
 Lambsquarters (*Chenopodium album*)  
 Mustard (*Brassica* spp.)  
 Pigweed (*Amaranthus* spp.)  
 Prickly sida (*Sida spirosoa*)  
 Purslane (*Portulaca oleracea*)  
 Ragweed (*Ambrosia* spp.)  
 Smartweed (*Polygonum pensylvanicum*)  
 Spurge, annual (*Euphorbia* spp.)  
 Velvetleaf (*Abutilon theophrasti*)

Early postemergence tank mixtures of PROWL plus Bicep will aid in the control of and reduce competition from:

Cocklebur (*Xanthium strumarium*)  
 Morningglory (*Ipomea* spp.)  
 Wild proso millet (*Panicum miliaceum*)  
 Woolly cupgrass (*Eriochloa villosa*)  
 Yellow nutsedge (*Cyperus esculentus*)

**Broadcast Rates per Acre of PROWL Plus Bicep  
 Preemergence and Early Postemergence Tank Mixtures In Corn**

(PROWL may be applied in sweet corn in Illinois, Minnesota,  
 New York and Wisconsin ONLY.)

PROWL (pints) + Bicep 6L\*\* (quarts)

SOIL* TEXTURE	Less than 1.5% Organic Matter	1.5% to 3.0% Organic Matter	More than 3.0% Organic Matter
	PROWL + Bicep 6L	PROWL + Bicep 6L	PROWL + Bicep 6L
COARSE	1.5 to 2 + 1.5	2 + 1.5	3 + 1.8
MEDIUM	2 + 1.8	3 + 1.8	3 + 2.4
FINE	2 + 2.4	3 + 2.4	3 + 2.4 to 3

\* See the PROWL label for soil textures classified as coarse, medium and fine.

\*\* When using Bicep 4.5L use equivalent rates. One quart of Bicep 6L equals 1.33 quarts of Bicep 4.5L.

NOTE: If crop loss occurs due to weather conditions and corn is replanted, seeding depth must be below retilled area.

Observe all cropping restrictions on the PROWL and Bicep labels.

This label must be in the possession of the user at the time of herbicide application.

PC 8348 3/10/89

 **CYANAMID**  
 American Cyanamid Company  
 Agricultural Division  
 Crop Protection Chemicals Department  
 Wayne, NJ 07470 C-1989

**NOT REVIEWED**  
EPA Reg. No. 241-243 In Accordance with PR Notice 82-2.  
Based on Leaf Labeling Dated 7/15/89

# FOR USE IN FIELD CORN GROWN IN NO-TILL SYSTEMS EAST OF THE MISSISSIPPI RIVER

OBSERVE ALL PRECAUTIONARY STATEMENTS IN  
THE PROWL LEAFLET LABEL BEFORE USING.

See the PROWL leaflet label for Application, Mixing, and Spraying Instructions, Use Rates, Weed Species Controlled, and Rotational Crop Restrictions.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PROWL alone and in tank mix combinations may be applied in field corn grown in no-till systems. Plant corn AT LEAST 1 1/2 inches deep to ensure good seed coverage. CORN SEED MUST BE COMPLETELY COVERED WITH SOIL. The use of no-till planters under conditions which do not allow good soil coverage of the corn seed can result in reduced crop stand or injury if PROWL contacts germinating corn seed. Check equipment to ensure good seed coverage. Apply in 20 or more gallons of spray solution per acre. CHEMIGATION: Refer to supplemental labeling entitled, PROWL leaflet label for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

DO NOT apply tank mix combinations unless each product is labeled for use in no-till systems

This label must be in possession of the user at the time of herbicide application.



\* Registered Trademark of American Cyanamid Company

PE 8352 3/89

NOT REVIEWED  
In Accordance with PR Notice 82-2.  
Based on Draft Labeling Dated 2/11/89

EPA Reg. No. 241-243

# FOR USE IN DRY, LIMA, AND SNAP BEANS

## PROWL PLUS Dual 8E<sup>1</sup> TANK MIXTURE

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL AND  
Dual 8E LEAFLET LABELS BEFORE USING.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Uniformly apply PROWL plus Dual 8E with properly calibrated ground equipment in 10 to 40 gallons of water per acre, or by aircraft in a minimum of 5 gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. DO NOT apply this product through any type of irrigation system.


DO NOT apply on peat or muck soils.

DO NOT use in California.

### PREPLANT INCORPORATED APPLICATIONS

Apply immediately before planting, or up to 14 days prior to planting. Incorporate uniformly into the top 2 inches of soil within 7 days after application. See INCORPORATION INSTRUCTIONS section of the PROWL label for acceptable incorporation techniques. Use PROWL and Dual 8E rates listed in the table on page 3.

<sup>1</sup>Dual 8E is a trademark of CIBA-GEIGY Corp



When applied as directed, PROWL plus Dual 8E tank mix will control the following grass and broadleaf weed species.

### GRASSES CONTROLLED

---

Barnyardgrass	( <i>Echinochloa crus-galli</i> )
Crabgrass	( <i>Digitaria</i> spp.)
Crowfootgrass	( <i>Dactyloctenium aegyptium</i> )
Fall panicum	( <i>Panicum dichotomiflorum</i> )
Field sandbur	( <i>Cenchrus incertus</i> )
Foxtails	
giant	( <i>Setaria faberi</i> )
green	( <i>Setaria viridis</i> )
yellow	( <i>Setaria glauca</i> )
Goosegrass	( <i>Eleusine indica</i> )
Johnsongrass (from seed)	( <i>Sorghum halepense</i> )
Red rice	( <i>Oryza sativa</i> )
Signalgrass	( <i>Brachiaria platyphylla</i> )
Southwestern cupgrass	( <i>Eriochloa gracilis</i> )
Texas panicum	( <i>Panicum texanum</i> )
Witchgrass	( <i>Panicum capillare</i> )
Yellow nutsedge	( <i>Cyperus esculentus</i> )

---

### BROADLEAF WEEDS CONTROLLED

---

Carpetweed	( <i>Mollugo verticillata</i> )
Florida pusley	( <i>Richardia scabra</i> )
Galinsoga	( <i>Galinsoga</i> spp.)
Kochia	( <i>Kochia scoparia</i> )
Lambsquarters	( <i>Chenopodium album</i> )
Pigweed	( <i>Amaranthus</i> spp.)
Purslane	( <i>Portulaca oleracea</i> )
Spurge, annual	( <i>Euphorbia</i> spp.)

---

Tank mixtures of PROWL plus Dual 8E will aid in the control of and reduce competition from:

---

Hairy nightshade	( <i>Solanum sarrachoides</i> )
Pennsylvania smartweed	( <i>Polygonum pennsylvanicum</i> )
Shattercane	( <i>Sorghum bicolor</i> )
Velvetleaf	( <i>Abutilon theophrasti</i> )
Volunteer sorghum	( <i>Sorghum vulgare</i> )
Wild proso millet	( <i>Panicum miliaceum</i> )

---

**Broadcast Rates per Acre of PROWL Plus Dual 8E  
Preplant Incorporated Tank Mixture In Dry, Lima or Snap Beans**

**PROWL (pints) + Dual 8E (pints)**

SOIL* TEXTURE	Northern States <sup>1</sup>		Southern States <sup>1</sup>	
	Up to 3% Organic matter	More than 3% Organic matter	Up to 3% Organic matter	More than 3% Organic matter
COARSE	1 to 2 + 1.5 to 2	2 + 2	1 to 1.5 + 1.5 to 2	1 to 1.5 + 2
MEDIUM	1.5 to 2.5 + 2 to 2.5	2.5 to 3 + 2 to 2.5	1.5 to 2 + 2 to 2.5	1.5 to 2 + 2 to 2.5
FINE	2 to 3 + 2 to 2.5	3 + 2.5 to 3	1.5 to 3 + 2 to 2.5	1.5 to 3 + 2.5 to 3

<sup>1</sup>See map at end of this label for specific states. \*See the PROWL label for soil textures classified as coarse, medium and fine

**NOTE:** If loss of dry, lima or snap bean crop occurs due to weather conditions, any crop registered for preplant incorporated use on the PROWL and DUAL 8E labels may be replanted the same year without adverse effects. If replanting is necessary, DO NOT rework the soil deeper than the treated zone.


**PRECAUTION:** DO NOT APPLY PREEMERGENCE (surface treatment) to dry, lima and snap beans or serious crop injury can result.

DO NOT graze or feed forage or fodder.

Observe all cropping restrictions on the PROWL and Dual 8E labels.

This label must be in the possession of the user at the time of herbicide application.

**REGIONAL MAP FOR RATE DETERMINATION**




**CYANAMID**  
 American Cyanamid Company  
 Agricultural Division  
 Crop Protection Chemicals Department  
 Wayne, NJ 07470 © 1989

NOT REVIEWED  
In Accordance with PR Notice 82-2.  
EPA Reg. No. 241-~~243~~ on Draft Labeling Dated

## FOR USE IN SOYBEANS

### PROWL PLUS CANOPY<sup>1</sup> TANK MIXTURE

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL AND CANOPY LEAFLET LABELS BEFORE USING.

See the PROWL leaflet label for Mixing Instructions, Incorporation Instruction., and Liquid Fertilizer Compatibility Determinations. Use one-half the teaspoons of herbicide specified for AAtrex Nine-0<sup>2</sup> or atrazine 4L when determining liquid fertilizer compatibility with Canopy.

#### GENERAL INFORMATION

Destroy existing weeds before applying PROWL tank-mix combinations. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Applied according to label directions and under normal growing conditions, PROWL tank-mix combinations will not cause crop injury. Over-application can result in crop stand loss, crop injury, or soil residues. Uneven application or improper soil incorporation can decrease weed control or cause crop injury.

Seedling diseases, cold weather, excessive moisture, deep planting, low or high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of crop damage from PROWL. Under these conditions, crop yields can be reduced.

<sup>1</sup>Canopy is a trademark of E. I. du Pont de Nemours & Co. (Inc.)  
<sup>2</sup>AAtrex Nine-0 is a trademark of Ciba-Geigy Corporation

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Uniformly apply PROWL plus Canopy with a properly calibrated low pressure (20 to 40 psi) fixed-boom sprayer equipped with fan type nozzles and screens no finer than 50 mesh. Use 10 to 40 gallons of water or 20 or more gallons of liquid fertilizer per acre. DO NOT apply this product through any type of irrigation system.

Continuous agitation in the spray tank is required to keep the material in suspension. Avoid overlap and shut off spray booms while starting, turning, slowing or stopping, or crop injury may result.

DO NOT apply on peat or muck soils or soils having a calcareous surface layer.

DO NOT apply to soils with a pH greater than 7.0, except as directed on the Canopy leaflet label.

Consult the Canopy label for sensitive soybean varieties. Before use on any soybean variety, tolerance to Canopy must first be determined.

#### PREPLANT INCORPORATED APPLICATIONS

Apply immediately before planting or up to two weeks prior to planting. Incorporate prior to planting and within 7 days of application. Follow the Incorporation Instructions in the PROWL leaflet label. Use the PROWL and Canopy rates listed in the table on page 2.

7 days after application.

If moisture is insufficient to activate the herbicides, a shallow cultivation (preferably with a rotary hoe) should be made after emergence of soybeans, but while weeds are still small enough to be controlled by mechanical cultivation.

DO NOT make applications of PROWL preemergence north of Interstate 80.

### MINIMUM AND NO-TILL SYSTEMS

PROWL plus Canopy may be applied preemergence in minimum or no-till soybeans. Emerged weeds will be suppressed by this tank mixture. Roundup<sup>3</sup> or paraquat may be added to the tank mixture to achieve control of emerged weeds.

Uniformly apply in 20 to 60 gallons of water per acre with ground equipment. Use the higher gallonage for fields with heavy weed infestations or excessive crop residues.

### WEED SPECIES CONTROLLED

When applied as directed, PROWL plus Canopy tank mix will control the following grass and broadleaf weed species:

#### GRASSES CONTROLLED

Barnyardgrass	( <i>Echinochloa crus-galli</i> )
Crabgrass	( <i>Digitaria</i> spp.)
Crowfootgrass	( <i>Dactyloctenium aegyptium</i> )
Field sandbur	( <i>Cenchrus incertus</i> )
Foxtails	
Giant	( <i>Setaria faberi</i> )
Green	( <i>Setaria viridis</i> )
Yellow	( <i>Setaria glauca</i> )
Goosegrass	( <i>Hesperis indica</i> )
Johnsongrass (from seed)	( <i>Sorghum halepense</i> )
Panicum	
Foil	( <i>Panicum dichotomiflorum</i> )
Texas	( <i>Panicum texanum</i> )
Signalgrass	( <i>Brachiaria platyphylla</i> )
Witchgrass	( <i>Panicum capillare</i> )

<sup>3</sup>Roundup is a trademark of Monsanto Co.

Lambsquarters	( <i>Chenopodium album</i> )
Morningglory*	
Entireleaf	( <i>Ipomoea hederacea</i> var. <i>intergruscula</i> )
Ivyleaf	( <i>Ipomoea hederacea</i> )
Pitted	( <i>Ipomoea lacunosa</i> )
Smallflower	( <i>Jacquemontia tamnifolia</i> )
Tall	( <i>Ipomoea purpurea</i> )
Pigweed	( <i>Amaranthus</i> spp.)
Prickly sida (teaweed)	( <i>Sida spinosa</i> )
Purslane	( <i>Portulaca oleracea</i> )
Ragweed	
Common	( <i>Ambrosia artemisiifolia</i> )
Giant*	( <i>Ambrosia trifida</i> )
Sicklepod*	( <i>Cassia obtusifolia</i> )
Smartweed	( <i>Polygonum pennsylvanicum</i> )
Spurge, annual	( <i>Euphorbia</i> spp.)
Sunflower	( <i>Helianthus annuus</i> )
Velvetleaf	( <i>Abutilon theophrasti</i> )

Tank mixtures of PROWL plus Canopy will aid in the control of and reduce competition from:

Itchgrass	( <i>Rottboellia exaltata</i> )
Red rice	( <i>Oryza sativa</i> )
Shattercane	( <i>Sorghum bicolor</i> )

\*Cultivation or application of a postemergence herbicide may be required for season long control

### Broadcast Rates per Acre of PROWL Plus Canopy Preplant Incorporated and Preemergence Tank Mixtures

	PROWL (pints)+ Canopy (ounces)	
	0.5%** to 3% Organic matter	More than 3% Organic matter
SOIL*		
TEXTURE	PROWL + Canopy	PROWL + Canopy
COARSE	1 + 6 to 10	1.5 + 6 to 10
MEDIUM	1.5 + 8 to 12	1.5 to 2 + 8 to 12
FINE	1.5 + 10 to 14	2 to 2.5 + 10 to 14

\*See the PROWL label for soil textures classified as coarse, medium and fine

\*\*DO NOT apply to soil with less than 0.5% organic matter

Use the higher rates on soils with higher organic matter or heavy weed pressure.

NOTE: If crop loss occurs due to weather conditions and corn is replanted, seeding depth must be below retiled area.

Observe all cropping restrictions on the PROWL and Canopy labels. Always follow the most restrictive label

This label must be in the possession of the user at the time of herbicide application.



**CYANAMID**

American Cyanamid Company

Agricultural Division

Crop Protection Chemicals Department

Wayne, NJ 07470

©1988

EPA Reg. No. 241-243

NOT REVIEWED

in accordance with EPA Notice 82-2.  
2/16/81

## FOR USE IN SOYBEANS

### PROWL PLUS PREVIEW<sup>1</sup> TANK MIXTURE

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL AND PREVIEW LEAFLET LABELS BEFORE USING.

See the PROWL leaflet label for Mixing Instructions, Incorporation Instructions, and Liquid Fertilizer Compatibility Determinations. Use one-half the teaspoons of herbicide specified for AAtrex Nine-0<sup>2</sup> of atrazine 4L when determining liquid fertilizer compatibility with Preview.

#### GENERAL INFORMATION

Destroy existing weeds before applying PROWL tank-mix combinations. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

Applied according to label directions and under normal growing conditions, PROWL tank-mix combinations will not cause crop injury. Over-application can result in crop stand loss, crop injury, or soil residues. Uneven application or improper soil incorporation can decrease weed control or cause crop injury.

Seedling diseases, cold weather, excessive moisture, deep planting, low or high soil pH, high soil salt concentration, or drought can weaken seedlings and plants and increase the possibility of crop damage from PROWL. Under these conditions, crop yields can be reduced.

<sup>1</sup>Preview is a trademark of E. I. du Pont de Nemours & Co. (Inc.)  
<sup>2</sup>AAtrex Nine-0 is a trademark of Ciba-Geigy Corporation.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Uniformly apply PROWL plus Preview with a properly calibrated low pressure (20 to 40 psi) fixed-boom sprayer equipped with fan type nozzles and screens no finer than 50 mesh. Use 10 to 40 gallons of water or 20 or more gallons of liquid fertilizer per acre. DO NOT apply this product through any type of irrigation system.

Continuous agitation in the spray tank is required to keep the material in suspension. Avoid overlap and shut off spray booms while starting, turning, slowing or stopping, or crop injury may result.

DO NOT apply on peat or muck soils, soils having a calcareous surface layer, or soils with a pH greater than 6.8.

Consult the Preview label for sensitive soybean varieties. Before use on any soybean variety, tolerance to Preview must first be determined.

#### PREPLANT INCORPORATED APPLICATIONS

Apply immediately before planting or up to two weeks prior to planting. Incorporate prior to planting and within 7 days of application. Follow the Incorporation Instructions on the PROWL leaflet label. Use the PROWL and Preview rates listed in the table on page 2.

7 days after application.

If moisture is insufficient to activate the herbicides, a shallow cultivation (preferably with a rotary hoe) should be made after emergence of soybeans, but while weeds are still small enough to be controlled by mechanical cultivation.

DO NOT make applications of PROWL preemergence north of Interstate 80.

### MINIMUM AND NO-TILL SYSTEMS

PROWL plus Preview may be applied preemergence in minimum or no-till soybeans. Emerged weeds will be suppressed by this tank mixture. Roundup<sup>3</sup> or paraquat may be added to the tank mixture to achieve control of emerged weeds.

Uniformly apply in 20 to 60 gallons of water per acre with ground equipment. Use the higher gallonage for fields with heavy weed infestations or excessive crop residues.

### WEED SPECIES CONTROLLED

When applied as directed, PROWL plus Preview tank mix will control the following grass and broadleaf weed species:

### GRASSES CONTROLLED

Barnyardgrass	( <i>Echinochloa crus-galli</i> )
Crabgrass	( <i>Digitaria</i> spp.)
Crowfootgrass	( <i>Dactyloctenium aegyptium</i> )
Field sandbur	( <i>Cenchrus incertus</i> )
Foxtails	
Giant	( <i>Setaria faberi</i> )
Green	( <i>Setaria viridis</i> )
Yellow	( <i>Setaria glauca</i> )
Goosegrass	( <i>Eleusine indica</i> )
Johnsongrass (from seed)	( <i>Sorghum halepense</i> )
Panicum	
Fall	( <i>Panicum dichotomiflorum</i> )
Texas	( <i>Panicum texanum</i> )
Signalgrass	( <i>Bracharia platyphylla</i> )
Witchgrass	( <i>Panicum capillare</i> )

<sup>3</sup>Roundup is a trademark of Monsanto Co.

Rigweed	( <i>Amaranthus</i> spp.)
Prickly sida (teaweed)	( <i>Sida spinosa</i> )
Purslane	( <i>Portulaca oleracea</i> )
Ragweed, common	( <i>Ambrosia artemisiifolia</i> )
Smartweed	( <i>Polygonum pensylvanicum</i> )
Spurge, annual	( <i>Euphorbia</i> spp.)
Sunflower*	( <i>Helianthus annuus</i> )
Velvetleaf	( <i>Abutilon theophrasti</i> )

Tank mixtures of PROWL plus Preview will aid in the control of and reduce competition from:

Black nightshade	( <i>Solanum nigrum</i> )
Burcucumber	( <i>Sicyos angulatus</i> )
Itchgrass	( <i>Rottboellia exaltata</i> )
Morningglory*	
Entireleaf	( <i>Ipomoea hederacea</i> var. <i>intergriuscula</i> )
Ivyleaf	( <i>Ipomoea hederacea</i> )
Pitted	( <i>Ipomoea lacunosa</i> )
Tall	( <i>Ipomoea purpurea</i> )
Nutsedge	( <i>Cyperus</i> spp.)
Ragweed, giant*	( <i>Ambrosia trifida</i> )
Red rice	( <i>Oryza sativa</i> )
Shattercane	( <i>Sorghum bicolor</i> )

\*Cultivation may be required to achieve season long control

### Broadcast Rates per Acre of PROWL Plus Preview Preplant Incorporated and Preemergence Tank Mixtures

SOIL* TEXTURE	PROWL (pints)+ Preview (ounces)	
	0.5%** to 3% Organic matter	More than 3% Organic matter
COARSE	1+6	1.5+7
MEDIUM	1.5+7	1.5 to 2+8
FINE	1.5+8	2 to 2.5+9 to 10

\*See the PROWL label for soil textures classified as coarse, medium and fine.

\*\*DO NOT apply to soil with less than 0.5% organic matter

NOTE: If crop loss occurs due to weather conditions and corn is replanted, seeding depth must be below retilled area.

Observe all cropping restrictions on the PROWL and Preview labels. Always follow the most restrictive label.

This label must be in the possession of the user at the time of herbicide application.



**CYANAMID**  
American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 ©1989