

FOR USE IN SWEET CORN IN ARIZONA AND CALIFORNIA

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL LEAFLET LABEL BEFORE USING.

See the PROWL leaflet label for Application, Mixing, and Spraying Instructions, Use Rates, Weed Species Controlled, and Rotational Crop Restrictions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. PROWL and PROWL tank-mix combinations may be applied preemergence in sweet corn in Arizona and California. DO NOT INCORPORATE or serious corn injury can result.

Plant into a seed bed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. Plant corn AT LEAST 1 1/2 inches deep to ensure good seed coverage.

PROWL or PROWL tank-mix combination treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage (such as rotary hoe), and make certain corn seeds are below the tilled area. If furrow irrigation is used, the beds must be thoroughly wetted or poor seed control can result.

See the PROWL leaflet label for approved tank mixtures in sweet corn. When PROWL is tank mixed with other herbicides, follow all cropping restrictions, precautionary statements, and limitations in the leaflet labels of the tank-mix partners. Always follow the most restrictive label.

When PROWL alone is used, if corn loss occurs due to weather conditions, corn or cotton can be replanted the same year without adverse effects. If corn is replanted, seeding depth must be below retilled area.


DO NOT use PROWL on peat or muck soils.

DO NOT apply this product through any type of irrigation system.

This label must be in the possession of the user at the time of herbicide application.

Registered Trademark of American Cyanamid Company

PE-8365



CYANAMID

American Cyanamid Company
Agricultural Division
Crop Protection Chemical Department
Wayne, NJ 07470
E-1009

PM23
211 213
1.75

FOR USE IN SWEET CORN IN ARIZONA AND CALIFORNIA

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL LEAFLET LABEL BEFORE USING.

See the PROWL leaflet label for Application, Timing, and Spraying Instructions, Use Rates, Weed Species Controlled, and Rotational Crop Restrictions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. PROWL and PROWL tank-mix combinations may be applied preemergence in sweet corn in Arizona and California. DO NOT INCORPORATE or serious corn injury can result.

Plant into a seed bed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. Plant corn AT LEAST 1 1/2 inches deep to ensure good seed coverage.

PROWL or PROWL tank-mix combination treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage (such as rotary hoe), and make certain corn seeds are below the tilled area. If furrow irrigation is used, the beds must be thoroughly wetted or poor seed control can result.

See the PROWL leaflet label for approved tank mixtures in sweet corn. When PROWL is tank mixed with other herbicides, follow all cropping restrictions, precautionary statements, and limitations in the leaflet labels of the tank-mix partners. Always follow the most restrictive label.

When PROWL alone is used, if corn loss occurs due to weather conditions, corn or cotton can be replanted the same year without adverse effects. If corn is replanted, seeding depth must be below retilled area.


DO NOT use PROWL on peat or muck soils.

DO NOT apply this product through any type of irrigation system.

This label must be in the possession of the user at the time of herbicide application.

©Registered Trademark of American Cyanamid Company

PE-8365



CYANAMID
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470

NOT REVIEWED

EPA Reg. No. 241-243

In Accordance with PR Notice 82-11

Based on Draft Labeling Dated

5/25/89

FOR USE IN SWEET CORN IN ARIZONA AND CALIFORNIA

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL LEAFLET LABEL BEFORE USING.

See the PROWL leaflet label for Application, Mixing, and Spraying Instructions, Use Rates, Weed Species Controlled, and Rotational Crop Restrictions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. PROWL and PROWL tank-mix combinations may be applied preemergence in sweet corn in Arizona and California. DO NOT INCORPORATE or serious corn injury can result.

Plant into a seed bed that is firm and free of clods and trash. Use only where adequate tillage is practiced to provide good soil coverage of the corn seed. Plant corn AT LEAST 1 1/2 inches deep to ensure good seed coverage.

PROWL or PROWL tank-mix combination treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application. If cultivation is necessary because of soil crusting, soil compaction or weed germination before rain or irrigation, use shallow tillage (such as rotary hoe), and make certain corn seeds are below the tilled area. If furrow irrigation is used, the beds must be thoroughly wetted or poor seed control can result.

See the PROWL leaflet label for approved tank mixtures in sweet corn. When PROWL is tank mixed with other herbicides, follow all cropping restrictions, precautionary statements, and limitations in the leaflet labels of the tank-mix partners. Always follow the most restrictive label.

When PROWL alone is used, if corn loss occurs due to weather conditions, corn or cotton can be replanted the same year without adverse effects. If corn is replanted, seeding depth must be below retilled area.

DO NOT use PROWL on peat or muck soils.

DO NOT apply this product through any type of irrigation system.

This label must be in the possession of the user at the time of herbicide application.

®Registered Trademark of American Cyanamid Company

PE-8363 4/19/89

 **CYANAMID**
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1989