

EPA Reg. No. 241-243

FOR USE IN OR ON GARLIC IN ARIZONA

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE PROWL LEAFLET LABEL BEFORE USING. SEE PROWL LEAFLET LABEL FOR APPLICATION INFORMATION AND WEED SPECIES CONTROLLED. OBSERVE ALL FOLLOW CROP RESTRICTIONS IN THE PROWL LEAFLET LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. PROWL herbicide controls most annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds. Destroy existing weeds before apply PROWL. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend it over a long period of time may reduce weed control.

PROWL will not cause crop injury when applied according to the label under normal growing conditions. Over-application may result in injury to crops, poor stands, or soil residues; conversely, uneven applications or incorporating the product too deeply may reduce weed control.

Disease, cold weather, excessive moisture, deep planting, low or high pH, salinity, or drought may weaken seedlings and plants and make them more susceptible to herbicidal damage.

**PROWL APPLIED PREMERGENCE,
POSTEMERGENCE OR PREEMERGENCE FOLLOWED
BY POSTEMERGENCE IN OR ON GARLIC**

Apply PROWL at a broadcast rate of 1.5 to 3 pints per acre. Uniformly apply PROWL in a minimum of 10 gallons of water per acre.

PROWL may be applied in the following ways:

- (a) Preemergence after planting but before crop and weeds emerge.
- (b) Postemergence to the garlic at the 1 to 5 true leaf growth stage.
- (c) As a split application, apply at both preemergence and postemergence timings.

DO NOT exceed 3 pints per acre per crop. **DO NOT** apply within 5 months of harvest. PROWL treatments are most effective when adequate rainfall or overhead irrigation is received within 7 days after application.

DO NOT USE on peat or muck soils.

NOTE: If loss of garlic crop occurs due to adverse weather conditions, any crop registered for PROWL pre-plant incorporated use can be replanted the same year into PROWL treated soil without adverse effects. If replanting is necessary, **DO NOT** rework the soil deeper than the PROWL treated zone or 2 inches deep if surface applied.

DO NOT feed or graze this crop.

This label must be in the possession of the user at the time of herbicide application.

*Registered Trademark of American Cyanamid Company
PE- 2407 10/1/90 DL



American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 © 1990

