

**RESTRICTED USE PESTICIDE**  
**Due to acute oral and dermal toxicity and bird toxicity.**


For retail sale to, and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.

American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

EPA REG. NO. 241-257

**DIRECTIONS FOR USE IN POTATOES**

**BEFORE USING, READ PRECAUTIONARY  
 STATEMENTS ON BAG LABEL**



It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. THIMET 20-G should be applied with a granular pesticide applicator properly calibrated to assure placement and proper dosage. See label for specific instructions.

Cover granules that may be exposed on the ends of the treated rows, turns and loading areas by deep discing immediately after treating fields.

Crop	Pests Controlled	Rate of THIMET 20-G	Application	Remarks
POTATOES At Planting	Aphids	Light or Sandy Soils:	Distribute the granules evenly in the furrow or band on each side of the row.	Wait 90 days after treatment before harvesting potatoes.
	Leafhoppers	11.3 lbs. per		
	Leaf miners	1,000 sq. ft.		
	Psyllids	row for any		
	Wireworms	row spacing		
	Flea beetle larvae and reduction of flea beetle adults	(minimum 32-inch spacing)		Do not use for Colorado potato beetle control in Long Island, New York and Del Marva Peninsula.
	Colorado potato beetle (early season control)	Heavy or Clay Soils:		
		17.3 lbs. per		
		1,000 sq. ft.		
		row for any		
		row spacing		
		(minimum 32-inch spacing)		

**THIS LABEL SHOULD BE IN THE POSSESSION OF THE  
 USER AT THE TIME OF PESTICIDE APPLICATION**

