

241-257
CONTAINER

1.1.14

RESTRICTED USE CLASSIFICATION

Due to acute oral and dermal toxicity and avian hazards

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification. American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

THIMET® 20-G
soil and systemic insecticide
LOCK'nLOAD™
CLOSED HANDLING SYSTEM

Active Ingredient: Phorate: 0,0-diethyl S-[(ethylthio) methyl] phcsphorodithioate.....	20.0%
Inert Ingredients.....	80.0%
Total.....	100.0%

EPA Reg. No. 241-257

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO!



POISON



PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

FIRST AID

If **swallowed**, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.


If **inhaled**, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get medical attention.

If **on skin**, wash thoroughly with soap and water. Remove contaminated clothing and shoes. Wash clothing and decontaminate shoes before reuse.

If **in eyes**, immediately flush eyes with plenty of water. Get medical attention.

Antidote: Atropine is an antidote.

See Side Panel For Additional Precautionary Statements



Handwritten notes:
1.1.14
1.1.14

LABEL NOT REFINISHED
PER PH NOTICE 88-6
DATE 10/2/90

NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, nausea, vomiting, diarrhea, and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine.

Use according to label directions.

IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.


SOLD BY
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1990

Net Weight: 40 lbs.

8/90

™Trademark of American Cyanamid Company

®Registered Trademark of American Cyanamid Company



2019

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER!**

Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

Do Not Breathe Dust

While emptying container into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

Do Not Contaminate Food or Feed Products

Once container has been opened, use it completely. Make sure the hoppers are emptied while still in the field. Refer to STORAGE AND DISPOSAL statement for further instructions.

Keep all unprotected persons out of operating areas.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Keep Out of Reach of Domestic Animals.


Not for Use or Storage in or Around the Home.

See leaflet for Re-entry Statement and Work Safety Rules

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife. Birds feeding in treated areas may be killed. Do not apply directly to water or wetlands (swamps, bogs, marshes, and pot-holes). Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Do not contaminate water when disposing of equipment washwaters.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See attached leaflet for complete Directions For Use. See leaflet for Disclaimer.

STORAGE AND DISPOSAL

Storage

Store pesticide products in a secure locked area where children, unauthorized persons and animals cannot enter. Do not store in the same area with food or feed. Do not store opened containers.

Prohibitions

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Cover or incorporate spills.

Pesticide Disposal


Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Return empty container to point of purchase.

General

Consult Federal, State or local disposal authorities for approved alternative procedures.



RESTRICTED USE CLASSIFICATION

Due to acute oral and dermal toxicity and avian hazards

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification. American Cyanamid Company endorses Certification to promote the responsible use of pesticides to insure the protection of man and the environment.

**THIMET® 20-G
soil and systemic insecticide
LOCK'nLOAD™
CLOSED HANDLING SYSTEM**

Active Ingredient: Phorate: 0,0-diethyl S-[(ethylthio) methyl] phosphorodithioate.....	20.0%
Inert Ingredients.....	80.0%
Total.....	100.0%

EPA Reg. No. 241-257

KEEP OUT OF REACH OF CHILDREN



DANGER/PELIGRO!



POISON

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

FIRST AID


If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Avoid alcohol. Get medical attention.

If inhaled, remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get medical attention.

If on skin, wash thoroughly with soap and water. Remove contaminated clothing and shoes. Wash clothing and decontaminate shoes before reuse.

If in eyes, immediately flush eyes with plenty of water. Get medical attention.

Antidote: Atropine is an antidote.



See Next Page For Additional Precautionary Statements

10/19
J


NOTE TO PHYSICIANS: Warning symptoms include weakness, headache, tightness in chest, blurred vision, nonreactive pinpoint pupils, salivation, nausea, vomiting, diarrhea, and abdominal cramps. Give atropine intramuscularly or intravenously, depending on severity of poisoning, 2 to 4 milligrams every 10 minutes until fully atropinized as shown by dilated pupils, dry flushed skin and tachycardia. Twenty to thirty milligrams, or more, may be required during the first 24 hours. Never give opiates or phenothiazine tranquilizers. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for at least 48 hours. Allow no further exposure to any cholinesterase inhibitor until cholinesterase regeneration has taken place as determined by blood tests.

Pralidoxime chloride (2-PAM; PROTOPAM chloride) may be effective as an adjunct to atropine.
Use according to label directions.

IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS PRODUCT, CALL COLLECT, DAY OR NIGHT, AREA CODE 201-835-3100.

SOLD BY
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1990

TM Trademark of American Cyanamid Company
® Registered Trademark of American Cyanamid Company



7-19
1

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER!**

Fatal if swallowed, inhaled or absorbed through the skin. Do not breathe dust. Do not get in eyes, on skin, or on clothing.

Do Not Breathe Dust

While emptying container into equipment, pour downwind and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach the breathing zone.

Do Not Contaminate Food or Feed Products

Once container has been opened, use it completely. Make sure the hoppers are emptied while still in the field. Refer to STORAGE AND DISPOSAL statement for further instructions.

Keep all unprotected persons out of operating areas.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons.

Keep Out of Reach of Domestic Animals.

Not for Use or Storage in or Around the Home.


REENTRY

After dusts have settled, do not enter or allow entry into the treated area until the 48 hour reentry interval has expired, unless the person entering the treated area is wearing the personal protective equipment listed below for early reentry.

For early reentry into treated areas wear a protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical-resistant gloves and chemical resistant shoes, coverings, or boots.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language, customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER. Area treated with THIMET on (date of application). Do not enter without protective clothing.



PERSONAL PROTECTIVE EQUIPMENT

WEAR THE FOLLOWING PROTECTIVE CLOTHING DURING LOADING, APPLICATION, EQUIPMENT REPAIR/CLEANING, AND DISPOSAL OF THE PESTICIDE. A protective suit of one or two pieces covering all parts of the body except head, hands, and feet; chemical resistant gloves; chemical resistant shoes, shoe covering, or boots.

Unless a chemical resistant protective suit is otherwise required, mixers/loaders must wear a chemical resistant apron and goggles or a face shield.

IF APPLICATION IS PERFORMED USING AN ENCLOSED CAB OR COCKPIT, THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT MAY BE WORN AS AN ALTERNATE: Long-sleeved shirt and long-legged pants; shoes and socks. Chemical resistant goves must be available in the cab or cockpit and must be worn when exiting. This clothing is inadequate to protect you during equipment repair or cleaning, reentry, or pesticide disposal work. Do not allow contaminated clothing to come in contact with the insides of the cab.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water. Always wash hands, face, and arms with soap and water before smoking, eating, drinking, or using the toilet.

AFTER WORK: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Wear clean clothes. Do not reuse contaminated clothing. Personal clothing worn during work must be must be laundered separately from household articles. Store protective clothing separately from personal clothing. Clean or launder protective clothing after each use. Clothing and protection equipment heavily contaminated must be destroyed according to state and local regulations. HEAVILY CONTAMINATED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

WORK SAFETY RULE

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

Persons working with this product should have frequent blood tests of their cholinesterase levels. If the cholinesterase level falls below a critical point, no further exposures should be allowed until it has been determined by means of blood tests that the cholinesterase level has return to normal. Before using this product consult the National Pesticide Telecommunications Network for recommendations regarding such blood tests, poisoning management, emergency treatment, and other information regarding the toxicity of phorate. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7378.



0.19

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away, see First Aid section.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife. Birds feeding in treated areas may be killed. Do not apply directly to water or wetlands (swamps, bogs, marshes, and pot-holes). Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application.

STORAGE AND DISPOSAL

Storage

Store pesticide products in a secure locked area where children, unauthorized persons and animals cannot enter. Do not store in the same area with food or feed. Do not store opened containers.

Prohibitions

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Cover or incorporate spills.

Pesticide Disposal


Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Return empty container to point of purchase.

General

Consult Federal, State or local disposal authorities for approved alternative procedures.



DISCLAIMER


The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein when used in accordance with the directions for use, subject to the risk referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.


BEFORE USING, READ PRECAUTIONARY STATEMENTS.

THIMET should be applied with a granular pesticide applicator properly calibrated to assure accurate placement and proper dosage. Cover granules that may be exposed on the ends of the treated rows and turns and loading areas by deep discing immediately after treating fields.



CROPS

Crop	Pests Controlled	Rate of THIMET 20-G	Application	Remarks
BEANS At Planting	Mexican bean beetles	4.9-9.4 ozs. per 1,000 ft. of row for any row (minimum 30-inch spacing)	Drill granules to the side of the seed	Do not place THIMET granules in direct contact with seed. Do not graze or feed treated hay or forage to livestock. Do not apply later than 60 days before harvest.
	Leafhoppers Aphids Lygus bugs Thrips Mites Seedcorn maggots	4.5-7.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch spacing)		
	Seedcorn maggots		Apply granules in a band over the row	



FIELD CORN
AND SWEET
CORN
(excluding
popcorn)

Corn
rootworms
Wireworms
White grubs
Seedcorn
maggots
Seedcorn
beetles
Flea beetles
Mites

6 ozs. per 1,000
ft. of row for any
row spacing
(minimum 30-inch
row spacing)

Place granules
in a 7-inch
band over the
row directly
behind the
planter shoe in
front of or
behind the
press wheel and
lightly
incorporate.

Do not use in-
furrow
application.
One additional
application can
be applied at
cultivation

At Planting

FIELD CORN
AND SWEET
CORN
(excluding
popcorn)

Corn
rootworms

Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil.

Do not make any
applications of
THIMET after
cultivation
treatment.

At
Cultivation

Do not harvest
within 30 days
of last
treatment.

FIELD CORN
(excluding
sweet and
popcorn)

Chinch bug
nymphs
(Colorado,
Kansas, and
Nebraska)

Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil.
Apply as soon
as chinch bug
nymphs are
observed at
base of plants.
Granules must
be placed at
the infested
region of the
plants and
a equately
covered with
soil.

Later
infestations of
chinch bugs may
not be
controlled.

At
Cultivation

Do not make any
applications of
THIMET after
cultivation
treatment.

Do not harvest
within 30 days
of last
treatment.

Do not graze or
cut for forage
within 30 days
of last
treatment.

Do not apply
more than 2
applications
per season.

<p>COTTON At Planting</p>	<p>Early season control of: Thrips Mites Leaf miners</p>	<p>3.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)</p>	<p>Distribute the granules evenly in the furrow at planting time. Soil temperature moisture and tilth should be favorable for good germination and emergence.</p>	<p>Suggested for areas where a limited period of protection against early season pests is necessary for good cotton growth. Use 1.5 lbs. THIMET 20-G per acre for hill-dropped cotton. Do not feed treated hay or forage to livestock.</p>
	<p>Mites Thrips Aphids Black cutworms Leafhoppers White flies Leaf miners</p>	<p>6.0-9.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)</p>	<p>Distribute the granules evenly in the furrow at planting time. Soil temperature, moisture and tilth should be favorable for good germination and emergence. Under adverse conditions plant stand may be affected.</p>	<p>Recommended for areas requiring and extended period of mite control as well as early season protection against other cotton pests. The higher rate is suggested for heavier soils. Do not feed treated hay or forage to livestock.</p>
<p>COTTON Side-Dressing (Irrigated Cotton Only)</p>	<p>Mites from spring to early summer</p>	<p>9.0-12.0 ozs. per 1,000 ft. of row for any row spacing (minimum 36-inch spacing)</p>	<p>Incorporate granules into the soil as a side-dress application deep enough to avoid disturbance by future cultivations.</p>	<p>Irrigate as soon as possible after treatment. Do not apply later than 60 days before harvest. Do not feed treated hay or forage to livestock. Do not apply more than 2 applications per season</p>

<p>PEANUTS At Planting</p>	<p>Thrips Leafhoppers</p>	<p>5.5 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)</p>	<p>Distribute the granules evenly in the furrow at planting time. For best results, plant peanuts when soil temperatures are favorable for establishing good peanut stands.</p>	<p>Do not graze or feed treated hay or forage to livestock. Wait 90 days after treatment before harvesting peanuts.</p>
<p>PEANUTS At Pegging</p>	<p>Southern corn rootworms and Leafhoppers</p>	<p>11.0 ozs. per 1,000 ft. of row for any row spacing (minimum 24-inch spacing)</p>	<p>Distribute the granules as a band over the fruiting zone at pegging time. Work into the top few inches of soil immediately.</p>	<p>Do not graze or feed treated hay or forage to livestock. Wait 90 days after treatment before harvesting peanuts. Do not apply more than 2 applications per season.</p>
<p>POTATOES At Planting</p>	<p>Aphids Leafhoppers Leaf miners Psyllids Wireworms Flea beetle larvae and reduction of flea beetle adults.</p>	<p>Light or Sandy Soils 11.3 ozs. per 1,000 ft. of row for any spacing (minimum 32-inch spacing)</p>	<p>Distribute the granules evenly in the furrow or band on each side of the row.</p>	<p>Wait 90 days after treatment before harvesting potatoes.</p>
	<p>Colorado potato beetle (early season control)</p>	<p>Heavy or Clay Soils: 17.3 ozs. per 1,000 ft. of row for any row spacing (minimum 32-inch spacing)</p>		<p>Do not use for Colorado potato beetle control in Long Island area of New York and the Del Marva Peninsula.</p>

SORGHUM
At Planting

Greenbugs

6.0 ozs. per 1,000
ft. of row for any
row spacing
(minimum 30-inch
spacing)

Place granules
in a 7-inch
band over the
row OR drill 1
inch below and
2 inches to the
side of the
seed.

Do not place
THIMET granules
in direct
contact with
seed. A second
application of
granules may be
applied at
cultivation
later in the
season.

SORGHUM
At
Cultivation

Chinch bug
nymphs
(Colorado,
Kansas,
Nebraska)

Apply granules
to base of
plants just
ahead of
cultivation
shovels so as
to cover
granules with
soil. Apply as
soon as chinch
bug nymphs are
observed at
base of plants.
Granules must
be placed at
the infested
region of the
plants and
adquately
covered with
soil.


Later
infestation of
chinch bugs may
not be
controlled.

Do not make any
application of
THIMET after
cultivation
treatment.

Do not feed
foliage before
grain harvest.

Do not apply
within 30 days
of harvest.

Do not apply
more than 2
applications
per season.




SOYBEANS At Planting	For the early season control of: Mexican bean beetles Leafhoppers Lygus bugs Seedcorn maggots Thrips Aphids Mites	9.0 ozs. per 1,000 ft. of row for any row spacing (minimum 30-inch row spacing)	Drill granules to the side of the seed.	Do not place THIMET granules in direct contact with seed. Do not feed the foliage of treated soybeans. Crop injury may result when preplant incorporated or preemergence applications of metribuzine herbicides are used in conjunction with this product or other soil-applied organo-phosphate pesticides.
	<hr/> Seedcorn maggots		<hr/> Place granules in a 7-inch band over the row directly behind the planter shoe and in front of the press wheel.	

SUGAR BEETS	Aphids Beets root maggots Leafhoppers Mites Leaf miners	4.5 ozs. per 1,000 ft. of row for any row spacing (minimum 20-inch spacing)	Drill granules to the side of the seed OR place granules in a band over the row.	Do not place THIMET granules in direct contact with the seed. Do not apply within 30 days of harvest. Do not feed sugar beet tops or silage to dairy cattle. Consult your state experiment station or state extension service for proper timing of application in your area.
--------------------	---	---	--	--



17-219
✓


SUGAR BEETS Post-emergence	Mites Aphids	4.9-7.5 lbs. per acre	Apply to foliage when plants are dry. Use the higher rate on dense foliage for good coverage.	Do not apply by air. Do not apply more than 2 applications per season.
SUGAR CANE (Florida)	Wireworms	19.5 lbs. per acre	Apply in a 10-12 inch band directly on the seed piece and surrounding soil in the open furrow immediately before covering.	Do not graze or feed treated hay or forage to livestock.
WHEAT At Planting	Grasshoppers ¹ Aphids (Western and North Central States only) Hessian fly	1.2 ozs. per 1,000 ft. of row for any row spacing (minimum 8-inch spacing)	Apply the granules in the seed furrow.	Do not feed or graze foliage within 70 days of treatment. Do not make any later applications after planting time treatment. ¹ Grasshoppers controlled in winter wheat only. Do not harvest within 70 days of application.



TYPICAL APPLICATION RATES FOR VARIOUS ROW SPACES

	Rate Oz./1,000 Ft. of Row									
	<u>1.2 oz</u>	<u>3 oz</u>	<u>4.5 oz</u>	<u>6 oz</u>	<u>8 oz</u>	<u>10 oz</u>	<u>12 oz</u>	<u>14 oz</u>	<u>16 oz</u>	<u>18 oz</u>
40"			3.7	4.9	6.5	8.2	9.8	11.4	13.1	14.7
38"			3.9	5.2	6.9	8.6	10.3	12.0	13.7	15.5
36"			4.1	5.4	7.3	9.1	10.9	12.7	14.5	16.3
34"			4.3	5.8	7.7	9.6	11.5	13.5	15.4	17.3
32"			4.6	6.1	8.2	10.2	12.2	14.3	16.3	18.4
30"			4.9	6.5	8.7	10.9	13.1	15.2	17.4	19.6
26"			5.7	7.5	10.0	12.6	15.1	17.6	20.1	22.6
24"			6.1	8.2	10.9	13.6	16.3	19.1	21.8	24.5
20"		4.9								
18"		5.4								
16"		6.1								
12"		8.2								
8"	4.9	12.2								
6"	6.5	16.3								

EQUIVALENT POUNDS PER ACRE



CALIBRATION INFORMATION

FIRST READ THE LABEL

It is important that applicator equipment be properly set to deliver the labeled rate. The table below contains suggested starting gauge settings for calibration at planting speeds of 5 or 7 mph and application rates of 4.5, 6.0 or 9.0 oz. 1,000 ft. of row.

NOTE: The settings in this table SHOULD ONLY BE USED AS STARTING POINTS. Continually check the amount of THIMET 20-G used against a known length of row and make further adjustments accordingly. Also check calibration occasionally to make sure equipment wear, changing moisture conditions, etc. have not caused a change in flow rate.

SUGGESTED GRANULAR APPLICATOR SETTING¹

Applicator	Planter Speed 5 mph			Planter Speed 7 mph		
	4.5 oz.	6.0 oz.	9.0 oz.	4.5 oz.	6.0 oz.	9.0 oz.
John Deere MaxEmerge 2	14	17	23	17	21	33
John Deere Max-Emerge (Odd Notches 5-15-25-35)	9	11	14	11	13	17
John Deere Max-Emerge (Even Notches 0-10-20-30)	9	13	20	13	19	26
John Deere 71 Flexi-Planter (John Deere Metal Hoppers)	1/23	1/26	2/3	1/26	1/32	2/14

¹Suggested starting point only.

