

APPLICATION INSTRUCTIONS

Add PROWL DG slowly to partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. **MAINTAIN CONTINUOUS AGITATION WHILE ADDING PROWL DG AND UNTIL SPRAYING IS COMPLETED.** If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. **DO NOT** apply when winds will cause drift.

PROWL DG treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application.

APPLICATION TIMING

PROWL DG treatments will not control established weeds. Applications must be made prior to germination of weeds.

For control of annual bluegrass, chickweed and henbit, apply in late summer to early fall before weeds germinate.

For control of crabgrass and other annual weeds listed above, apply in the spring before weeds germinate.

Two applications per year may be made. Allow at least 4 months between applications.

APPLICATION RATE

Apply 4.5 lbs of PROWL DG per acre (1.0 lb/10,000 sq. ft.).

NOTE: Use only on well established turfgrass with a dense and uniform stand. **DO NOT** use on newly seeded or sprigged turfgrass. **DO NOT** use on winter-overseeded turfgrass. **DO NOT** use on turfgrass severely thinned by winter stress.

PROWL provides effective residual weed control which may affect new turfgrass seedlings or sprigs. **DO NOT** reseed or winter overseed turfgrass within 4 growing months after application. **DO NOT** sprig turfgrass within 5 months after application.

DO NOT use on bentgrass or dichondra.



CYANAMID

Agricultural Division
Bio-Tech Products Department
Wayne, NJ 07470 © 1984

PROWL[®] DG herbicide

SUPPLEMENTAL LABELING

EPA Reg. No. 241-268


FOR USE IN LAWNS AND TURF

PROWL DG can be used for weed control in lawns and turf.

TURFGRASS TYPES

PROWL DG should only be applied to well established lawns and turf. PROWL DG can be used on the following turfgrasses:

- Bahia
- Bermudagrass
- Centipedegrass
- St. Augustinegrass
- Tall fescue
- Zoysiagrass



WEED SPECIES CONTROLLED


When applied as directed, PROWL DG will control the following weed species:

GRASSES CONTROLLED

- Annual bluegrass (Poa annua)
- Barnyardgrass (Echinochloa crus-galli)
- Crabgrass (Digitaria spp.)
- Fall panicum (Panicum dichotomiflorum)
- Foxtail (Setaria spp.)
- Goosegrass (Eleusine indica)

BROADLEAF WEEDS CONTROLLED

- Common chickweed (Stellaria media)
- Hop clover (Trifolium procumbens)
- Henbit (Lamium amplexicaule)
- Mouseear chickweed (Cerastium vulgatum)
- Prostrate spurge (Euphorbia supina)
- Yellow wood sorrel (Oxalis stricta)



EPA Reg. No. 241-268

FOR USE IN LAWNS AND TURF

PROWL DG can be used for weed control in lawns and turf.

NOT REVIEWED
In Accordance with PR Notice 82-2.
Based on Draft Labeling Dated
7/5/84

TURFGRASS TYPES

PROWL DG should only be applied to well established lawns and turf. PROWL DG can be used on the following turfgrasses:

Bahia
Bermudagrass
Centi pedegrass

St. Augustinegrass
Tall fescue
Zoysiagrass

WEED SPECIES CONTROLLED

When applied as directed, PROWL DG will control the following weed species:

GRASSES CONTROLLED

Annual bluegrass (*Poa annua*)
Barnyardgrass (*Echinochloa crus-galli*)
Crabgrass (*Digitaria* spp.)
Fall panicum (*Panicum dichotomiflorum*)
Foxtail (*Setaria* spp.)
Goosegrass (*Eleusine indica*)

BROADLEAF WEEDS CONTROLLED

Common chickweed (*Stellaria media*)
Hop clover (*Trifolium procumbens*)
Henbit (*Lamium amplexicaule*)
Mouseear chickweed (*Cerastium vulgatum*)
Prostrate spurge (*Euphorbia supina*)
Yellow wood sorrel (*Oxalis stricta*)

APPLICATION INSTRUCTIONS

Add PROWL DG slowly to partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. MAINTAIN CONTINUOUS AGITATION WHILE ADDING PROWL DG AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

PROWL DG treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application.

APPLICATION TIMING

PROWL DG treatments will not control established weeds. Applications must be made prior to germination of weeds.

For control of annual bluegrass, chickweed and henbit, apply in late summer to early fall before weeds germinate.

For control of crabgrass and other annual weeds listed above, apply in the spring before weeds germinate.

Two applications per year may be made. Allow at least 4 months between applications.

APPLICATION RATE

Apply 4.5 lbs. of PROWL DG per acre (1.0 lb/10,000 sq. ft.).

NOTE: Use only on well established turfgrass with a dense and uniform stand. DO NOT use on newly seeded or sprigged turfgrass. DO NOT use on winter-overseeded turfgrass. DO NOT use on turfgrass severely thinned by winter stress.

PROWL provides effective residual weed control which may affect new turfgrass seedlings or sprigs. DO NOT reseed or winter overseed turfgrass within 4 growing months after application. DO NOT sprig turfgrass within 5 months after application.

DO NOT use on bentgrass or dichondra.



**Agricultural Division
Bio-Tech Products Department
Wayne, NJ 07470 ©1984**

PROWL® DG herbicide

SUPPLEMENTAL LABELING

EPA Reg. No. 241-268


FOR USE IN LAWNS AND TURF

PROWL DG can be used for weed control in lawns and turf.

TURFGRASS TYPES

PROWL DG should only be applied to well established lawns and turf. PROWL DG can be used on the following turfgrasses:

Bahia
Bermudagrass
Centipedegrass
St. Augustinegrass
Tall fescue
Zoysiagrass



WEED SPECIES CONTROLLED

When applied as directed, PROWL DG will control the following weed species:

GRASSES CONTROLLED

Annual bluegrass (Poa annua)
Barnyardgrass (Echinochloa crus-galli)
Crabgrass (Digitaria spp.)
Fall panicum (Panicum dichotomiflorum)
Foxtail (Setaria spp.)
Goosegrass (Eleusine indica)

BROADLEAF WEEDS CONTROLLED

Common chickweed (Stellaria media)
Hop clover (Trifolium procumbens)
Henbit (Lamium amplexicaule)
Mouseear chickweed (Cerastium vulgatum)
Prostrate spurge (Euphorbia supina)
Yellow wood sorrel (Oxalis stricta)

BEST AVAILABLE COPY

APPLICATION INSTRUCTIONS

Add PROWL DG slowly to partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. MAINTAIN CONTINUOUS AGITATION WHILE ADDING PROWL DG AND UNTIL SPRAYING IS COMPLETED. If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying.

Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. DO NOT apply when winds will cause drift.

PROWL DG treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application.

APPLICATION TIMING

PROWL DG treatments will not control established weeds. Applications must be made prior to germination of weeds.

For control of annual bluegrass, chickweed and henbit, apply in late summer to early fall before weeds germinate.

For control of crabgrass and other annual weeds listed above, apply in the spring before weeds germinate.

Two applications per year may be made. Allow at least 4 months between applications.

APPLICATION RATE

Apply 4.5 lbs of PROWL DG per acre (1.0 lb/10,000 sq. ft.).

NOTE: Use only on well established turfgrass with a dense and uniform stand. DO NOT use on newly seeded or sprigged turfgrass. DO NOT use on winter-overseeded turfgrass. DO NOT use on turfgrass severely thinned by winter stress.

PROWL provides effective residual weed control which may affect new turfgrass seedlings or sprigs. DO NOT reseed or winter overseed turfgrass within 4 growing months after application. DO NOT sprig turfgrass within 5 months after application.

DO NOT use on bentgrass or dichondra.

BEST AVAILABLE COPY



CYANAMID

Agricultural Division
New York Products Department
Wayne, N.J. 07178 • 1964