

EPA Reg. No. 241-268

ACCEPTED

MAR 19 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-268

FOR USE IN LAWNS AND TURF

PROWL DG can be used for weed control in lawns and turf.

TURFGRASS TYPES

PROWL DG should only be applied to well established lawns and turf.
PROWL DG can be used on the following turfgrasses:

Bahia	Fine fescue	St. Augustinegrass
Bermudagrass	Kentucky bluegrass	Tall fescue
Centipedegrass	Perennial ryegrass	Zoysiagrass

WEED SPECIES CONTROLLED

When applied as directed, PROWL DG will control the following weed species:

GRASSES CONTROLLED

Annual bluegrass (Poa annua)
Barnyardgrass (Echinochloa crus-galli)
Crabgrass (Digitaria spp.)
Fall panicum (Panicum dichotomiflorum)
Foxtail (Setaria spp.)
Goosegrass (Eleusine indica)

BROADLEAF WEEDS CONTROLLED

Common chickweed (Stellaria media)
Cudweed (Gnaphalium spp.)
Hop clover (Trifolium procumbens)
Henbit (Lamium amplexicaule)
Mouseear chickweed (Cerastium vulgatum)
Prostrate spurge (Euphorbia supina)
Yellow wood sorrel (Oxalis stricta)
Evening primrose (Oenothera biennis)

Add PROWL DG slowly to partially-filled tank (1/2 to 3/4 full) of water while agitating and then fill the remainder of the tank with water. **MAINTAIN CONTINUOUS AGITATION WHILE ADDING PROWL DG.** If the spray mixture is allowed to settle for any period of time, thorough agitation is essential to resuspend the mixture before spraying is resumed. Continue agitation while spraying as needed.

Apply with properly calibrated equipment in sufficient water per acre to provide uniform spray distribution. Low pressure (25-50 psi) sprayers are recommended. Nozzle screens must be no finer than 50 mesh. Avoid application when winds may cause drift.

PROWL DG treatments are most effective in controlling weeds when adequate rainfall or overhead irrigation is received within 7 days after application.

Avoid contact of spray solution with driveways, stone, wood or other porous surfaces. Rinse immediately to avoid staining.

COMPATABILITY

PROWL DG is compatible with most commonly used herbicides. When tank-mixing be sure to add PROWL DG to the partially filled tank first, mix thoroughly, and then add combination products to the mixture. **DO NOT** apply tank mix combinations unless previous experience indicates the mixture is effective and will not result in application problems or plant injury. **DO NOT** attempt to mix PROWL DG directly with liquid fertilizer. Premix one part of PROWL DG with two parts water and add this diluted mixture slowly into the tank while agitating.

GENERAL INFORMATION

Special Precautions

PROWL DG treatments will not control established weeds. Applications must be made prior to germination of weeds.

Two applications per year may be made. Allow at least 4 months between applications except where indicated under DIRECTIONS FOR USE.

Use only on well established turfgrass with a dense and uniform stand. **DO NOT** use on newly seeded or sprigged turfgrass. **DO NOT** use on winter-overseeded turfgrass. **DO NOT** use on turfgrass severely thinned by winter stress.

PROWL provides effective residual weed control which may affect new turfgrass seedlings or sprigs. **DO NOT** reseed or winter overseed turfgrass within 4 growing months after application. **DO NOT** sprig turfgrass within 5 months after application.

DO NOT use on bentgrass or dichondra maintained as desirable turf.

DIRECTIONS FOR USE-

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

TURF SPECIES	WEEDS CONTROLLED	RATES OF PROWL DG	COMMENTS
NORTHERN GRASSES: Fine fescue Kentucky bluegrass Perennial ryegrass Tall fescue	Barnyardgrass Crabgrass Fall panicum Foxtail Hop clover Prostrate spurge Yellow wood sorrel Cudweed	2.5 to 3.3 lbs per Acre or .9 to 1.2 oz. per 1000 ft ²	Apply in spring before weeds germinate.
	Goosegrass Evening primrose	5 lbs per Acre or 1.8 oz per 1000 ft ²	Apply in spring before weeds germinate. Repeat application if necessary in 8 weeks at 2.5 lbs per Acre or .9 oz per 1000 ft ²
	Annual bluegrass Common chickweed Mouseear chickweed Henbit	2.5 to 3.3 lbs per Acre or .9 to 1.2 oz. per 1000 ft ²	Apply in late summer to early fall, before weeds germinate.
SOUTHERN GRASSES: Bahia Bermudagrass Centipedegrass St. Augustinegrass Tall fescue Zoysiagrass	Barnyardgrass Crabgrass Fall panicum Foxtail Prostrate spurge Yellow wood sorrel Cudweed	2.9 to 3.3 lbs per Acre or 1.0 to 1.2 oz. per 1000 ft ²	Apply in spring before weeds germinate.
	Goosegrass Hop clover Evening primrose	4.5 to 5.5 lbs per Acre or 1.6 to 2.0 oz. per 1000 ft ²	
	Henbit Annual bluegrass Common chickweed Mouseear chickweed	2.5 to 3.3 lbs per Acre or .9 to 1.2 oz. per 1000 ft ²	Apply in late summer to early fall before weeds germinate.