

ASSERT!



herbicide



BEST AVAILABLE COPY

2

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER

Corrosive. Causes Irreversible Eye Damage. **DANGER! DO NOT** get in eyes. Wear goggles and gloves when handling. Avoid contact with skin and clothing. Avoid breathing vapor or spray mist. Use with adequate ventilation. Wash thoroughly after handling with soap and water. Remove contaminated clothing and wash separately from noncontaminated clothes before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ASSERT should be used only in accordance with recommendations on this label.

DO NOT apply this product through any type of irrigation system.

Observe all cautions and limitations on this label and on the labels of products used in combination with ASSERT. American Cyanamid Company will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by American Cyanamid. User assumes all risks associated with such non-recommended use. Keep container closed to avoid spills and contamination.

STORAGE AND DISPOSAL

Keep from freezing. **DO NOT STORE BELOW 40°F (5°C)**. Ice will form in the container at 5°F (-15°C). If ice occurs from prolonged storage at temperatures of 5°F (-15°C) or below, place container at room temperature until ice melts. Stability of ASSERT is not affected by freezing and thawing.

DO NOT contaminate water, food or feed by storage or disposal. Store ASSERT only in original container. **DO NOT** mix or store in unlined containers.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR 2.5 GALLONS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR BULK: Return empty container for reuse.

GENERAL INFORMATION

Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

EQUIPMENT APPLICATION INSTRUCTIONS

The spray equipment must be clean and properly calibrated before treatments are applied. For best results, use flat fan nozzles, and a pressure of

20 to 40 psi to achieve uniform spray distribution and minimize drift. A drift retardant agent may be added to the tank if needed. **DO NOT** use flood jet nozzles. Keep the by-pass line on or near the bottom of the tank to minimize foaming.

During aerial applications, a flagman should be located at each end of the field, or attach an automatic mechanical flagging unit to the aircraft to ensure uniform application. **DO NOT** overlap spray by aerial or ground application.

MIXING INSTRUCTIONS

A NON-IONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT MUST BE USED WITH ASSERT HERBICIDE.

ADD clean water to the spray tank one-half to three quarters full.

When using ASSERT alone, add ASSERT to the partially filled tank while agitating. When ASSERT is thoroughly mixed add the surfactant to the tank at a rate of 2 pints surfactant per 100 gallons of spray solution. Fill the remainder of tank with clean water while continuing agitation.

Crop oil can be added to an ASSERT application at a rate of 2 pints (1 quart) per acre of spray solution. A non-ionic surfactant plus crop oil is acceptable in tank mix with an ASSERT application.

When tank mixing ASSERT with a labeled tank mix partner, first fill the spray tank two thirds full with clean water. Add the partner herbicide(s) to the spray tank first while agitating the solution. When the herbicide solution is thoroughly mixed, add ASSERT to the tank. An antifoaming agent may be added if necessary. After the ASSERT and tank mix partners are thoroughly mixed, add the non-ionic surfactant at a rate of 2 pints per 100 gallons spray solution. Finally, crop oil and/or drift retardant should be added last. Complete filling the tank with water while continuing agitation. Maintain continuous agitation until spraying is completed.

HOW TO USE ASSERT PLUS SUN-IT II spray adjuvant in wheat and barley: SUN-IT II spray adjuvant may be used instead of a non-ionic surfactant or instead of a non-ionic surfactant plus crop oil in tank mix with an ASSERT application.

NOTE: When using other brands of crop oil, or petroleum or vegetable-based crop oils, a non-ionic surfactant must be included with ASSERT.

Use SUN-IT II at the rate of 1.5 to 2.0 pints per acre. Use the higher rate when weeds are at the maximum label size or under stress. SUN-IT II is recommended when weeds are under moisture or temperature stress. When using ASSERT with a non-phosphorous liquid fertilizer, follow instructions on the ASSERT label for the compatibility test of SUN-IT II. When tank mixing ASSERT with a labeled herbicide tank mix partner, determine whether SUN-IT II spray adjuvant (crop oil) is approved on that tank mix partner label.

APPLICATIONS WITH LIQUID FERTILIZERS

GENERAL

ASSERT can be applied with a non-phosphorus liquid fertilizer such as 28-0-0. Non-phosphorus liquid fertilizers can be applied with ASSERT alone or in combination with MCPA ester², 2,4-D ester², or Bromoxynil + MCPA ester² (Bronate[®]) to wheat and barley. Follow all ASSERT label recommendations regarding timing of application, special instructions and precautions. Apply the ASSERT/non-phosphorus liquid fertilizer combination with a minimum of 8 gallons per acre with ground equipment or a minimum of 5 gallons of spray solution per acre with aircraft.

NOTE: Herbicides can increase contact burn of fertilizers on plant foliage. Reduced gallons of fertilizer per acre may decrease leaf burn. **DO NOT** allow ASSERT to remain overnight in a liquid fertilizer solution.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the ASSERT/liquid fertilizer mixture.

BEST AVAILABLE COPY

LIQUID FERTILIZER COMPATIBILITY DETERMINATIONS

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of ASSERT alone or with other herbicides in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint of liquid fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant to one jar.
3. (a) When using ASSERT alone, add to each jar the correct amount of ASSERT as specified in the table below.
(b) When using ASSERT tank mixtures, first add the specified quantity of MCPA ester, 2,4-D ester, or Bromoxynil + MCPA ester and then add the correct amount of ASSERT.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
5. Determine compatibility.
 - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
 - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.
 - (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
 - (d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

Gallons of Liquid Fertilizer plus Water to be Applied per acre	Teaspoons of Specific Herbicide to be Added to 1 pint of Liquid Fertilizer Solution			
	ASSERT	MCPA ester	2,4-D ester	Bronate
5	3	2	2	2
10	1 1/2	1	1	1
15	3/4	1/2	1/2	1/2

SPRAYING INSTRUCTIONS

Uniformly apply the recommended ASSERT or ASSERT tank mixture in 8 or more gallons of water per acre with ground equipment or in 5 or more gallons of water per acre by aircraft. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop or a rotational crop could result. DO NOT allow spray to drift onto adjacent crops as injury may occur. For optimum weed control, ASSERT must absorb into plant leaves for 3 hours prior to overhead irrigation or rainfall.

IMPORTANT

DO NOT make more than one application of ASSERT per growing season. DO NOT graze treated fields or cut treated forage for silage or hay. Wheat and barley straw may be fed or used for bedding.

FOLLOW CROP RESTRICTIONS

The following rotational crops may be planted the season after applying ASSERT at recommended rates in wheat, barley and sunflowers:

Soybean	Corn	Edible beans	Sunflowers
Safflower	Wheat	Barley	

If ASSEN[®] and Glean[®], ASSERT and Ally[®], or ASSERT and Finites[®] are applied together in any type of tank mix during the same year, follow the most restrictive precautions, directions and recropping/rotation limitations that appear on the respective product labels.

DO NOT plant sugar beets for at least 20 months following an ASSERT application. DO NOT plant other rotational crops than those listed above for 15 months following an ASSERT application.

Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

WHEAT AND BARLEY

GENERAL INFORMATION

ASSERT is a selective herbicide for the postemergence control of wild oat, roughstalk bluegrass, interrupted windgrass, or grain broadleaved weeds in wheat and barley. Apply ASSERT postemergence in wheat (including durum) and barley from the 2 leaf stage of the crop, but before development of the first internode (jointing). DO NOT apply ASSERT when freezing temperatures are forecast.

Wild oats will be controlled with ASSERT applied when wild oats are in the 1 to true leaf stage.

If unfavorable growing conditions exist, ASSERT may cause slight discoloration and delayed growth of durum wheat. The crop will recover under normal growing conditions and yields will not be affected.

ASSERT alone or in combination can be applied by ground equipment or aircraft. Maximum weed control with ASSERT alone or in combination is obtained when temperature, moisture, fertility and cultural practices provide favorable conditions for active plant growth. Thorough uniform spray coverage is required to maximize weed control. For high populations of wild oats (in excess of 25 plants per square foot), use a minimum spray volume of 15 gallons per acre.

ASSERT rapidly inhibits growth of susceptible weeds; however, typical symptoms (discoloration) of dying weeds may not be noticeable until 2-4 weeks after application. Weeds hardened by cold weather or drought stress may not be fully controlled and regrowth may occur; increase spray solution (gallons per acre) and surfactant under these conditions.

RATES* AND TIMING FOR WEED CONTROL IN WHEAT AND BARLEY					
WEED	Application Leaf Stage	STATES			
		MIN. ND. & SD	MT & WY	So. ID. CO & UT	WA, OR. & No. ID.
		pints/acre			
GRASSES:					
WILD OATS	1 - 4	1.0 - 1.2	1.2 - 1.5	1.3 - 1.5	1.5 ^b
Roughstalk bluegrass	1 - 4	1.2	1.5	1.5	1.5
Interrupted windgrass	2 - 4	—	—	—	1.5
BROADLEAVES:					
Wild mustard	1 - 6	1.0	1.2	1.3	1.5
London rocket	1 - 6	1.0	1.2	1.3	1.5
Field pennycress	1 - 4	1.2	1.5	1.5	1.5
Flaxweed	1 - 4	1.2	1.5	1.5	1.5
Tansymustard	1 - 6	1.2	1.5	1.5	1.5
Catchweed bedstraw ^c	1 - 3	1.5	1.5	1.5	1.5
Tartary buckwheat ^c	1 - 3	1.5	1.5	1.5	1.5
Wild buckwheat ^c	1 - 3	1.5	1.5	1.5	1.5

* When two or more weed species are present, use the recommended rate to control the more difficult to control weed. When a rate range is possible, use the higher rate when weed density is high and/or weeds are large.

BEST AVAILABLE COPY

DISCLAIMER

The label instructions for the use of this product reflect the opinion on research and field use. The directions are believed to be reliable followed carefully. However, it is impossible to eliminate all risks associated with use of this product. Crop injury, ineffectiveness or consequences may result because of such factors as weather conditions, other materials, or the use of, or application of the product instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for loss resulting from use of this product in any manner not set forth or assumed all risks associated with the use of this product in specifically set forth on this label.

American Cyanamid Company warrants only that the material conforms to the chemical description on the label and is reasonably described when used in accordance with the directions on the label referred to above. CYANAMID DOES NOT MAKE OR AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT. IN NO CASE SHALL CYANAMID OR THE SELLER BE LIABLE FOR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

USES WITH OTHER PRODUCTS (TANK MIX)

If this product is used in combination with any other product except recommended in writing by American Cyanamid Company, then American Cyanamid Company shall have no liability for any loss, damage, or injury arising in any such combination not so specifically recommended. If used in combination with any other product recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not by the inclusion of the American Cyanamid Company in such combination in any event shall be limited to return of the amount of the purchase price of the product.

¹ Registered Trademark of American Cyanamid Company

² SUN-IT II is a trademark of AGSCO, Inc.

³ MCPA ester, 2,4-D ester, and Bromoxynil + MCPA ester are sold under various names.

⁴ Bromate is a trademark of Rhone-Poulenc.

⁵ Ally, Express, Finesse, Glan, Harmony, and Harmony Extra are trademarks of Monsanto and Co., Inc.

⁶ Curtaw M is a trademark of DowElanco.

⁷ Banvel is a trademark of Sandak Crop Protection Corp.

⁸ Dicamba is an active ingredient of various herbicides sold under various names.

⁹ In eastern Washington and Northern Idaho a rate of 1.3 to 1.5 pints/acre of ASSERT may be applied to spring wheat or barley growing under favorable conditions.

¹⁰ When ASSERT is applied up to the 3 leaf stage, plants stop growing, resulting in a non-competitive plant.

HERBICIDE COMBINATIONS

Mixtures of ASSERT and one or more of the following broadleaf herbicides may be tank mixed to obtain maximum wild oat and broadleaf control:

MCPA ester ²	Ally ⁴
2,4-D ester ²	Harmony ⁴
Bromoxynil + MCPA ester ² (Bromate ³)	Harmony Extra ⁴
Curtaw M ⁶	Express ⁴
Glan ⁴	Finesse ⁴

¹¹ E. Follow the most restrictive precautions, directions and recropping/rotation limitations that appear on the respective product labels. Tank mixtures must be applied prior to the development of the first internode (jointing) of the crop.

DO NOT tank mix ASSERT with 2,4-D ester unless the crop is fully tillered. DO NOT tank mix ASSERT with Banvel⁷, any product containing dicamba⁸, MCPA amine, or 2,4-D amine formulations. A waiting period of 4 days should be observed before applying any herbicide not listed as a tank mix partner.

ASSERT/AVENGE⁹ tank mix for wild oat control:

ASSERT herbicide may be tank mixed with AVENGE wild jet herbicide for the control of wild oats only. This combination may be used in the following states: Idaho, California, Arizona, Nevada, Oregon, Utah, Washington and Wyoming. Follow all varietal restrictions present on the AVENGE label. ASSERT should be mixed at a rate of 0.54 pints per acre with AVENGE at 2 pints per acre.

This mixture can be applied from the 2-5 true leaf stage of wild oats. DO NOT apply this tank mix when the wheat/barley flagleaf is exposed. The tank mix should be applied in 8 or more gallons of water per acre (gpa) by ground equipment and in 5 or more gpa by aircraft. For wild oat populations in excess of 25 plants per square foot, use a minimum spray volume of 15 gpa. Use a non-ionic surfactant containing at least 80% active ingredient at a rate of 2 pints per 100 gallons spray solution up to an application rate of 15 gpa. For application rates greater than 15 gpa consult the SURFACTANT ADDITION TABLE on the AVENGE label for surfactant requirements.

WEED SPECIES NAMES

Common Names	Scientific Name
Wild oats	<i>Avena fatua</i>
Roughstalk bluegrass	<i>Poa trivialis</i>
Interrupted windgrass	<i>Apera interrupta</i>
Wild mustard	<i>Brassica kaber</i>
London rocket	<i>Sisymbrium irio</i>
Field pennycress, Fanweed	<i>Thlaspi arvense</i>
Finweed	<i>Descurainia sophia</i>
Tansymustard	<i>Descurainia pinnata</i>
Catchweed bedstraw	<i>Galium aparine</i>
Tartary buckwheat	<i>Fagopyrum tataricum</i>
Wild buckwheat	<i>Polygonum convolvulus</i>

ACCEPTED

MAR 26 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 241-285

INTEGTM is protected by the following Patents: U.S.A. 4744161, 4680080, 4582572, 4675082, Canada 422,783, and others

BEST AVAILABLE COPY

ASSERT!
herbicide

In case of an emergency endangering life or property involving this product, call collect, day or night (201) 835-3100.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER

Corrosive, Causes Irreversible Eye Damage. Danger! DO NOT get in eyes. Wear goggles and gloves when handling. Avoid contact with skin and clothing. Avoid breathing vapor or spray mist. Use with adequate ventilation. Wash thoroughly after handling with soap and water. Remove contaminated clothing and wash separately from noncontaminated clothes before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ASSERT should be used only in accordance with recommendations on the leaflet label.

DO NOT apply this product through any type of irrigation system.

Observe all cautions and limitations on the leaflet label and on the labels of products used in combination with ASSERT. American Cyanamid Company will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by American Cyanamid. User assumes all risks associated with such non-recommended use. Keep container closed to avoid spills and contamination.

See Complete Directions For Use Inside Leaflet Label.

STORAGE AND DISPOSAL

Keep from freezing. DO NOT STORE BELOW 40°F (5°C). Ice will form in the container at 5°F (-15°C). If ice occurs from prolonged storage at temperatures of 5°F (-15°C) or below, place container at room temperature until ice melts. Stability of ASSERT is not affected by freezing and thawing.

DO NOT contaminate water, food or feed by storage or disposal. Store ASSERT only in original carton. DO NOT mix or store in unlabeled containers.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL FOR 2.5 GALLONS: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR BULK: Return empty container for reuse.

See Disclaimer Inside Leaflet Label.

BEST AVAILABLE COPY