

Label for the use of this product in Minnesota, North Dakota, and South Dakota is hereby approved for distribution and use in Minnesota, North Dakota, and South Dakota.

241-285

FOR USE IN SUNFLOWER
(Expires April 30, 1996)

**FOR DISTRIBUTION AND USE IN MINNESOTA,
NORTH DAKOTA, AND SOUTH DAKOTA**

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE ASSERT LABEL AND ON TANK MIX PARTNER LABELS. FOLLOW THE MOST RESTRICTIVE USE PRECAUTIONS, OR DIRECTIONS FOR APPLICATION THAT APPEAR ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF HERBICIDE APPLICATION. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

ASSERT herbicide is a selective herbicide for the postemergence control of wild mustard in sunflowers. APPLY ASSERT POSTEMERGENCE TO SUNFLOWERS NOT UNDER DROUGHT/HEAT STRESS, FROM THE 2 TO 8-LEAF STAGE OF GROWTH (SUNFLOWERS MUST BE LESS THAN 15 INCHES HIGH). Maximum weed control with ASSERT is obtained when temperature, moisture, fertility and cultural practices provide favorable conditions for active plant growth. UNIFORM SPRAY COVERAGE IS REQUIRED FOR MAXIMUM WEED CONTROL AND REDUCED INJURY POTENTIAL.

Before applying ASSERT to sunflowers, thoroughly rinse and empty the spray tank of any herbicide containing solution. Then, wash all lines prior to adding ASSERT to the tank. Herbicide residues remaining in the spray tank can injure sunflowers at extremely low concentrations.

A NON-IONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT MUST BE USED WITH ASSERT HERBICIDE. Refer to the ASSERT container

leaflet label for general use, mixing, application and follow crop instructions.

DO NOT tank mix insecticides with ASSERT in sunflowers.

WEED CONTROLLED

Wild mustard (*Brassica kaber*)

TIMING AND RATES OF APPLICATION

For optimum wild mustard control apply ASSERT at 0.6 to 0.8 pints per acre (9.6-12.8 fl. oz/acre) when the majority of wild mustard plants are in the rosette stage and prior to bloom.

ASSERT may be applied following application of a soil applied herbicide registered for use in sunflowers such as PROWL® herbicide, Amiben¹, Eptam², Treflan³, or trifluralin⁴.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF ASSERT ON SUNFLOWERS. Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of ASSERT and therefore, injury to sunflowers and rotational crops is always possible.

REFER TO THE ASSERT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.

¹Amiben is a trademark of Rhone-Poulenc

²Eptam is a trademark of Zeneca.

³Treflan is a trademark of DowFlanco.

⁴Trifluralin is sold under various brand names

®Registered Trademark of American Cyanamid Company

PE-13064 Rev. 2 9/20/94



CYANAMID

American Cyanamid Company
Agricultural Products Division
Specialty Products Department
Wayne, NJ 07470

©1994

BEST AVAILABLE COPY