


# ASSERT<sup>®</sup>

PULL HERE TO OPEN  
PRESS TO RESEAL

herbicide

7.312  
Inches



5.500 Inches

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS**

**DANGER**

Corrosive. Causes Irreversible Eye Damage. **DANGER! DO NOT get in eyes.** Avoid contact with skin and clothing. Avoid breathing vapor or spray mist. Use with adequate ventilation.

**Personal Protective Equipment (PPE):**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category B on an EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:**

- long-sleeved shirt and long pants
- chemical-resistant gloves, such as barrier-laminate or butyl rubber ≥ 14 mils
- shoes plus socks
- protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations:**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

**DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ASSERT should be used only in accordance with recommendations on this label.

**DO NOT** apply this product through any type of irrigation system.

Observe all cautions and limitations on this label and on the labels of products used in combination with ASSERT. American Cyanamid Company will not be responsible for losses or damages resulting from the use of this product in any manner not specifically recommended by American Cyanamid. User assumes all risks associated with such non-recommended use. Keep container closed to avoid spills and contamination.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical-resistant gloves, such as barrier-laminate or butyl rubber ≥ 14 mils
- shoes plus socks
- protective eyewear.

**STORAGE AND DISPOSAL**

Keep from freezing. **DO NOT STORE BELOW 40°F (5°C).** Ice will form in the container at 5°F (-15°C). If ice occurs from prolonged storage at temperatures of 5°F (-15°C) or below, place container at room temperature until ice melts. Stability of ASSERT is not affected by freezing and thawing.

**DO NOT** contaminate water, food or feed by storage or disposal. Store ASSERT only in original container. **DO NOT** mix or store in unlined containers.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL FOR 2.5 GALLONS:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONTAINER DISPOSAL FOR BULK:** Return empty container for reuse.

7.312  
Inches

5.469 Inches

### DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

**BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF ASSERT.** In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

### GENERAL INFORMATION

Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Following the use of this product and chemically related products with the same mode of action, naturally occurring biotypes\* of some of the weeds listed on this label cannot be effectively controlled by this and related products. This product should be tank-mixed or applied sequentially with an appropriate registered herbicide having a different mode of action to ensure control of resistant biotypes.

\* A weed biotype is a naturally occurring individual within a given species that has a slightly different, but distinct genetic makeup from other individuals.

See your local Cyanamid representative for more information.

### MIXING INSTRUCTIONS

**A NON-IONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT MUST BE USED WITH ASSERT HERBICIDE.**

**ADD clean water to the spray tank one-half to three quarters full.**

When using ASSERT alone, add ASSERT to the partially filled tank while agitating. When ASSERT is thoroughly mixed add the surfactant to the tank at a rate of 2 pints surfactant per 100 gallons of spray solution. Fill the remainder of tank with clean water while continuing agitation.

Crop oil can be added to an ASSERT application at a rate of 2 pints (1 quart) per acre of spray solution. When using crop oil as an adjuvant, it is required to also add an approved non-ionic surfactant (NIS) to the tank, unless the crop oil includes a NIS at a concentration which will give a final spray solution concentration of at least 0.25% NIS.

**DO NOT** use a surfactant that also acts as a buffering agent, or ASSERT may precipitate. If a precipitate forms due to incompatible tank-mix partners, contact your Cyanamid representative. If test of clean water supply indicates that a precipitate forms due to highly-buffered alkaline or hard water, precondition your water with sodium bisulfate. Contact your Cyanamid representative for instructions.

When tank mixing ASSERT with a labeled tank mix partner, first fill the spray tank two thirds full with clean water. Add the partner herbicide(s) to the spray tank first while agitating the solution. When the herbicide solution is thoroughly mixed, add ASSERT to the tank. An antifoaming agent may be added if necessary. After the ASSERT and tank mix partners are thoroughly mixed, add the non-ionic surfactant at a rate of 2 pints per 100 gallons spray solution. Finally, crop oil and/or drift retardant should be added last. Complete filling the tank with water while continuing agitation. Maintain continuous agitation until spraying is completed.

**HOW TO USE ASSERT PLUS SUN-IT II<sup>®</sup> spray adjuvant\*** In wheat and barley: SUN-IT II spray adjuvant may be used instead of a non-ionic surfactant or instead of a non-ionic surfactant plus crop oil in tank mix with an ASSERT application.

**NOTE:** When using other brands of crop oil, or petroleum or vegetable-based crop oils, a non-ionic surfactant must be included with ASSERT.

Use SUN-IT II at the rate of 1.5 to 2.0 pints per acre. Use the higher rate when weeds are at the maximum label size or under stress. SUN-IT II is recommended when weeds are under moisture or temperature stress. When using ASSERT with a non-phosphorous liquid fertilizer, follow instructions on the ASSERT label for the compatibility test of SUN-IT II. When tank mixing ASSERT with a labeled herbicide tank mix partner, determine whether SUN-IT II spray adjuvant (crop oil) is approved on that tank mix partner label.

\* **DO NOT** tank mix ASSERT with any product not registered in specific states of intended use.


### APPLICATIONS WITH LIQUID FERTILIZERS

#### GENERAL

ASSERT can be applied with a non-phosphorus liquid fertilizer such as 28-0-0. Non-phosphorus liquid fertilizers can be applied with ASSERT alone or in combination with MCPA ester<sup>2</sup>, 2,4-D ester<sup>2</sup>, or Bromoxynil + MCPA ester<sup>2</sup> (Bronate<sup>2</sup>) to wheat and barley. Follow all ASSERT label recommendations regarding timing of application, special instructions and precautions. Apply the ASSERT/non-phosphorus liquid fertilizer combination with a minimum of 8-10 gallons per acre with ground equipment or a minimum of 5 gallons of spray solution per acre with aircraft.

**NOTE:** Herbicides can increase contact burn of fertilizers on plant foliage. Reduced gallons of fertilizer per acre may decrease leaf burn. **DO NOT** allow ASSERT to remain overnight in a liquid fertilizer solution.

7.312  
Inches



All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the ASSERT/liquid fertilizer mixture.

### LIQUID FERTILIZER COMPATIBILITY DETERMINATIONS

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of ASSERT alone or with other herbicides in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint of liquid fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant to one jar.
3. (a) When using ASSERT alone, add to each jar the correct amount of ASSERT as specified in the table below.  
(b) When using ASSERT tank mixtures, first add the specified quantity of MCPA ester, 2,4-D ester, or Bromoxynil + MCPA ester and then add the correct amount of ASSERT.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
5. Determine compatibility.
  - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
  - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.
  - (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
  - (d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

Gallons of Liquid Fertilizer plus Water to be Applied per acre	Teaspoons of Specific Herbicide to be Added to 1 pint of Liquid Fertilizer Solution			
	ASSERT	MCPA ester	2,4-D ester	Bronate
5	3	2	2	2
10	1 1/2	1	1	1
15	3/4	1/2	1/2	1/2

### APPLICATION INSTRUCTIONS

The spray equipment must be clean and properly calibrated before treatments are applied. For best results, use flat fan nozzles, and a pressure of 20 to 40 psi to achieve uniform spray distribution and minimize drift. A drift retardant agent may be added to the tank if needed. DO NOT use flood jet nozzles. Keep the by-pass line on or near the bottom of the tank to minimize foaming.

### Ground Application

Uniformly apply the recommended ASSERT or ASSERT tank mixture in a minimum of 8-10 gallons of water per acre with ground equipment. **FOR HIGH POPULATIONS OF WILD OATS (IN EXCESS OF 25 PLANTS PER SQUARE FOOT), USE A MINIMUM SPRAY VOLUME OF 15 GALLONS PER ACRE BY GROUND.**

### Aerial Application

Uniformly apply with a minimum of 5 gallons of spray solution per acre by aircraft for all levels of wild oat populations.

During aerial applications, a flagman should be located at each end of the field, or attach an automatic mechanical flagging unit to the aircraft to ensure uniform application. DO NOT overlap spray by aerial or ground application.

### THOROUGH UNIFORM SPRAY COVERAGE IS REQUIRED TO MAXIMIZE WEED CONTROL.

Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping, or injury to the crop or a rotational crop could result. DO NOT allow spray to drift onto adjacent crops as injury may occur. For optimum weed control, ASSERT must absorb into plant leaves for 3 hours prior to overhead irrigation or rainfall.

DO NOT apply ASSERT when freezing temperatures have occurred or are forecast. Allow at least 2 days of non-freezing temperatures before and after ASSERT application or reduced weed control may occur.

For optimum weed control under cool conditions, it is recommended to apply ASSERT using the following additional instructions:

- 1) Use a minimum spray volume of 15 gallons per acre by ground or 5 gallons per acre by aerial application.
- 2) Use the maximum rate of ASSERT within a rate range.
- 3) Use SUN-IT II or non-ionic surfactant plus crop oil concentrate.

### IMPORTANT

DO NOT make more than one application of ASSERT per growing season. DO NOT graze treated fields or cut treated forage for silage or hay. Wheat and barley straw may be fed or used for bedding.

### FOLLOW CROP RESTRICTIONS



The following rotational crops may be planted the season after applying ASSERT at recommended rates in wheat, barley and sunflowers:

Soybean	Corn	Edible beans	Sunflowers
Safflower	Wheat	Barley	

If ASSERT and Glean<sup>®</sup>, ASSERT and Aly<sup>®</sup>, or ASSERT and Finesse<sup>®</sup> are applied together in any type of tank mix during the same year, follow the most restrictive precautions, directions and recropping/rotation limitations that appear on the respective product labels.

DO NOT plant sugar beets for at least 20 months following an ASSERT application. DO NOT plant other rotational crops than those listed above for 15 months following an ASSERT application.

Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.



### WHEAT AND BARLEY

#### GENERAL INFORMATION

ASSERT is a selective herbicide for the postemergence control of wild oats, roughstalk bluegrass, interrupted windgrass, and certain broadleaved weeds in wheat and barley. Apply ASSERT postemergence to wheat (including durum) and barley from the 2 leaf stage of the crop, but before development of the first internode (jointing). DO NOT apply ASSERT when freezing temperatures have occurred or are forecast. Allow at least 2 days of non-freezing temperatures before and after ASSERT application or reduced weed control may occur.

Wild oats will be controlled with ASSERT applied when wild oats are in the 1 to 4 true leaf stage.

If unfavorable growing conditions exist, ASSERT may cause slight discoloration and delayed growth of durum wheat. The crop will recover under normal growing conditions and yields will not be affected.

ASSERT alone or in combination can be applied by ground equipment or aircraft.

Maximum weed control with ASSERT alone or in combination is obtained when temperature, moisture, fertility and cultural practices provide favorable conditions for active plant growth. THOROUGH UNIFORM SPRAY COVERAGE IS REQUIRED TO MAXIMIZE WEED CONTROL. FOR HIGH POPULATIONS OF WILD OATS (IN EXCESS OF 25 PLANTS PER SQUARE FOOT), USE A MINIMUM SPRAY VOLUME OF 15 GALLONS PER ACRE BY GROUND OR 5 GALLONS PER ACRE BY AERIAL APPLICATION.

ASSERT rapidly inhibits growth of susceptible weeds; however, typical symptoms (discoloration) of dying weeds may not be noticeable until 2-4 weeks after application. Weeds hardened by cold weather or drought stress may not be fully controlled and regrowth may occur; increase spray solution (gallons per acre) and surfactant under these conditions.

RATES* AND TIMING FOR WEED CONTROL IN WHEAT AND BARLEY					
WEED	Application Leaf Stage	STATES			
		MN, ND, & SD	MT & WY	So. ID, CO & UT	WA, OR. & No. ID
		pints/acre			
<b>GRASSES:</b>					
WILD OATS	1 - 4	1.0 - 1.5 <sup>a</sup>	1.2 - 1.5	1.3 - 1.5	1.5 <sup>a</sup>
Roughstalk bluegrass	1 - 4	1.2	1.5	1.5	1.5
Interrupted windgrass	2 - 4	—	—	—	1.5
<b>BROADLEAVES:</b>					
Wild mustard	1 - 6	1.0	1.2	1.3	1.5
London rocket	1 - 6	1.0	1.2	1.3	1.5
Field pennycress	1 - 4	1.2	1.5	1.5	1.5
Flixweed	1 - 4	1.2	1.5	1.5	1.5
Tansymustard	1 - 6	1.2	1.5	1.5	1.5
Catchweed bedstraw <sup>b</sup>	1 - 3	1.5	1.5	1.5	1.5
Tartary buckwheat <sup>c</sup>	1 - 3	1.5	1.5	1.5	1.5
Wild buckwheat <sup>d</sup>	1 - 3	1.2	1.2	1.2	1.2


\* When two or more weed species are present, use the recommended rate to control the more difficult to control weed. When a rate range is possible, use the higher rate when weed density is high and/or weeds are large.

<sup>a</sup> In eastern Washington and Northern Idaho a rate of 1.3 to 1.5 pints/acre of ASSERT may be applied to spring wheat or barley growing under favorable conditions.

<sup>b</sup> When ASSERT is applied up to the 3 leaf stage, plants stop growing, resulting in a non-competitive plant.

<sup>c</sup> When ASSERT is applied at a rate of 1.2 pints/acre, it will provide suppression of wild buckwheat.

<sup>d</sup> In MN, ND, and SD when wild oats are 3 leaves or greater and populations are in excess of 25 plants per square foot, use the highest label rate of 1.5 pts per acre.



### HERBICIDE COMBINATIONS

Mixtures of ASSERT and one or more of the following broadleaf herbicides may be tank mixed to obtain maximum wild oat and broadleaf control:

- MCPA ester<sup>2</sup>
- 2,4-D ester<sup>2</sup>
- Bromoxynil + MCPA ester<sup>2</sup>
- ~~(Bronate<sup>3</sup>)~~
- Glean<sup>4</sup>
- Aily<sup>4</sup>
- Harmony<sup>4</sup>
- Harmony Extra<sup>4</sup>
- Express<sup>4</sup>
- Finesse<sup>4</sup>

NOTE: DO NOT tank mix ASSERT with any product not registered in specific states of intended use. Follow the most restrictive precautions, directions and recropping/rotation limitations that appear on the respective product labels. Tank mixtures must be applied prior to the development of the first internode (jointing) of the crop.

DO NOT tank mix ASSERT with 2,4-D ester unless the crop is fully tilled. DO NOT tank mix ASSERT with Banvel<sup>5</sup>, any product containing dicamba<sup>6</sup>, MCPA amine, or 2,4-D amine formulations. A waiting period of 4 days should be observed before applying any herbicide not listed as a tank mix partner.

**ASSERT®/AVENGE® tank mix for wild oat control:**  
ASSERT herbicide may be tank mixed with AVENGE wild oat herbicide for the control of wild oats only. Follow all varietal restrictions present on the AVENGE label. ASSERT should be mixed at a rate of 3/4 pints per acre with AVENGE at 2 pints per acre.

This mixture can be applied from the 2-5 true leaf stage of wild oats. DO NOT apply this tank mix when the wheat/barley flagleaf is exposed. The tank mix should be applied in 8-10 gallons of water per acre (gpa) by ground equipment and in 5 or more gpa by aircraft. **FOR WILD OAT POPULATIONS IN EXCESS OF 25 PLANTS PER SQUARE FOOT, USE A MINIMUM SPRAY VOLUME OF 15 GALLONS PER ACRE BY GROUND OR 5 GALLONS PER ACRE BY AERIAL APPLICATION.** Use a non-ionic surfactant containing at least 80% active ingredient at a rate of 2 pints per 100 gallons spray solution up to an application rate of 15 gpa. For application rates greater than 15 gpa consult the SURFACTANT ADDITION TABLE on the AVENGE label for surfactant requirements.

### WEED SPECIES NAMES

Common Names	Scientific Name
Wild oats	<i>Avena fatua</i>
Roughstalk bluegrass	<i>Poa trivialis</i>
Interrupted windgrass	<i>Apera interrupta</i>
Wild mustard	<i>Brassica kaber</i>
London rocket	<i>Sisymbrium irio</i>
Field pennycress, Fanweed	<i>Thlaspi arvense</i>
Flixweed	<i>Descurainia sophia</i>
Tansymustard	<i>Descurainia pinnata</i>
Catchweed bedstraw	<i>Galium aparine</i>
Tartary buckwheat	<i>Fagopyrum tataricum</i>
Wild buckwheat	<i>Polygonum convolvulus</i>

### USES WITH OTHER PRODUCTS (TANK MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company, then American Cyanamid Company shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in a combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use and in any event shall be limited to return of the amount of the purchase price of the product.

For additional information regarding the use of ASSERT herbicide call telephone number (800) 942-0500.

<sup>1</sup> Registered Trademark of American Cyanamid Company  
<sup>2</sup> SUN-IT II is a trademark of AGSCCO, Inc.  
<sup>3</sup> MCPA ester, 2,4-D ester, and Bromoxynil + MCPA ester are sold under various brand names.  
<sup>4</sup> Bronate is a trademark of Rhone Poulenc.  
<sup>5</sup> Aily, Express, Finesse, Glean, Harmony, and Harmony Extra are trademarks of E. I. Du Pont de Nemours and Co., Inc.  
<sup>6</sup> Banvel is a trademark of Sandax Crop Protection Corp.  
<sup>7</sup> Dicamba is an active ingredient of various herbicides sold under various brand names.



**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

**KILLS:** Bur Clover, Carpetweeds, Chickweeds, Creeping Charlie, Cudweed, Curly Dock, Dandelion, English Daisy, False Dandelion, Filaree, Florida Pusley, Heartleaf Drymary, Henbit, Oxalis, Pennywort, Plantains, Purslane, Red Sorrel, Sheep Sorrel, Spurges, Spurweed, Thistles, Toadflax, White Clover, Wild Carrot, Wild Geranium, Wild Onion.

**HOW TO USE:** Adjust spray nozzle to give a coarse spray. Aim at center of weed and spray to wet. Spray any time weeds are actively growing. Do not apply when rain or temperatures over 85° are expected within 24 hours. Hard to control weeds may require repeat application in 2-3 weeks. Weeds in newly seeded lawns may be sprayed after lawn has been mowed 3 times.

**IMPORTANT:** Do not use this product for controlling weeds in vegetable gardens, flower beds or around shrubs or ornamental plants. Temporary discoloration may occur on St. Augustine, Centipede, and Bentgrass lawns.

*[FOR 24 oz. size only] To Prepare Refill: Add 1 tablespoonful of WEED-B-GON Lawn Weed Killer, EPA Reg. No. 239-2342, to this empty container, then add 24 oz. (1-1/2 pints) tap water and use according to directions on this label.*

**STORAGE AND DISPOSAL**

*[For conventional trigger sprayers]*

**STORAGE:** Rotate nozzle to closed position.

*[For conventional trigger sprayers larger than 24 oz]* Place trigger sprayer under handle on container to keep sprayer above the level of contents

*[For proprietary sprayer and bottle]*

**STORAGE:** Flip down spout to close. No need to disconnect trigger sprayer. Close nozzle on trigger sprayer. Snap sprayer back in place.

Keep from freezing. Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area.

**DISPOSAL:** Do not reuse empty jug. Wrap container and put in trash. *[PR notice 84-1 Disposal copy for packages 1 gallon or less printed without refill instructions]*

**PRODUCT-**Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. **CONTAINER-**Do not reuse empty bottle except as specified in the directions to prepare refill. Rinse thoroughly before discarding in trash.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION:** Harmful if absorbed through skin. **APPLYING:** When applying this product wear goggles or face shield, long pants, long sleeved shirt, socks, shoes, and chemical resistant gloves. Wash nondisposable gloves thoroughly with soap and water before removing. Remove contaminated clothing and launder separately before reuse. Promptly and thoroughly wash hands and exposed skin with soap and water after using this product. Remove saturated clothing as soon as possible and shower. *[Child/Pet Icon]* Do not allow others such as children and pets on treatment area during application or to reenter treated areas until spray has dried. **FIRST AID:** If on skin, wash with plenty of soap and water. Note to Physicians: Emergency Information-call 1-800-454-2333. **ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Exercise caution when disposing of waste as groundwater contamination may result from careless handling or spills. Do not contaminate water when disposing of equipment washwaters.

**NOTICE:** Buyer assumes all responsibility for safety and use not in accordance with directions.

☐ Medical Information call 1-800-454-2333  
 Product Information call 1-800-225-2883

©Trademark of Monsanto Company  
 ©Monsanto Company 1994

Manufactured for  
 The SOLARIS Group of  
 Monsanto Company  
 P.O. Box 5008  
 San Ramon CA 94583-0808  
 Form RS789-A Product 5488  
 EPA Reg. No. 239-2499  
 EPA Est. 239-IA-3<sup>1</sup>, 58996-MO-1<sup>A</sup>  
 Superscript is first letter of lot number

Made in USA  
 UPC