

ASSERT®**Supplemental
Labeling****ACCEPTED**

herbicide

JAN 13 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-285

EPA Reg. No. 241-285

FOR USE IN WHEAT AND BARLEY

(Expires December 31, 2000)

P112

**TANK-MIXTURE APPLICATIONS OF ASSERT AND STARANE¹ EC,
STARANE PLUS SALVO², AND STARANE PLUS SWORD² FOR ENHANCED
WEED CONTROL IN WHEAT AND BARLEY**

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE ASSERT LEAFLET LABEL AND ON THE STARANE EC, STARANE PLUS SALVO, AND STARANE PLUS SWORD LABELS. SEE THE ASSERT LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. The use of ASSERT herbicide not consistent with this labeling may result in injury to crops. **DO NOT** apply this product through any type of irrigation system.

This labeling must be in the possession of the user at the time of herbicide application.

GENERAL INFORMATION

Starane EC, Starane Plus Salvo or Starane Plus Sword herbicide may be tank mixed with ASSERT herbicide for postemergence weed control in wheat and barley. Refer to the respective product labels for application rates and weeds controlled. Follow the most restrictive precautions, directions and recropping/rotation limitations that appear on the respective product labels. When tank-mixing herbicides, always follow the most restrictive label.

Apply this tank mix to wheat and barley **before** development of the first internode (jointing). **DO NOT** tank mix ASSERT with 2,4-D ester unless the crop is fully tillered.

USE RATE

Apply ASSERT at the rate of 1-1.5 pints/A.
See the Starane EC, Starane Plus Salvo or Starane Plus Sword labels for recommended rates.

MIXING INSTRUCTIONS

1. Fill spray tank one-half to two-thirds full with clean water.
2. Add Starane EC, Starane Plus Salvo or Starane Plus Sword herbicide to the spray tank while agitating the solution.

(see reverse side)

3. After the Starane EC, Starane Plus Salvo or Starane Plus Sword is thoroughly mixed, add ASSERT to the partially filled tank while continuing agitation
4. When the herbicide solution is thoroughly mixed, add the non-ionic surfactant (NIS) to the tank at a rate of 2 pints per 100 gallons of final spray solution.
5. Fill remainder of the tank with clean water.

SURFACTANTS/ADJUVANTS: Use only a non-ionic surfactant containing at least 80% active ingredient when using this tank mix. Apply the surfactant at a rate of 2 pints per 100 gallons of spray solution. **Do not use a crop oil concentrate or fertilizer as an adjuvant, or crop injury may result.**

Refer to the ASSERT product label for additional information on application instructions.

ROTATIONAL CROP INFORMATION

Refer to rotational crop restrictions listed in the ASSERT and Starane EC, Starane Plus Salvo or Starane Plus Sword product labels.

Always follow the most restrictive label.

Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.



8341

¹Trademark of Dow AgroSciences LLC
²Registered Trademark of Platte Chemical Co.
 © Registered Trademark of American Cyanamid Company
 PE-XXXXX 12/7/98 SD