

241-285
ASSERT[®]
herbicide

01/28/2000

1/3

Supplemental Labeling

EPA Reg. No. 241-285

FOR USE IN SUNFLOWER

FOR DISTRIBUTION AND USE IN MINNESOTA,
NORTH DAKOTA, AND SOUTH DAKOTA

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE ASSERT LABEL AND ON TANK MIX PARTNER LABELS. FOLLOW THE MOST RESTRICTIVE USE PRECAUTIONS, OR DIRECTIONS FOR APPLICATION THAT APPEAR ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF HERBICIDE APPLICATION. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

ASSERT herbicide is a selective herbicide for the postemergence control of wild mustard in sunflowers. APPLY ASSERT POSTEMERGENCE TO SUNFLOWERS NOT UNDER DROUGHT/HEAT STRESS, FROM THE 2 TO 8-LEAF STAGE OF GROWTH (SUNFLOWERS MUST BE LESS THAN 15 INCHES HIGH). Maximum weed control with ASSERT is obtained when temperature, moisture, fertility and cultural practices provide favorable conditions for active plant growth. UNIFORM SPRAY COVERAGE IS REQUIRED FOR MAXIMUM WEED CONTROL AND REDUCED INJURY POTENTIAL.

Before applying ASSERT to sunflowers, thoroughly rinse and empty the spray tank of any herbicide containing solution. Then, wash all lines prior to adding ASSERT to the tank. Herbicide residues remaining in the spray tank can injure sunflowers at extremely low concentrations.

A NON-IONIC SURFACTANT CONTAINING AT LEAST 80% ACTIVE INGREDIENT MUST BE USED WITH ASSERT HERBICIDE. Refer to the ASSERT container leaflet label for general use, mixing, application and follow crop instructions.


DO NOT tank mix insecticides or herbicides with ASSERT in sunflowers.

WEED CONTROLLED

Wild mustard (*Brassica kaber*)

TIMING AND RATES OF APPLICATION

For optimum wild mustard control, apply ASSERT at 0.6 to 0.8 pints per acre (9.6-12.8 fl. oz/acre) when the majority of wild mustard plants are in the rosette stage and prior to bloom.



(See other side)

CANOLA FOLLOW CROP

CLEARFIELD* (imidazolinone tolerant) varieties of canola, such as Pioneer 45A71 and Pioneer 46A72, may be planted as a rotational crop the season after an application of ASSERT herbicide at label rates on wheat, barley and sunflowers.

Non CLEARFIELD (non-imidazolinone tolerant) canola may be planted as a rotational crop the season after an application of ASSERT herbicide at the 1 pint rate on wheat, barley, and sunflowers in the counties of Pembina, Cavalier, Ramsey, Walsh, Towner and Rolette in North Dakota and the counties of Kittson, Marshall, Roseau, Pennington, and Red Lake in Minnesota.

The following rotational crops may be planted the season after applying ASSERT at recommended rates in wheat, barley, and sunflowers:

Barley	Corn	Edible Beans	Safflower
Sunflower	Soybeans	Wheat	

For other follow crops, follow instructions on the ASSERT leaflet label.

DISCLAIMER

THE USER ASSUMES ALL RISKS AND CONSEQUENCES ASSOCIATED WITH APPLICATIONS OF ASSERT ON SUNFLOWERS. Use of ASSERT herbicide in accordance with label directions is expected to result in normal growth of crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of ASSERT and therefore, injury to sunflowers and rotational crops is always possible.

REFER TO THE ASSERT LEAFLET LABEL FOR ADDITIONAL DISCLAIMER STATEMENTS.



* Trademarks of American Cyanamid Company

ASSERT[®]

herbicide

3/3
Supplemental
Labeling

EPA Reg. No. 241-285

FOR USE IN WHEAT, BARLEY AND SUNFLOWERS

OILSEED RAPE (CANOLA) FOLLOW CROPS

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE ASSERT LEAFLET LABEL AND ON TANK MIX PARTNER LABELS. FOLLOW THE MOST RESTRICTIVE USE PRECAUTIONS OR DIRECTIONS FOR APPLICATION THAT APPEAR ON ANY PRODUCT LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system.

This label must be in the possession of the user at the time of herbicide application.

CANOLA FOLLOW CROP

CLEARFIELD* (imidazolinone tolerant) varieties of canola, such as Pioneer 45A71 and Pioneer 46A72, may be planted as a rotational crop the season after an application of ASSERT herbicide at label rates on wheat, barley and sunflowers.

Non-CLEARFIELD (non-imidazolinone tolerant) canola may be planted as a rotational crop the season after an application of ASSERT herbicide at the 1 pint rate on wheat, barley, and sunflowers in the counties of Pembina, Cavalier, Ramsey, Walsh, Towner and Rolette in North Dakota and the counties of Kittson, Marshall, Roseau, Pennington, and Red Lake in Minnesota.

The following rotational crops may be planted the season after applying ASSERT at recommended rates in wheat, barley, and sunflowers:

Barley	Safflower	Sunflower
Corn	Soybeans	Wheat
Edible Beans		

For other follow crops, follow instructions on the ASSERT leaflet label.

If ASSERT is used on a field treated during the same year with Ally¹, Amber², Canvas¹, Express¹, Finesse¹, Glean¹, Harmony¹ Extra or Peak², follow the most restrictive precautions, directions and recropping/rotation limitations that appear on the respective product labels.


Excessive moisture, drought and/or cold temperatures may affect the performance of ASSERT as well as the residual effect on rotational crops. **DO NOT** plant rotational crops in spray overlap areas, as follow crop injury may occur. The end user assumes all risks and consequences resulting from sprayer overlaps from ground and aerial applications.



CYANAMID

American Cyanamid Company
Global Agricultural Products Division
Crop Protection Products Department
One Campus Drive
Parsippany, NJ 07054

©1999



¹ Trademarks of E. I. du Pont de Nemours and Co., Inc.

² Trademarks of Novartis, Inc.

®/*Trademarks of American Cyanamid Company

PE- 13105 Rev. 2 12/1/99