

SCEPTER® herbicide

Supplemental Labeling

EPA Reg. No. 241-289


FOR USE IN SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER LEAFLET LABEL, BEFORE USING.


SCEPTER may be applied in Minnesota, North Dakota, South Dakota, northern Michigan and Wisconsin. Refer to the SCEPTER leaflet label for all use directions.

USE AREA

SCEPTER can be applied only in the states or parts of states shaded in the following map:



® Registered Trademark of American Cyanamid Company



40633

SCEPTER<sup>(R)</sup> herbicide

Supplemental Labeling

EPA Reg. No. 241-289

FOR USE IN LOUISIANA SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER LEAFLET LABEL BEFORE USING.

EARLY PLANNED POSTEMERGENCE

Apply SCEPTER early postemergence to soybeans after crop emergence, but before weeds exceed the 2 leaf stage of growth. Apply at a broadcast rate of 2/3 pints per acre.

THE ADDITION OF A NONIONIC SURFACTANT OR CROP OIL CONCENTRATE IS REQUIRED. THE NONIONIC SURFACTANT SHOULD CONTAIN AT LEAST 80% ACTIVE INGREDIENT AND SURFACTANT SHOULD BE APPLIED AT A RATE OF 2 PINTS PER 100 GALLONS OF SPRAY MIXTURE. APPLY THE CROP OIL CONCENTRATE (COC) AT THE RATE STATED ON THE COC LABEL.

There must be an interval of at least 90 days between the last SCEPTER application and soybean harvest.

When applied as directed, SCEPTER early planned postemergence treatments will control the following weeds:

- |                          |                           |
|--------------------------|---------------------------|
| COCKLEBUR                | (Xanthium strumarium)     |
| PIGWEEED                 |                           |
| PALMER                   | (Amaranthus palmeri)      |
| REDROOT                  | (Amaranthus retroflexus)  |
| SMOOTH                   | (Amaranthus hybridus)     |
| SPINY                    | (Amaranthus spinosus)     |
| TALL WATERHEMP           | (Amaranthus tuberculatus) |
| RED RICE*                | (Oryza sativa)            |
| MEXICANWEED              | (Cyperonia castaniifolia) |
| ECLIPTA                  | (Eclipta alba)            |
| POINSETTIA               | (Euphorbia heterophylla)  |
| SMALLFLOWER MORNINGGLORY | (Jacquemontia tamnifolia) |
| SMARTWEED, PENNSYLVANIA  | (Polygonum pensylvanicum) |

\* For control of this species, the SCEPTER early planned postemergence application must follow the application of a PROWL, Treflan, trifluralin, Lasso or Dual. SCEPTER should be applied before the height of the red rice exceeds 3 inches.

(R) Registered Trademark of American Cyanamid Company

ACCEPTED

FEB - 5 1987

U.S. Dept. of Agriculture  
 Agricultural Chemicals Act  
 registered for the pesticide  
 registered on  
 EPA Reg. No. 241-289

 **CYANAMID**

American Cyanamid Company  
 Agricultural Division  
 Crop Protection Chemicals Department  
 Wayne, NJ 07470

