

EPA Reg. No. 241-289

FOR USE IN LOUISIANA SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER LEAFLET LABEL BEFORE USING.

EARLY PLANNED POSTEMERGENCE

Apply SCEPTER early postemergence to soybeans after crop emergence, but before weeds exceed the 2 leaf stage of growth. Apply at a broadcast rate of 2/3 pints per acre.

THE ADDITION OF A NONIONIC SURFACTANT OR CROP OIL CONCENTRATE IS REQUIRED. THE NONIONIC SURFACTANT SHOULD CONTAIN AT LEAST 80% ACTIVE INGREDIENT AND SURFACTANT SHOULD BE APPLIED AT A RATE OF 2 PINTS PER 100 GALLONS OF SPRAY MIXTURE. APPLY THE CROP OIL CONCENTRATE (COC) AT THE RATE STATED ON THE COC LABEL.


There must be an interval of at least 90 days between the last SCEPTER application and soybean harvest.

When applied as directed, SCEPTER early planned postemergence treatments will control the following weeds:

- | | |
|--------------------------|---------------------------|
| COCKLEBUR | (Xanthium strumarium) |
| FIGWEED | |
| PALMER | (Amaranthus palmeri) |
| REDROOT | (Amaranthus retroflexus) |
| SMOOTH | (Amaranthus hybridus) |
| SPINY | (Amaranthus spinosus) |
| TALL WATERHEMP | (Amaranthus tuberculatus) |
| RED RICE* | (Oryza sativa) |
| MEXICANWEED | (Caperonia castaniifolia) |
| ECLIPTA | (Eclipta alba) |
| POINSETTIA | (Euphorbia heterophylla) |
| SMALLFLOWER MORNINGGLORY | (Jacquemontia tamnifolia) |
| SMARTWEED, PENNSYLVANIA | (Polygonum pensylvanicum) |

* For control of this species, the SCEPTER early planned postemergence application must follow the application of a PROWL, Treflan, trifluralin, Lasso or Dual. SCEPTER should be applied before the height of the red rice exceeds 3 inches.

(R) Registered Trademark of American Cyanamid Company


 FEB-6 1967
 Agricultural Division
 Crop Protection Chemicals Department
 EPA Reg. No. 241-289


CYANAMID
 American Cyanamid Company
 Agricultural Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470

