

EPA Reg. No. 241-289

NOT REVIEWED
In Accordance with FR Notice 82-2.
Based on Draft Labeling Dated 12/17/87

FOR USE IN SOYBEANS

**SCEPTER® PLUS PROWL® PLUS COMMAND¹ PREPLANT
INCORPORATED TANK MIXTURE**

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
SCEPTER LEAFLET LABEL BEFORE USING**

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
PROWL LEAFLET LABEL BEFORE USING**

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
COMMAND LEAFLET LABEL BEFORE USING**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The SCEPTER plus PROWL plus Command 4 EC tank mixture may be used when heavy infestations of velvetleaf are anticipated. Uniformly apply the tank mixture with properly calibrated ground equipment in 10 to 40 gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. NOTE: The use of an agriculturally approved drift reducing agent is required at finished spray volumes of 10 to 15 gallons per acre.

Apply SCEPTER plus PROWL plus Command 4 EC to the soil surface and uniformly incorporate to a depth of 1 to 2 inches. Incorporate within 3 hours of application. DO NOT apply aerially or through irrigation equipment. DO NOT apply this tank mixture to overly moist or wet soils.

SPECIAL PRECAUTION

Command 4 EC is a volatile compound. Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of some plants. Prior to using Command, read and strictly follow all precautions and application instructions on the Command 4 EC label

WEED SPECIES CONTROLLED

The SCEPTER plus PROWL plus Command tank mixture controls heavy infestations of velvetleaf as well as all of the weeds listed on the SCEPTER leaflet label for combination preplant incorporated treatments of SCEPTER plus PROWL.

¹Registered Trademark of American Cyanamid Company
Command 4 EC is a trademark of FMC Corporation

Less than 3% Organic Matter			
Coarse	3/4	1 to 1 1/2	1/4 to 1/2
Medium	3/4	1 1/2 to 2	1/4 to 1/2
Fine	3/4	1 1/2 to 2	1/4 to 1/2
More than 3% Organic Matter			
Coarse	3/4	1 1/2	1/4 to 1/2
Medium	3/4	1 1/2 to 2	1/4 to 1/2
Fine	3/4	2 to 2 1/2	1/4 to 1/2

When heavy grass populations are anticipated, apply PROWL at the high rate for each soil texture.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying the SCEPTER plus PROWL plus Command tank mixture at recommended rates in soybeans:

1. Nine months after last SCEPTER plus PROWL plus Command tank mix application:

Rice

2. Eleven months after last SCEPTER plus PROWL plus Command tank mix application:

Corn* Dry Beans Peanuts
Grain Sorghum Snap Beans Tobacco

3. Eighteen months after last SCEPTER plus PROWL plus Command tank mix application:

Cotton
Small Grains

* Do Not plant seed corn in the spring of the year following the application of SCEPTER plus PROWL plus Command tank mixture

Only rotational crops harvested at maturity may be used for feed or food.

DO NOT plant other rotational crops prior to 18 months after last SCEPTER plus PROWL plus Command tank mix application.

DO NOT apply Classic², Canopy², Gemini², Preview², or Lorox Plus² the same year as SCEPTER plus PROWL plus Command tank mix application or injury to follow crops may occur.


In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.

² Classic, Canopy, Gemini, Preview, and Lorox Plus are trademarks of E. I. du Pont de Nemours and Company



EPA Reg. No. 241-289

NOT REVIEWED
In Accordance with PR Notice 82-2.
Based on Draft Labeling Dated 12/17/87

FOR USE IN SOYBEANS

SCEPTER® PLUS TRIFLURALIN PLUS COMMAND¹ PREPLANT¹ INCORPORATED TANK MIXTURE

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
SCEPTER LEAFLET LABEL BEFORE USING**

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
TRIFLURALIN LEAFLET LABEL BEFORE USING**

**OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
COMMAND LEAFLET LABEL BEFORE USING**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The SCEPTER plus Trifluralin plus Command 4 EC tank mixture may be used when heavy infestations of velvetleaf are anticipated. Uniformly apply the tank mixture with properly calibrated ground equipment in 10 to 40 gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. NOTE: The use of an agriculturally approved drift reducing agent is required at finished spray volumes of 10 to 15 gallons per acre.

Apply SCEPTER plus trifluralin plus Command 4 EC to the soil surface and uniformly incorporate to a depth of 2 inches. Incorporate within 3 hours of application. DO NOT apply aerially or through irrigation equipment. DO NOT apply this tank mixture to overly moist or wet soils.

SPECIAL PRECAUTION

Command 4 EC is a volatile compound. Off-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of some plants. Prior to using Command, read and strictly follow all precautions and application instructions on the Command 4 EC label.

WEED SPECIES CONTROLLED

The SCEPTER plus trifluralin plus Command tank mixture controls heavy infestations of velvetleaf as well as all of the weeds listed on the SCEPTER leaflet label for combination preplant incorporated treatments of SCEPTER plus trifluralin.

¹Registered Trademark of American Cyanamid Company
²Command 4 EC is a trademark of FMC Corporation