

FOR USE IN SOYBEANS

NOT REVIEWED
 In Accordance with EPA Not 108 84-2a
 Based on Draft Registration Dated 12/1/76

Observe all Precautionary Statements in the SCEPTER® Leaflet label before using.

PU 25
 241-289
 pg. 1 of 2

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

The following rotational crops may be planted after applying SCEPTER at recommended rates in soybeans:

1. Wheat may be planted in the fall, four months after last SCEPTER application in soybeans. Rice may be planted in the spring of the year following SCEPTER application (See Northern Use Area Restrictions*)

2. Eleven months after last SCEPTER application:

Barley (See Northern Use Area Restrictions*)
 Oats (See Northern Use Area Restrictions*)
 Field Corn (See Northern Use Area* or Southern Use Area Restrictions**)
 Edible beans
 Grains sorghum
 Peanuts
 Tobacco

*NORTHERN USE AREA RESTRICTIONS:

For the Northern Use Area as defined in the USE AREA section of the label; field corn, wheat, barley or oats may be planted as a rotational crop in the fall or in the spring of the year following SCEPTER application ONLY if SCEPTER was applied as a post-emergence application at 1/3 pint/acre AND at least 10 inches of rainfall or irrigation has been received from the date of application through October of the year of application.

**SOUTHERN USE AREA RESTRICTIONS:

For the Southern Use Area as defined in the USE AREA section of the label; field corn may be planted in the spring of the year following SCEPTER application unless extreme drought conditions develop and less than 15 inches (10 inches for 1/3 pint/acre post-emergence applications) of rainfall or irrigation is received from the date of application through October of the year of application. For eastern Oklahoma, Arkansas, the bootheel of Missouri, Tennessee, North Carolina and states south; field corn may be planted in the spring of the year following SCEPTER application unless extreme drought conditions develop and less than 15 inches (10 inches for 1/3 pint/acre post-emergence applications) of rainfall or irrigation is received within 6 months following the date of application.

*Registered Trademarks of American Cyanamid Company
 PE-12088 3/23/76

Only rotational crops harvested at maturity may be used for feed or food. DO NOT plant other rotational crops prior to 18 months after last SCEPTER application.


DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Gemini, Lorox Plus, Preview, etc.) or imazethapyr (e.g. PURSUIT® or PURSUIT® PLUS) the same year as SCEPTER or injury to follow crops may occur.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches. DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

*Classic, Canopy, Gemini, Lorox Plus and Preview are trademarks of E. I. duPont de Nemours and Company.

USE AREA

SCEPTER can be applied only in the states or parts of states shaded in the following map:



 = Northern Use Area (includes: Nebraska, east of U.S. 81, north of U.S. 34; South Dakota, east of U.S. 81, Illinois, north of S.R. 116 for area west of Peoria; north of U.S. 24 for area east of Peoria, Indiana, north of U.S. 24 for area west of Huntington, north of U.S. 224, for area east of Huntington; Ohio, north of U.S. 224; and all other states or portions of states shaded in the Northern Use Area Map.)

 = Southern Use Area (includes Kansas, east of U.S. 81; southern portions of Illinois, Indiana, Ohio and eastern Nebraska not covered by Northern Use Area description; and all other states or portions of states shaded in the Southern Use Area map.)

This label must be in the possession of the user at the time of herbicide application.

ACCEPTED

MAY - 3, 1990

Under Federal Insecticide, Fungicide, and
 Rodenticide Act, as amended, for use
 pesticide registration under
 EPA Reg. No. 241-289



CYANAMID

American Cyanamid Company
 Agricultural Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470

EPA Reg. No. 241-289

FOR USE IN LOUISIANA SOYBEANS

EARLY PLANNED POSTEMERGENCE

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SCEPTER LABEL. See the SCEPTER label for mixing, spraying, application information, weed species controlled and fellow crop restrictions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label should be in the possession of the user at the time of pesticide application.

DO NOT apply this product through any type of irrigation system.



USE RECOMMENDATIONS

Apply SCEPTER herbicide early postemergence to soybeans after crop emergence, but before weeds exceed the 2 leaf stage of growth. Apply at a broadcast rate of $\frac{2}{3}$ pints per acre. THE ADDITION OF A NONIONIC SURFACTANT OR CROP OIL CONCENTRATE IS REQUIRED. THE NONIONIC SURFACTANT SHOULD CONTAIN AT LEAST 80% ACTIVE INGREDIENT, AND THE SURFACTANT SHOULD BE MIXED AT THE RATE OF 2 PINTS PER 100 GALLONS OF SPRAY MIXTURE. APPLY THE CROP OIL CONCENTRATE (COC) AT THE RATE STATED ON THE COC LABEL.

There must be an interval of at least 90 days between the last SCEPTER application and soybean harvest.

When applied as directed, SCEPTER herbicide early planned postemergence treatments will control the following weeds:

Alligatorweed	(<u>Alternanthera philoxeroides</u>)
Rice Flatsedge	(<u>Cyperus iria</u>)
Dayflower	(<u>Commelina communis</u>)
Redweed	(<u>Melochia corchorifolia</u>)
Texasweed	(<u>Cyperonia palustris</u>)



 **CYANAMID**
American Cyanamid Company
Agriculture Division
Crop Protection Chemical Department
Wayne NJ 07470

2/9/89