

EPA Reg. No. 241-289

PM25
241-289
pg. 193

FOR USE IN SOYBEANS

SCEPTER PREPLANT INCORPORATED OR PREEMERGENCE APPLICATION FOLLOWED BY CLASSIC¹ POSTEMERGENCE FOR CONTROL OF SICKLEPOD

For distribution and use only in Alabama, Arkansas, Florida, Georgia,
Louisiana, Mississippi, North Carolina, South Carolina, Tennessee and Texas.

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER AND CLASSIC LABELS BEFORE
USING. SEE THE SCEPTER AND CLASSIC LABELS FOR SPRAYING AND APPLICATION INFORMATION.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

Apply SCEPTER at 2/3 pint per acre preplant incorporated or preemergence in soybeans. A single postemer-
gence application of Classic at a rate of 1/2 to 3/4 ounce per acre may be applied to control sicklepod.

Refer to the Classic label for weed height limitations. Use the high rate of Classic if sicklepod seedlings are taller
than two inches. A cultivation approximately 14 days after Classic application may be required to control sicklepod
escapes.

NOTE: If replanting is necessary in a field previously treated with SCEPTER, the field may be replanted to
soybeans. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of
SCEPTER.

¹Classic is a Trademark of E.I. du Pont de Nemours and Company
²Registered Trademark of American Cyanamid Company

1. Soybeans may be planted anytime.

2. Fifteen months after last herbicide application:

Barley	Grain sorghum	Rice
Edible beans	Oats	Tobacco
Field corn	Peanuts	Wheat

3. Sugarbeets may be planted 26 months after last herbicide application.

4. All other crops, including cotton, may be planted eighteen months after the last herbicide application, unless specified otherwise in the PRECAUTIONS section of the Classic label.

Use of SCEPTER herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

ONLY rotational crops harvested at maturity may be used for feed or food.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label must be in the possession of the user at the time of herbicide application.

1000000000
1000000000
1000000000

JUL 25, 1990

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
25 U.S.C. 133, it is hereby
declared that the use of this
product is necessary to protect
the public health, safety, and
interests.


24-789


PE-12137 3/13/90



CYANAMID

American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07479 21990





U.S.A.