

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

PM 25 241-289 2 192

JUN 5 1992

Mr. Mark Galley
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08543

Dear Mr. Galley:

Subject: Supplemental Labeling
Planting of Field Corn at 11 Months After Use
SCEPTER® HERBICIDE
EPA Reg. No. 241-289
Your letter dated March 12, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:6/5/92:3056800

CONCURRENCES

SYMBOL	SURNAME	DATE						
	<i>ed</i>							

EPA Reg. No. 241-289

FOR USE IN SOYBEANS

SCEPTER ROTATIONAL CROP RESTRICTIONS

FOR DISTRIBUTION AND USE ONLY IN SCEPTER USE REGION 1 AND USE REGION 2

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER LABEL BEFORE USING. SEE THE SCEPTER LABEL FOR SPRAYING AND APPLICATION INFORMATION, WEEDS CONTROLLED, AND USE RATES. WHEN TANK MIXING, ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label should be in the possession of the user at the time of pesticide application. DO NOT apply this product through any type of irrigation system.

ROTATIONAL CROP RESTRICTIONS

When a single SCEPTER application at 2/3 pint per acre is made preplant incorporated or preemergence to soybeans, field corn may be planted as a rotational crop 11 months after herbicide application if one of the following criteria is met:

1. If at least 15 inches of rainfall or irrigation is received from two weeks prior to the last application through November 15 of the same year (within 6 months following the date of the last application in Use Region 1), or:
2. If a soil sample taken before field corn planting and in accordance with the guidelines set forth in "Soil Sampling for Determination of Residues" and analyzed at an American Cyanamid approved laboratory indicates that less than 10 parts per billion of imazaquin are present in the soil. Contact your chemical dealer or American Cyanamid representative for a copy of the soil sampling guidelines and a list of approved analytical laboratories.

If neither of the above requirements is met, only field corn hybrids which are classified as tolerant or resistant to SCEPTER and/or other imidazolinone herbicides may be planted in the spring of the year following a SCEPTER application. Contact your chemical dealer, seed supplier, or American Cyanamid to obtain a list of field corn hybrids. Other field corn hybrids may be planted 18 months after SCEPTER application.

Consult the SCEPTER label for rotational crop restrictions involving other crops, other SCEPTER application options, sequential herbicide applications, or tank mixtures with other herbicides.

Use of SCEPTER herbicide in accordance with directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of this product and therefore, rotational crop injury is always possible.

*Registered Trademark of American Cyanamid Company
PB-12235 2/25/92 SD

