

JUL 30 1993

Ms. Desiree L. Little
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08543

Dear Ms. Little:

Subject: Final Printed Labels
SCEPTER® HERBICIDE
EPA Reg. No. 241-289
Your submission dated July 16, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR part 158, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:7/30/93:3056800

PH 25 241-289
EPA Reg. No. 241-289

2013

FOR USE IN SOYBEANS

SCEPTER PLUS PURSUIT® HERBICIDE FOR SUPPRESSION OF VOLUNTEER CORN

FOR USE IN SCEPTER USE REGION 2 AND USE REGION 3
(EXCLUDING IOWA, MINNESOTA AND WISCONSIN)

THIS LABEL EXPIRES DECEMBER 31, 1993

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER AND PURSUIT LABELS BEFORE USING. SEE THE PURSUIT LABEL FOR HERBICIDE APPLICATION INFORMATION AND WEEDS CONTROLLED. WHEN TANK MIXING, ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

GENERAL INFORMATION

Apply SCEPTER at 2.0 fluid ounces per acre in combination with PURSUIT at 4.0 fluid ounces per acre postemergence for suppression of volunteer corn.

Refer to the SCEPTER leaflet label for a listing of those states included in USE REGION 2 and the approved use area of USE REGION 3.

WEEDS SUPPRESSED - VOLUNTEER CORN

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

This tank-mixture will provide suppression of volunteer corn in soybeans in addition to those weeds controlled by a postemergence application of PURSUIT. Refer to the PURSUIT label for a listing of weeds controlled by a postemergence application of PURSUIT.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

Follow all MIXING, SPRAYING and APPLICATION instructions in the PURSUIT leaflet label.

A postemergence application of SCEPTER will NOT suppress those field corn hybrids which possess tolerance or resistance to SCEPTER or other imidazolinone herbicides (i.e. PURSUIT).

USE RATE AND APPLICATION INSTRUCTIONS

Apply SCEPTER at 2.0 fluid ounces per acre in combination with PURSUIT at 4.0 fluid ounces per acre postemergence. Apply before the corn is ten inches tall.

ROTATIONAL CROP RESTRICTIONS

SCEPTER USE REGION 2: Follow all rotational crop restrictions on the PURSUIT and SCEPTER leaflet labels for that area designated as SCEPTER USE REGION 2 on the SCEPTER leaflet label.


USE AREA

This tank-mix application of SCEPTER and PURSUIT may be used in Region 2 and Region 3 (excluding Iowa, Minnesota and Wisconsin) as described in the USE AREA section of the SCEPTER label.

FOR THIS SUPPLEMENTAL LABEL ONLY; THE ENTIRE STATE OF ILLINOIS MAY FOLLOW THE ROTATIONAL CROP RESTRICTION GUIDELINES FOR SCEPTER USE REGION 2.

®Registered Trademark of American Cyanamid Company

(See other side)



1. Four months after SCEPTER plus PURSUIT application:
Soybeans
2. Four months after SCEPTER plus PURSUIT application:
Wheat
3. Eleven months after SCEPTER plus PURSUIT application:
Field corn (resistant)*
4. Eighteen months after SCEPTER plus PURSUIT application:
- | | |
|------------|------------|
| Alfalfa | Popcorn |
| Cotton | Safflower |
| Field corn | Sorghum |
| Lettuce | Sweet corn |
| Oats | |
5. Twenty six months after SCEPTER plus PURSUIT application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after SCEPTER plus PURSUIT application:
Sugar beets

Other field corn hybrids may be planted 10 months after application of this tank-mixture.

NOTE: Availability of imidazolinone resistant field corn seed cannot be assured.

Use of SCEPTER and PURSUIT herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of these products and, therefore, rotational crop injury is always possible.

EXCEPT as stated above, **DO NOT** use this tank mixture in the same year as an application of SCEPTER, SQUADRON®, TRI-SCEPT®, SCEPTER® O.T.®, PURSUIT, or PURSUIT PLUS®. Also, **DO NOT** use the above tank mixture in the same year as an application of a product containing chlorimuron ethyl (e.g. Classic¹, Canopy¹, Gemini¹, Lorox Plus¹, Preview¹, etc.)

ACCEPTED
with **COMMENTS**
in EPA Letter Dated:

JUL 30 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

241-289

*Registered Trademark of American Cyanamid Company
¹Trademark of E. I. duPont de Nemours & Company, Inc.
PE-0599 4/5/93 SD



CYANAMID

American Cyanamid Company
Agricultural Products Division
Crop Protection & Chemicals Department
Wayne, NJ 07433 ©1993