

FEB 2 1994

Mrs. Desiree L. Little  
American Cyanamid Co.  
P.O. Box 400  
Princeton, NJ 08540

Dear Mrs. Little:

Subject: For Use on Soybeans  
Alternate Tradename  
SCEPTER® Herbicide  
EPA Reg. No. 241-289  
Your submission dated Jan. 20, 1994

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor  
Product Manager 25  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

Enclosure  
RD:RTaylor:eja:93:305-6800

**BEST AVAILABLE COPY**

HERBICIDE 1

FOR USE IN SOYBEANS

ACTIVE INGREDIENT:

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)* .....	17.3%
--	-------

INERT INGREDIENTS** .....	82.7%
---------------------------	-------

TOTAL .....	100.0%
-------------	--------

\*Equivalent to 16.4% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid.

\*\*Contains Petroleum Distillates

(1 gallon contains 1.5 pounds of active ingredient as the free acid)

EPA Reg. No. 241-289

EPA Est. No. 241-MO-1

KEEP OUT OF REACH OF CHILDREN

CAUTION!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Flush eyes with plenty of water.

In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

See Additional Precautionary Statements Inside  
See Directions For Use Inside

AMERICAN CYANAMID COMPANY  
AGRICULTURAL PRODUCTS DIVISION  
CROP PROTECTION CHEMICALS DEPARTMENT  
WAYNE, NJ 07470 ©1993

Net Contents:

~~1000 ml~~  
~~1.0 L~~  
EPA Reg. No. 241-289

FEB 2 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
registered for the pesticide  
EPA Reg. No.  
241-289

**BEST AVAILABLE COPY**

TABLE OF CONTENTS

	Page
INGREDIENTS .....	cover
STATEMENT OF PRACTICAL TREATMENT .....	cover
PRECAUTIONARY STATEMENTS .....	3
STORAGE AND DISPOSAL .....	4
DISCLAIMER .....	5
GENERAL INFORMATION .....	6
USE AREA .....	8
WEEDS CONTROLLED (Soil-Applied Treatments).....	9
WEEDS CONTROLLED (Postemergence Treatments) .....	12
DIRECTIONS FOR CONVENTIONAL, MINIMUM, AND NO-TILL APPLICATIONS.....	13
APPLICATION RATE.....	13
SOIL APPLICATIONS .....	13
PREPLANT INCORPORATED APPLICATIONS .....	13
PREEMERGENCE APPLICATIONS (including no-till) .....	13
POSTEMERGENCE APPLICATIONS .....	14
HERBICIDE COMBINATIONS .....	15
WEEDS CONTROLLED (Combination Treatments).....	16
HERBICIDE 1 Plus PROWL® 3.3EC Tank Mixture.....	17
HERBICIDE 1 Plus Trifluralin.....	18
HERBICIDE 1 Plus Lasso <sup>1</sup> .....	19
HERBICIDE 1 Plus Dual <sup>2</sup> .....	20
HERBICIDE 1 Plus PROWL 3.3EC or Trifluralin Plus Command <sup>3</sup> 4EC Tank Mixture .....	21
HERBICIDE 1 Plus Metribuzin Tank Mixture .....	22
Tank Mixtures with 2,4-D.....	23
Tank Mixtures with 2,4-DB .....	23
Tank Mixtures with Bronco <sup>1</sup> .....	23
Tank Mixtures with Roundup <sup>1</sup> .....	24
Tank Mixtures with Gramoxone Extra <sup>4</sup> .....	24
MIXING INSTRUCTIONS .....	25
SPRAYING INSTRUCTIONS .....	25
APPLICATIONS WITH FERTILIZER.....	27
APPLICATIONS WITH LIQUID FERTILIZER.....	27
APPLICATIONS WITH DRY BULK FERTILIZER .....	29
ROTATIONAL CROP RESTRICTIONS.....	31
WEED SCIENTIFIC NAMES.....	34

<sup>1</sup>Registered trademarks of Monsanto Agricultural Products Company.

<sup>2</sup>Registered Trademark of Ciba-Geigy Corporation.

<sup>3</sup>Registered Trademark of FMC Corporation.

<sup>4</sup>Registered Trademark of Zeneca, Inc.

®Registered Trademark of American Cyanamid Company

**BEST AVAILABLE COPY**

**HERBICIDE 1  
PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**

**CAUTION!**

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

**Personal Protective Equipment (PPE):**

Applicators and other handlers must wear:

- \* Long-sleeved shirt and long pants
- \* Waterproof gloves
- \* Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations:**  
Users should:

- \* Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- \* Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.


**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with HERBICIDE 1. Do not use HERBICIDE 1 other than in accordance with the instructions set forth on this label. The use of HERBICIDE 1 not consistent with this label may result in injury to crops. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

**BEST AVAILABLE COPY**



DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- \* Coveralls
- \* Waterproof gloves
- \* Shoes plus socks

**STORAGE AND DISPOSAL**

**PROHIBITIONS:**

**KEEP FROM FREEZING**

DO NOT store below 32°F.

DO NOT contaminate water, food, or feed by storage or disposal of wastes.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

**BEST AVAILABLE COPY**

**CONTAINER DISPOSAL FOR MINI BULK:** HERBICIDE 1 is sold in System 60™ returnable mini-bulk designed for reuse; return to point of purchase for repackaging. If no repackaging program is available or tank is no longer suitable for reuse, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**CONTAINER DISPOSAL FOR 2.5 GALLONS:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**DISCLAIMER**

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed populations, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. **CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.**

**BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF HERBICIDE 1.** In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

**USES WITH OTHER PRODUCTS (TANK MIXES)**

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company then American Cyanamid Company shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in a combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event, shall be limited to return of the amount of the purchase price of the product.

™Trademark of American Cyanamid Company

**BEST AVAILABLE COPY**

**GENERAL INFORMATION**

HERBICIDE 1 is a uniquely flexible broad-spectrum herbicide which may be applied preplant incorporated, preemergence or postemergence in soybeans.

When HERBICIDE 1 is applied to the soil, susceptible weeds emerge, growth stops, and then the weeds either die or are not competitive with the crop.

When HERBICIDE 1 is applied postemergence, absorption occurs through both the foliage and roots. Susceptible weeds stop growing and either die or are not competitive with the crop.

A timely cultivation may aid in the control of certain weeds or improve general weed control when adequate moisture is not received after application. Cultivation should be shallow. After postemergence treatments, wait at least 10 days before cultivating.

The mode of weed killing activity involves uptake of HERBICIDE 1 by weed roots and/or foliage and rapid translocation to growing points. Therefore, adequate soil moisture is important for optimum HERBICIDE 1 activity. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, then a cultivation or postemergence herbicide application may be needed to improve weed control. When adequate moisture is received after dry conditions, HERBICIDE 1 will provide residual control of susceptible germinating weeds; activity on established weeds will depend on the weed species and the location of its root system in the soil.

A soil treatment of HERBICIDE 1 followed by a postemergence application of HERBICIDE 1, SCEPTER® 70DG, SCEPTER® O.T.® or Classic<sup>5</sup> will control certain problem weeds. Use sequential treatments in recommended states only.

Occasionally, internode shortening of soybean plants may be observed with HERBICIDE 1 applications. This has no effect on soybean yields.

Within any weed population, there may be the potential for the occurrence of weeds resistant to herbicides when compounds that affect the same site of action are used continuously on the same acre. Therefore, any multiple year weed management program should include, when local agronomic practices allow, as many of the following practices as possible:

- use scouting and weed thresholds to determine the need for herbicide treatment
- utilize tank-mixes or sequential applications of different mode-of-action herbicides to control the same target weeds
- combine cultivation with herbicide treatments

See your Cyanamid representative for additional information.

**BEST AVAILABLE COPY**

•Registered Trademarks of American Cyanamid Company  
<sup>5</sup>Registered Trademark of E. I. duPont de Nemours and Company

**PRECAUTIONS:**

**DO NOT** use on crops other than soybeans. Crops other than soybeans, such as cotton, corn and vegetables, may be injured by spray drift or other indirect contact with HERBICIDE 1.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in **SPRAYING INSTRUCTIONS** on pages 25 and 26.

**Do not use HERBICIDE 1 other than in accordance with the instructions set forth on this label.**

To avoid injury to sensitive crops, spray equipment used for HERBICIDE 1 applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops. See pages 31 to 33 for **ROTATIONAL CROP RESTRICTIONS**.

There must be an interval of at least 90 days between the last HERBICIDE 1 application and soybean harvest.

**DO NOT** graze or feed treated soybean forage, hay or straw to livestock.


Use of HERBICIDE 1 herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, crop injury, is always possible. See pages 31 to 33 for **ROTATIONAL CROP RESTRICTIONS**.

**Replanting:** If replanting is necessary in a field previously treated with HERBICIDE 1, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. **DO NOT** apply a second treatment of HERBICIDE 1.

**BEST AVAILABLE COPY**

**USE AREA**

HERBICIDE I can be applied only in the states or parts of states shaded in the following map:



**USE REGION 1** includes eastern Oklahoma, Arkansas, the Missouri bootheel, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and eastern Texas.

**USE REGION 2** includes eastern Kansas (east of U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois (south of S.R. 116 west of Peoria; south of U.S. 24 east of Peoria), Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware, and New Jersey.

**USE REGION 3** includes Nebraska (east of U.S. 81, north of U.S. 34 and also that area east of U.S. 283, south of U.S. 30, and west of U.S. 81), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24 east of Peoria), Wisconsin, Iowa, and southern Minnesota.

**Note:** See **USE RATES** and **ROTATIONAL CROP RESTRICTIONS** sections for recommendations applying to each Use Region.

**BEST AVAILABLE COPY**

**WEEDS CONTROLLED  
HERBICIDE 1 Alone**

When applied as directed, soil-applied treatments of HERBICIDE 1 will control the following weeds:

WEED	LEVEL OF CONTROL
<b><u>BROADLEAF WEEDS</u></b>	
Alligatorweed	Control
Beggarweed, Florida	Control <sup>1</sup>
Bristly Starbur	Control
Burcucumber	Control <sup>2</sup>
Cocklebur, Common	Control
Copperleaf, Hophornbeam	Control <sup>3</sup>
Jimsonweed	Control
Lambsquarters, Common	Control
Mallow, Venice	Control
Mexicanweed	Control <sup>1</sup>
Morningglory	
Entireleaf	Control <sup>4</sup>
Ivyleaf	Control <sup>4</sup>
Palmleaf	Control
Pitted	Control
Smallflower	Control
Tall	Control <sup>4</sup>
Mustard species	Control
Nightshade, Eastern Black	Control
Pigweed	
Palmer	Control
Redroot	Control
Smooth	Control
Spiny	Control
Waterhemp, Tall	Control
Poinsettia, Wild	Control
Puncturevine	Control
Pusley, Florida	Control
Ragweed,	
Common	Control
Giant	Control <sup>5</sup>
Redweed	Control <sup>2</sup>
Sesbania, Hemp	Control <sup>3</sup>
Sicklepod	Control <sup>6</sup>
Sida, Prickly (teaweed)	Control

Smartweed,	
Ladysthumb	Control
Pennsylvania	Control
Spurge, Spotted	Suppression
Sunflower, Common	Control
Texasweed	Control <sup>1</sup>
Velvetleaf	Control <sup>7</sup>

**GRASSES**

Barnyardgrass	Suppression
Corn, Volunteer	Suppression <sup>8</sup>
Foxtail	
Giant	Control
Green	Control
Yellow	Control
Goosegrass	Suppression
Johnsongrass, seedling	Control
Shattercane	Suppression
Signalgrass, Broadleaf	Suppression

**SEDGES**

Nutsedge, Yellow	Suppression <sup>2</sup>
------------------	--------------------------

---

<sup>1</sup> In USE REGION 1 ONLY. A 2/3 pint per acre soil application of HERBICIDE 1 will provide suppression of these weeds. A postemergence application of SCEPTER at 2/3 pint per acre or SCEPTER 70DG at 2.8 ounces per acre must be applied following a soil application of HERBICIDE 1 at 2/3 pint per acre in a planned sequential program to control these weeds. Apply SCEPTER before weeds exceed the 1 to 2 true leaf stage.

<sup>2</sup> To obtain consistent control or suppression of these weeds under a wide range of environmental conditions, a preplant incorporated application is required.

<sup>3</sup> USE REGION 1 and USE REGION 2. SCEPTER O.T. herbicide must be applied postemergence at 1 pint per acre following a soil application of HERBICIDE 1 for control of these weeds. The total amount of SCEPTER O.T. applied should not exceed 1 pint per acre in Use Region 2 or 2 pints per acre in Use Region 1 in one season when following 2/3 pint per acre of HERBICIDE 1. Refer to the SCEPTER O.T. label for application instructions. In Use Region 2, refer to the Rotational Crop Restrictions section of this label for special instructions on planting follow crops when using this program.

<sup>4</sup> A 2/3 pint per acre soil application of HERBICIDE 1 will provide suppression of these weeds. For control of these morningglory species, the sequential program in footnote #3 should be used.

**BEST AVAILABLE COPY**

<sup>5</sup> Some broadleaf weeds such as giant ragweed germinating deep in the soil or with subsequent flushes may require a cultivation or postemergence herbicide application for season long control.

<sup>6</sup> Only light to moderate infestations of sicklepod are controlled. Later sicklepod flushes may require a cultivation or a postemergence herbicide application. For heavy sicklepod infestations, the sequential program in footnote #1 may be used or Classic may be applied postemergence at 1/2 to 3/4 ounces per acre following a soil application of HERBICIDE 1 at 2/3 pint acre. Apply the higher rate of Classic if sicklepod seedlings are taller than 2 inches. A cultivation 14 days after Classic application may be required to control sicklepod escapes. Refer to the Rotational Crop Restrictions section of this label for instructions on planting follow crops when using this program. The Classic sequential program may be used only in Use Region 1 (except Oklahoma).

<sup>7</sup> A preplant incorporated application is recommended for velvetleaf control. A preemergence application of HERBICIDE 1 will aid in the control and reduce competition from velvetleaf.

<sup>8</sup> A soil application of HERBICIDE 1 will suppress only those field corn hybrids which DO NOT possess tolerance or resistance to HERBICIDE 1 or other imidazolinone herbicides (i.e. PURSUIT®).

NOTE: The total amount of HERBICIDE 1 applied should not exceed 1 1/3 pints per acre in one year.

**BEST AVAILABLE COPY**

®Registered Trademark of American Cyanamid Company

**WEEDS CONTROLLED BY POSTEMERGENCE APPLICATIONS**

When applied as direct postemergence treatments of HERBICIDE 1 will control the following weeds:

WEEDS CONTROLLED*	MAXIMUM WEED HEIGHT CONTROLLED (inches)	
	1/3 pint/acre	2/3 pint/acre
Cocklebur, Common	8	12
Pigweed		
Palmer	N	12
Redroot	4	12
Smooth	4	12
Spiny	N	12
Tall waterhemp	N	12
Poinsettia, Wild	N	6
Sunflower, Common	4	8
Volunteer Corn**	8	12

N = Not Controlled

Apply when weeds are actively growing. DO NOT apply HERBICIDE 1 postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes.

\* Use a nonionic surfactant or crop oil concentrate (see page 14).

\*\* A postemergence application of HERBICIDE 1 will control only those field corn hybrids which DO NOT possess tolerance or resistance to HERBICIDE 1 or other imidazolinone herbicides (i.e. PURSUIT).

14736

**DIRECTIONS FOR CONVENTIONAL, MINIMUM, AND NO-TILL APPLICATIONS**

**APPLICATION RATE**

Apply HERBICIDE 1 at a broadcast rate of 2/3 pint per acre for preplant incorporated and preemergence (including minimum and no-till systems) applications. At this broadcast rate, one gallon of HERBICIDE 1 will treat 12 acres of soybeans. Apply HERBICIDE 1 at a broadcast rate of 1/3 or 2/3 pint per acre for postemergence applications.

**NOTE:** For preplant incorporated and preemergence applications on sands and loamy sands, 1/2 pint per acre can be used without a substantial reduction in the control of only cocklebur and pigweed.

**NOTE:** In Use Region 3 as defined by the Use Area section of this label, apply HERBICIDE 1 prior to July 1.

**APPLICATION INSTRUCTIONS**

**SOIL APPLICATIONS**

HERBICIDE 1 may be applied in conventional, minimum, or no-till as a preplant incorporated or preemergence application up to 45 days (30 days in Use Region 1) before planting. Preemergence applications may also be applied during or after planting but before the crop emerges.

Adequate moisture is required for activation of HERBICIDE 1.

**PREPLANT INCORPORATED APPLICATIONS**

Apply HERBICIDE 1 before planting and incorporate uniformly into the top 1 to 2 inches of soil.

HERBICIDE 1 may be applied immediately before planting or up to 30 days prior to planting in Use Region 1. For Use Region 2 and Use Region 3, HERBICIDE 1 may be applied 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or rolling cultivator.

**PREEMERGENCE APPLICATIONS (including no-till)**

Apply HERBICIDE 1 up to 45 days (30 days in Use Region 1) before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate HERBICIDE 1 in the soil. If sufficient rainfall or irrigation to activate HERBICIDE 1 is not received within 7 days after application, a thorough shallow tillage, cultivation, or postemergence treatment may be required for control of emerged weeds.

## **POSTEMERGENCE APPLICATIONS**

**HERBICIDE 1** applied postemergence not only controls the emerged weed species listed previously, but can also provide residual control of susceptible weeds that may emerge after application. Application should be made to small soybeans so more herbicide reaches the soil surface. The level of residual control will depend on weed density and canopy closure which affect the amount of **HERBICIDE 1** reaching the soil surface and environmental conditions, such as rainfall, at the time of and following application.

To maximize weed control following a postemergence **HERBICIDE 1** application, wait at least 10 days before cultivating. This timely cultivation will enhance residual weed control, especially under dry conditions.

Use a nonionic surfactant or crop oil concentrate. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

There must be an interval of at least 90 days between the last **HERBICIDE 1** application and soybean harvest.

## HERBICIDE COMBINATIONS

A soil treatment of HERBICIDE 1 is active against many grass species, but for maximum grass control HERBICIDE 1 should be tank mixed with PROWL 3.3EC herbicide, trifluralin, Lasso or Dual. HERBICIDE 1 may also be applied as a sequential preemergence or postemergence treatment following application of one of the above grass herbicides.

For no-till uses, HERBICIDE 1 may be tank mixed with Gramoxone Extra, Roundup, 2,4-D, or 2,4-DB to kill existing vegetation. Gramoxone Extra, Roundup, 2,4-D or 2,4-DB should be deleted from the tank mixture if vegetation is absent at the time of application. HERBICIDE 1 may also be tank mixed with Bronco, a product which gives both preemergence control of annual grasses and postemergence control of existing vegetation. Use a minimum of 10 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues. Use ground equipment only. Plant soybeans at least one inch deep and adjust planters to ensure adequate seed coverage.

HERBICIDE 1 may be followed by a postemergence application of Fusilade<sup>6</sup> or Poast<sup>7</sup>. DO NOT tank mix HERBICIDE 1 with postemergence grass herbicides.

When used in combination with another herbicide, HERBICIDE 1 should be used only in accordance with recommendations on this label. Always follow the more restrictive label limitations and precautions. HERBICIDE 1 cannot be mixed with any product containing a label prohibition against such mixing.

### Soil Texture

Although the rate of HERBICIDE 1 does not vary with soil texture, the recommended rates of other herbicides used with this product generally do vary with soil texture. Combination rate tables in this label refer to three soil texture groups: COARSE, MEDIUM, and FINE. The following table lists soil textures included in each of these three groups:

<u>COARSE</u>	<u>MEDIUM</u>	<u>FINE</u>
sands	sandy clay loams*	silty clay loams*
loamy sands	sandy clays	silty clays
sandy loams	loams	clay loams
	silt loams	clays
	silts	

\*Sometimes considered transitional soils.

<sup>6</sup>Registered Trademark of Zeneca, Inc.

<sup>7</sup>Registered Trademark of BASF Wyandotte Corporation.

**ADDITIONAL WEEDS CONTROLLED  
BY COMBINATION TREATMENTS**

When applied as directed, HERBICIDE 1 preplant incorporated or preemergence combination treatments with PROWL 3.3EC, trifluralin, Lasso or Dual will control, as indicated, weeds listed in the following table in addition to those controlled by HERBICIDE 1 alone.

<b>WEED</b>	<b>PROWL 3.3EC</b>	<b>Trifluralin<sup>c</sup></b>	<b>Lasso</b>	<b>Dual</b>
<b>GRASSES</b>				
BARNYARDGRASS	X	X	X	X
CRABGRASS, SMOOTH	X	X	X	X
CRABGRASS, LARGE	X	X	X	X
CROWFOOT GRASS	X	X		
GOOSEGRASS	X	X	X	X
PANICUM, FALL	X	X <sup>a</sup>	X	X
PANICUM, TEXAS	X	X		
RICE, RED				X <sup>b</sup>
SANDBUR, FIELD	X	X		
SHATTERCANE	X <sup>c,d</sup>	X <sup>a</sup>		
SIGNALGRASS	X	X	X	X
WITCHGRASS	X	X	X	X
<b>BROADLEAF WEEDS</b>				
CARPETWEED	X	X	X	X
GALINSOGA			X	X
KOCHIA	X	X		
PURSLANE	X	X	X	
SPURGE, ANNUAL	X			

<sup>a</sup>See Note on page 18.

<sup>b</sup>HERBICIDE 1 plus Dual will provide suppression of red rice.

<sup>c</sup>Preplant incorporated treatments only.

<sup>d</sup>See Note on page 17.

**HERBICIDE 1 Plus PROWL 3.3EC Tank Mixture**

Apply HERBICIDE 1 plus PROWL 3.3EC preplant incorporated or preemergence using rates in the table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. For preplant incorporated applications, incorporate into the soil within 7 days after application, and plant soybeans within 45 days after application (30 days in Use Region 1). For preemergence applications, apply from up to 45 days prior to planting (30 days in Use Region 1) to a maximum of 2 days after planting. DO NOT apply if soybeans have emerged.

**Broadcast Rate of HERBICIDE 1 Plus PROWL 3.3EC\***

Soil Texture	Less Than 3% Organic Matter		3% or More Organic Matter	
	HERBICIDE 1 pints/acre	+ PROWL 3.3EC pints/acre	HERBICIDE 1 pints/acre	+ PROWL 3.3EC pints/acre
Coarse	2/3	1.2 to 1.8	2/3	1.8
Medium	2/3	1.8 to 2.4	2/3	1.8 to 2.4
Fine	2/3	1.8 to 2.4	2/3	2.4 to 3.0

NOTE: Use the higher rate of PROWL 3.3EC if heavy weed populations are anticipated or if shattercane will be present.

Observe all precautions and limitations on PROWL 3.3EC label.

\* Based on 3.3 pounds of active ingredient per gallon. If the 4.0 pounds of active ingredient per gallon formulation is used, multiply the above PROWL 3.3EC rates by 0.82.

### HERBICIDE 1 Plus Trifluralin Tank Mixture

Apply HERBICIDE 1 plus trifluralin preplant incorporated using rates in the table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. Incorporate into the soil within 24 hours after application, and plant soybeans within 45 days after application (30 days in Use Region 1). Use equipment that provides uniform 2 inch incorporation.

#### Broadcast Rate of HERBICIDE 1 Plus trifluralin\*

Soil Texture	HERBICIDE 1 pints/acre	Trifluralin* pints/acre
Coarse	2/3	1
Medium	2/3	1.5
Fine	2/3	2

NOTE: If fall panicum or shattercane will be present, refer to trifluralin label for higher rates to control these weeds.

Observe all precautions and limitations on the trifluralin label.

\*Based on 4 pounds active ingredient per gallon.

**HERBICIDE 1 Plus Lasso Tank Mixture**

Apply HERBICIDE 1 plus Lasso preplant incorporated or preemergence using rates in the table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. For preplant incorporated applications, apply and incorporate within 7 days prior to planting. For preemergence applications, apply up to 7 days prior, during or immediately after planting.

**Broadcast Rate of HERBICIDE 1 Plus Lasso**

Soil Texture	<u>Less than 3% Organic Matter</u>			<u>3% or More Organic Matter</u>		
	<u>HERBICIDE 1</u>	<u>Lasso (quarts/acre)</u>		<u>HERBICIDE 1</u>	<u>Lasso (quart/acre)</u>	
	(pints/acre)	Preplant Incorporated	Preemergence	(pints/acre)	Preplant Incorporated	Preemergence
Coarse	2/3	2.0 to 2.5	2.0	2/3	2.0 to 2.5	2.0
Medium	2/3	2.5 to 3.0	2.0 to 2.5	2/3	2.5 to 3.0	2.0 to 2.5
Fine	2/3	2.5 to 3.0	2.0 to 3.0*	2/3	2.5 to 3.0	2.5 to 3.0

\*On silty clay and heavy clay soils of the Mississippi Delta, use 2 to 2-1/2 quarts of Lasso.

**NOTE:** For preplant incorporated applications, use the higher rate of Lasso if heavy weed populations are anticipated.

Use the higher rate of Lasso (1) when heavy grass infestations are anticipated, (2) when fall panicum or crabgrass will be present or (3) when previous crop residues are extremely dense. Apply up to 7 days before, during or immediately after planting.

Observe all precautions and limitations on the Lasso label.

**HERBICIDE 1 Plus Dual Tank Mixture**

Apply HERBICIDE 1 plus Dual preplant incorporated or preemergence using rates in table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. For preplant incorporated applications, apply up to 29 days prior to planting. Incorporate within 7 days after application. For preemergence applications, apply up to 29 days prior, during or immediately after planting.

**Broadcast Rate of HERBICIDE 1 Plus Dual**

Soil Texture	<u>Less Than 3% Organic Matter</u>		<u>3% or More Organic Matter</u>	
	HERBICIDE 1 pints/acre	DUAL pints/acre	HERBICIDE 1 pints/acre	DUAL pints/acre
Coarse	2/3	1.25	2/3	1.5
Medium	2/3	1.5	2/3	2.0
Fine	2/3	2.0	2/3	2.0 to 2.5*

\*Use the higher rate of Dual if heavy weed populations are anticipated.

NOTE: Observe all precautions and limitations on Dual label.

**HERBICIDE 1 Plus PROWL 3.3EC or Trifluralin Plus Command 4EC Tank Mixture**

Observe all precautionary statements and limitations in the PROWL 3.3EC or trifluralin and Command 4EC labels before using. The HERBICIDE 1 plus PROWL 3.3EC or trifluralin plus Command 4EC preplant incorporated tank mixture may be used when heavy infestations of velvetleaf are anticipated. Uniformly apply the tank mixture with properly calibrated ground equipment in 10 to 40 gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application.

Apply HERBICIDE 1 plus PROWL 3.3EC or trifluralin plus Command 4EC to the soil surface and uniformly incorporate. Refer to the Command 4EC label for incorporation instructions. DO NOT apply aerially or through irrigation equipment. DO NOT apply this tank mixture to overly moist or wet soils.

**SPECIAL PRECAUTION**

Command 4EC is a volatile compound. Off-site movement of spray drift or vapors of Command 4EC can cause foliar whitening or yellowing of some plants. Prior to using Command 4EC, read and strictly follow all precautions and application instructions of the Command 4EC label. The use of an agriculturally approved drift reducing agent is required at finished spray volumes of 10 to 15 gallons per acre.

**BROADCAST RATE PER ACRE OF HERBICIDE 1 PLUS PROWL 3.3EC OR TRIFLURALIN PLUS COMMAND 4EC**

Apply HERBICIDE 1 at 2/3 pint per acre. Apply Command 4EC at 1/4 to 1/2 pint per acre. Refer to HERBICIDE COMBINATIONS section for PROWL 3.3EC (page 17) or trifluralin (page 18) use rates.

**ROTATIONAL CROP RESTRICTIONS:** Follow all rotational crop restrictions on the HERBICIDE 1, PROWL 3.3EC or trifluralin, and Command 4EC label.

**ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.**

### HERBICIDE 1 Plus Metribuzin Tank Mixtures

Before using, observe all precautionary statements in the HERBICIDE 1, Sencor<sup>8</sup>, Lexone<sup>9</sup>, PROWL 3.3EC, Lasso, Dual, and trifluralin leaflet labels. HERBICIDE 1 may be applied in tank mix combination with Sencor or Lexone plus one of the following grass herbicides: PROWL 3.3EC, Lasso, Dual, or trifluralin; either preplant incorporated or preemergence (excluding trifluralin). HERBICIDE 1 may also be applied in tank mix with any one of the above grass herbicides as a preplant incorporated application followed by a preemergence application of Sencor or Lexone. For preemergence applications, apply during or immediately after planting. These treatments will control Florida beggarweed, hemp sesbania, hophornbeam copperleaf and sicklepod in addition to the weeds controlled by a soil application of HERBICIDE 1 plus the grass herbicide.

Apply Sencor or Lexone at the broadcast rate per acre specified in the table below, in tank mix combination with 2/3 pint per acre of HERBICIDE 1 plus one of the above grass herbicides.

#### Sencor or Lexone Rate \*

	4.0L	75%DF
Coarse Soils	1/2 pint	1/3 lb.
Medium Soils	1/2 pint	1/3 lb.
Fine Soils	2/3 pint	1/2 lb.

\* Do not use Sencor or Lexone on coarse soils of less than 2% organic matter. Use 1/2 lb. or 2/3 pint of Sencor or Lexone on medium soils of greater than 4% organic matter.

<sup>8</sup>Registered Trademark of Miles Company

<sup>9</sup>Registered Trademark of E.I. duPont de Nemours and Company

### Tank Mixtures with 2,4-D

2,4-D may be used with HERBICIDE 1 alone or in combination with other HERBICIDE 1 tank mixes prior to planting for control of some annual broadleaf weeds. Use the following rates of 2,4-D in tank mixtures with HERBICIDE 1:

<u>2,4-D Formulation</u>	<u>Rate (lb a.i./A)</u>	<u>Minimum Days Before Planting</u>
Ester	0.5	7
Amine	0.5	15
Ester or Amine	1.0	30

Refer to 2,4-D label for weeds controlled. Observe all cautions and limitations on the 2,4-D label.

### Tank Mixtures With 2,4-DB

2,4-DB may be used with HERBICIDE 1 alone or in combination with other HERBICIDE 1 tank mixes for the control of some annual broadleaf weeds. Refer to the 2,4-DB label for specific use recommendations, rates and weeds controlled.

Observe all cautions and limitations on the 2,4-DB label.

### Tank Mixtures With Bronco

Use the following rates of Bronco for HERBICIDE 1 plus Bronco tank mixtures:

<u>Soil Texture</u>	<u>Bronco (quarts/acre)</u>
Coarse	3 to 5
Medium	4 to 5
Fine	4 to 5

Use the higher rate of Bronco (1) when previous crop residues are extremely dense, (2) when emerged weeds are over 6 inches tall, (3) when heavy infestations of weeds are expected, and (4) when soils contain more than 3% organic matter. Apply up to 7 days before, during or immediately after planting.

Observe all precautions and limitations on Bronco label.

### **Tank Mixtures With Roundup**

Roundup may be used with HERBICIDE 1 alone or in combination with other tank mixes for the control of certain emerged grasses and broadleaf weeds. Refer to the Roundup label for specific use recommendations, rates and weeds controlled.

**NOTE:** The Roundup rate for tank mixes with HERBICIDE 1 is the same as the rate for Roundup used alone.

Observe all cautions and limitations on the Roundup label.

### **Tank Mixtures With Gramoxone Extra**

Apply 1-1/2 to 2-1/2 pints of Gramoxone Extra per acre in HERBICIDE 1 tank mixtures. Use the 2-1/2 pint rate if weeds are 4 to 6 inches tall. Weeds over 6 inches may not be controlled with this treatment. Apply up to 14 days before, during or immediately after planting.

When Gramoxone Extra is included in a tank mixture, add a nonionic spreader surfactant at a rate of 8 fl. oz. per 100 gallons of spray mixture as the last ingredient in the tank.

Gramoxone Extra will control most annual emerged weeds and suppress many emerged perennials. Refer to the Gramoxone Extra label for specific use recommendations and weeds controlled.

Observe all precautions and limitations on Gramoxone Extra label.

### MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. While agitating add the required amount of HERBICIDE 1 and then fill the remainder of the tank with water. For postemergence applications, add the nonionic surfactant or the crop oil concentrate as last ingredient in the tank. An antifoaming agent may be added to the tank if needed. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank mixing HERBICIDE 1 with recommended herbicides, add HERBICIDE 1 to the spray tank first and make sure it is thoroughly mixed before adding the other herbicide.

When tank mixing Roundup or Gramoxone Extra with HERBICIDE 1 and PROWL 3.3EC, Lasso or Dual, add herbicides to the tank in the following order: (1) HERBICIDE 1 (2) PROWL 3.3EC, Lasso or Dual, (3) Roundup or Gramoxone Extra, and (4) nonionic surfactant.

### SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

#### GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. To minimize drift, use a maximum spray pressure of 40 psi.

To ensure thorough coverage, use a minimum of 20 gallons of water per acre when applying HERBICIDE 1 as a postemergence treatment. Use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply with ground equipment when wind velocity is greater than 10 mph.

For band treatment, apply the broadcast equivalent rate and volume per acre.

To determine these:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Rate per acre} = \text{Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per acre} = \text{Band VOLUME per acre}$$

**AERIAL APPLICATIONS:**

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following **SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS:**

- Use nozzles which produce a coarse spray.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph.

Applicator is responsible for any loss or damage which results from spraying HERBICIDE 1 in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

## APPLICATIONS WITH FERTILIZERS

### APPLICATIONS WITH LIQUID FERTILIZERS

HERBICIDE 1 can be applied to the soil in liquid fertilizers, alone or in combination with PROWL 3.3EC, trifluralin, Lasso, Dual, Command 4EC or metribuzin. Follow all HERBICIDE 1 label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the HERBICIDE 1/liquid fertilizer mixture.

### LIQUID FERTILIZER COMPATIBILITY DETERMINATIONS

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of HERBICIDE 1 alone or with other herbicides in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant to one jar. See recommended adjuvants below.

**NOTE:** Compreh-Kalo Laboratories Incorporated, Kansas City, MO; Sponyo 168 D-Witco-Retzloff Chemical Company, Houston, TX; Unite-Hopkins Agricultural Chemical Company, Madison, WI; or other compatible adjuvants.

3. (a) When using HERBICIDE 1 alone, add to each jar the correct amount of HERBICIDE 1 as specified in the table below.
- (b) When using HERBICIDE 1 tank mixtures, first add the specified quantity of HERBICIDE 1 and then add the correct amount of PROWL 3.3EC, trifluralin, Lasso, Dual, Command 4EC or metribuzin.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer of globules, sludge, flakes or other precipitates.
  - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
  - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.

- (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
- (d) If separation of the mixture occurs and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

**Teaspoons of Specific Herbicide to be added to  
1 Pint of Liquid Fertilizer Solution**

Gallons of Liquid Fertilizer to be Applied per Acre	HERBICIDE 1	PROWL 3.3EC	Trifluralin	Lasso	Dual
20	1/2	3	2-1/2	5	2-1/2
30	1/4	2	1-3/4	3-1/2	1-3/4
40	1/4	1-1/2	1-1/4	2-1/2	1-1/4

## APPLICATIONS WITH DRY BULK FERTILIZERS

HERBICIDE 1 may be impregnated on dry bulk fertilizers. When applied as directed, HERBICIDE 1/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of HERBICIDE 1 applied in water.

Apply HERBICIDE 1/dry bulk fertilizer mixtures only with ground equipment.

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the HERBICIDE 1/dry bulk fertilizer mixture.

A minimum of 200 pounds and a maximum of 450 pounds of dry bulk fertilizer impregnated with the recommended amount of HERBICIDE 1 must be applied per acre.

DO NOT impregnate HERBICIDE 1 onto coated ammonium nitrate or limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with HERBICIDE 1. A minimum of 200 pounds of impregnated dry bulk fertilizer, excluding the weight of ammonium nitrate or limestone, must be applied per acre.

Use the following table to determine the amount of HERBICIDE 1 to be impregnated on a ton of dry bulk fertilizer based on the rate of fertilizer which will be applied per acre.

### RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZER WITH HERBICIDE 1

(Pints of HERBICIDE 1 per Ton of Fertilizer)

HERBICIDE 1 Rate Per Acre	Fertilizer Rate lbs/acre	Pints per Ton
2/3 Pint	200	6-2/3
	250	5-1/3
	300	4-1/2
	350	3-3/4
	400	3-1/3
	450	3

For those rates not listed in this table, calculate the pints of HERBICIDE 1 to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times \frac{2/3 \text{ pint of HERBICIDE 1 per acre (recommended rate)}}{1} = \text{Pints of HERBICIDE 1 per ton of fertilizer}$$

To impregnate HERBICIDE 1 on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of HERBICIDE 1 onto the fertilizer during mixing.

Apply the HERBICIDE 1/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The HERBICIDE 1/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

### ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying HERBICIDE 1 at recommended rates in soybeans:

<u>CROP</u>	<u>USE REGION 1</u>	<u>USE REGION 2</u> (except Michigan <sup>3</sup> )	<u>USE REGION 3</u>
Soybeans	No restrictions	No restrictions	No restrictions
Wheat	4 months	4 months <sup>4</sup>	See FOOTNOTE 5 & 6
Rice	Spring following HERBICIDE 1 application	Spring following HERBICIDE 1 application <sup>4</sup>	---
Barley	11 months	11 months <sup>4</sup>	See FOOTNOTE 5
Field Corn	11 months <sup>1</sup>	11 months <sup>2,4</sup>	See FOOTNOTE 5 & 6
Edible Beans	11 months	11 months <sup>4</sup>	11 months
Grain Sorghum	11 months	11 months <sup>4</sup>	11 months
Oats	11 months	11 months <sup>4</sup>	See FOOTNOTE 5
Peanuts	11 months	11 months <sup>4</sup>	11 months
Tobacco	11 months	11 months <sup>4</sup>	11 months
Sugar Beets	26 months	26 months	26 months
Other Crops	18 months	18 months	See FOOTNOTE 7

<sup>1</sup> For USE REGION 1 as defined in the USE AREA section of this label; field corn may be planted in the spring of the year following HERBICIDE 1 application unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received within 6 months following the date of last application). A minimum of 10 inches of rainfall is needed for a postemergence application of HERBICIDE 1 only if the total amount of HERBICIDE 1 applied in one season does not exceed 1/3 pint per acre.

<sup>2</sup> For USE REGION 2 (except Michigan<sup>3</sup>) as defined in the USE AREA section of this label; field corn may be planted in the spring of the year following HERBICIDE 1 application unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of last application through November 15 of the same year). A minimum of 10 inches of rainfall is needed for a postemergence application of HERBICIDE 1 only if the total amount of HERBICIDE 1 applied in one season does not exceed 1/3 pint per acre.

If the minimum rainfall requirement is not met, only field corn hybrids which possess tolerance or resistance to HERBICIDE 1 and other imidazolinone herbicides may be planted the spring of the year following an application including HERBICIDE 1 (SEE NOTE).

3 Growers in the counties of Berrien, Cass, St. Joseph, Branch, Hillsdale, Lenawee, Monroe, Van Buren, Kalamazoo, Calhoun, Jackson, Washtenaw, and Wayne may use the rotational crop restrictions for Use Region 2.

Growers in other Michigan counties may NOT plant oats or barley in the fall or spring of the year following a HERBICIDE 1 application. In this geography, only field corn hybrids which possess tolerance or resistance to HERBICIDE 1 and other imidazolinone herbicides may be planted the spring of the year following an application including HERBICIDE 1 (SEE NOTE).

4 **ROTATIONAL CROP RESTRICTIONS for a sequential application of SCEPTER O.T. following a soil application of HERBICIDE 1 (USE REGION 2 only):**

For USE REGION 2 as defined in the USE AREA section of this label; soybeans may be planted anytime. Barley, edible beans, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application.

Only field corn hybrids which possess tolerance or resistance to HERBICIDE 1 and other imidazolinone herbicides may be planted the spring of the year following a sequential application (SEE NOTE). Other field corn varieties may be planted 15 months after the last herbicide application.

5 For USE REGION 3 as defined in the USE AREA section of this label; field corn, wheat, barley, or oats may be planted as a rotational crop in the fall or in the spring of the year following HERBICIDE 1 application ONLY if HERBICIDE 1 was applied as a postemergence application at 1/3 pint per acre and at least 10 inches of rainfall or irrigation has been received from the date of application through October of the same year.

6 In Nebraska, east of U.S. 283, south of U.S. 30, and west of U.S. 81; wheat may be planted 4 months after a HERBICIDE 1 application. In this geography, only field corn hybrids which possess tolerance or resistance to HERBICIDE 1 and other imidazolinone herbicides may be planted the spring of the year following an application including HERBICIDE 1 (SEE NOTE).

7 For USE REGION 3 as defined in the USE AREA section of this label; canola, table beets, strawberries, cabbage, tomatoes, potatoes, carrots, celery, cole crops, garlic, onions, spinach, asparagus, cauliflower, and broccoli may be planted 26 months after HERBICIDE 1 application. Other crops may be planted 18 months after HERBICIDE 1 application.

NOTE: Contact your chemical dealer, seed supplier, or American Cyanamid to obtain information regarding the availability of imidazolinone tolerant field corn hybrids which are adapted to your area.

**ROTATIONAL CROP RESTRICTIONS for sequential  
Classic application following HERBICIDE 1 soil treatments (USE REGION 1 only):**

Soybeans may be planted anytime. Barley, edible beans, field corn, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application. Refer to rotational crop restrictions listed in the Classic label. Always follow the more restrictive label.

Only rotational crops harvested at maturity may be used for feed or food.

Use of HERBICIDE 1 herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, crop injury, is always possible.

Except where stated on a Supplemental Label or in this label, DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy<sup>10</sup>, Gemini<sup>10</sup>, Lorox Plus<sup>10</sup>, Preview<sup>10</sup>, etc.) or imazaquin (HERBICIDE 1, SQUADRON<sup>®</sup>, TRI-SCEPT<sup>®</sup>, SCEPTER O.T.) the same year as HERBICIDE 1. DO NOT apply products containing imazethapyr (e.g. PURSUIT, PURSUIT<sup>®</sup> PLUS or PASSPORT<sup>®</sup>) the same year as HERBICIDE 1.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

<sup>10</sup> Registered Trademarks of E.I. du Pont de Nemours and Company.

<sup>®</sup>Registered Trademarks of American Cyanamid Company.

## WEED SCIENTIFIC NAMES

BROADLEAF WEEDS

Alligatorweed	( <i>Alternanthera philoxeroides</i> )
Beggarweed, Florida	( <i>Desmodium tortuosum</i> )
Bristly Starbur	( <i>Acanthospermum hispidum</i> )
Burcucumber	( <i>Sicyos angulatus</i> )
Carpetweed	( <i>Mollugo verticillata</i> )
Cocklebur, Common	( <i>Xanthium strumarium</i> )
Copperleaf, Hophornbeam	( <i>Acalypha ostryacifolia</i> )
Galinsoga	( <i>Galinsoga</i> spp.)
Jimsonweed	( <i>Datura stramonium</i> )
Kochia	( <i>Kochia scoparia</i> )
Lambsquarters, Common	( <i>Chenopodium album</i> )
Mallow, Venice	( <i>Hibiscus trionum</i> )
Mexicanweed	( <i>Caperonia castanifolia</i> )
Morningglory	
Entireleaf	( <i>Ipomoea hederacea</i> var. <i>intergriuscula</i> )
Ivyleaf	( <i>Ipomoea hederacea</i> )
Palmleaf	( <i>Ipomoea wrightii</i> )
Pitted	( <i>Ipomoea lacunosa</i> )
Smallflower	( <i>Jacquemontia tamnifolia</i> )
Tall	( <i>Ipomoea purpurea</i> )
Mustard species	( <i>Brassica</i> spp.)
Nightshade, Eastern Black	( <i>Solanum ptycanthum</i> )
Pigweed	
Palmer	( <i>Amaranthus palmeri</i> )
Redroot	( <i>Amaranthus retroflexus</i> )
Smooth	( <i>Amaranthus hybridus</i> )
Spiny	( <i>Amaranthus spinosus</i> )
Waterhemp, Tall	( <i>Amaranthus tuberculatus</i> )
Poinsettia, Wild	( <i>Euphorbia heterophylla</i> )
Puncturevine	( <i>Tribulus terrestris</i> )
Purslane	( <i>Portulaca oleracea</i> )
Pusley, Florida	( <i>Richardia scabra</i> )
Ragweed,	
Common	( <i>Ambrosia artemisiifolia</i> )
Giant	( <i>Ambrosia trifida</i> )
Redwee.†	( <i>Melochia corchorifolia</i> )
Sesbania, Hemp	( <i>Sesbania exaltata</i> )
Sicklepod	( <i>Cassia obtusifolia</i> )
Sida, Prickly (teaweed)	( <i>Sida spinosa</i> )
Smartweed,	
Ladysthumb	( <i>Polygonum persicaria</i> )
Pennsylvania	( <i>Polygonum pennsylvanicum</i> )
Spurge	
Annual	( <i>Euphorbia</i> spp.)
Spotted	( <i>Euphorbia maculata</i> )
Sunflower, Common	( <i>Helianthus annuus</i> )
Texasweed	( <i>Caperonia palustris</i> )
Velvetleaf	( <i>Abutilon theophrasti</i> )

GRASSES

Barnyardgrass  
 Corn, Volunteer  
 Crabgrass  
   Smooth  
   Large  
 Crowfoot Grass  
 Foxtail  
   Giant  
   Green  
   Yellow  
 Goosegrass  
 Johnsongrass, seedling  
 Panicum  
   Fall  
   Texas  
 Rice, Red  
 Sandbur, Field  
 Shattercane  
 Signalgrass, Broadleaf  
 Witchgrass

(*Echinochloa crus-galli*)  
 (*Zea mays*)  
  
 (*Digitaria ischaemum*)  
 (*Digitaria sanguinalis*)  
 (*Dactyloctenium aegyptium*)  
  
 (*Setaria faberi*)  
 (*Setaria viridis*)  
 (*Setaria glauca*)  
 (*Eleusine indica*)  
 (*Sorghum halepense*)  
  
 (*Panicum dichotomiflorum*)  
 (*Panicum texanum*)  
 (*Oryza sativa*)  
 (*Cenchrus incertus*)  
 (*Sorghum bicolor*)  
 (*Brachiaria platyphylla*)  
 (*Panicum capillare*)

SEDGES

Nutsedge, Yellow

(*Cyperus esculentus*)