

AUG 23 1995

Under the Federal Insecticide,
Fungicide, and Plant Growth Regulator
Act of 1947, as amended, this product is
registered for use as a herbicide in
soybeans.
EPA Reg. No. 241-289

PM 25 241-289
EPA Reg. No. 241-289

10/4

FOR USE IN SOYBEANS

SCEPTER AND ROUNDUP¹ FOR WEED CONTROL IN SOYBEANS AND SCEPTER USE IN ROUNDUP READY² SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER LEAFLET LABEL BEFORE USING. SEE THE SCEPTER LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application. The use of SCEPTER herbicide not consistent with this labeling may result in injury to crops. Keep container closed to avoid spills and contamination. **DO NOT** apply this product through any type of irrigation system.

GENERAL INFORMATION

SCEPTER provides control of broadleaf and grass weeds when applied preplant, preemergence and postemergence. SCEPTER can be applied to either conventional or no-till soybeans. When applied as a burndown application, Roundup may be tank-mixed with SCEPTER for control of weeds common to no-till production, such as marestail.

SCEPTER may also be applied to Roundup Ready soybeans.

Refer to the SCEPTER label and the Roundup label for specific use recommendations, rates, and weeds controlled.

No-till Soybeans (non-glyphosate resistant)

Apply SCEPTER plus Roundup for weed control in no-till soybeans. Roundup will enhance the control of weeds including marestail and prickly lettuce. Always include a nonionic surfactant in the spray solution. **DO NOT** apply Roundup postemergence to non-glyphosate resistant soybeans.

Roundup Ready Soybeans (glyphosate resistant)

SCEPTER may be used for weed control in Roundup Ready soybeans. Apply SCEPTER either preplant soil applied, preplant as a burndown treatment, or for postemergence weed control. SCEPTER may be applied either prior to Roundup, or following Roundup for general weed control and residual activity on broadleaf and grass weeds. For sequential treatments, a sufficient time period should occur between treatments to allow an appropriate assessment of weed control needs.

SCEPTER may be tank-mixed with other herbicides (not Roundup) on Roundup Ready soybeans when applied postemergence. Refer to the SCEPTER label for recommended herbicides. Refer to the other herbicide labels for weeds controlled and recommended application rates.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. When tank-mixing SCEPTER with recommended herbicides, while agitating, add the required amount of SCEPTER to the spray tank first and make sure it is thoroughly mixed before adding the other herbicide; then fill the remainder of the tank with clean water. Maintain agitation while

10/4

SCEPTER HERBICIDE USE AREA

See SCEPTER leaflet label for use area and a description of Use Regions.

APPLICATION INSTRUCTIONS

SOIL APPLICATIONS

SCEPTER may be applied in conventional, minimum, or no-till soybeans as a preplant incorporated or preemergence application up to 45 days (30 days in Use Region 1) before planting. Preemergence applications may also be applied during or after planting but before the crop emerges.

Adequate moisture is required for activation of SCEPTER.

PREPLANT INCORPORATED APPLICATIONS

Apply SCEPTER before planting and incorporate uniformly into the top 1 to 2 inches of soil.

SCEPTER may be applied immediately before planting or up to 30 days prior to planting in Use Region 1. For Use Region 2 and Use Region 3, SCEPTER may be applied 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or rolling cultivator.

PREEMERGENCE APPLICATIONS (including no-till)

Apply SCEPTER up to 45 days (30 days in Use Region 1) before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate SCEPTER in the soil. If sufficient rainfall or irrigation to activate SCEPTER is not received within 7 days after application, a thorough shallow tillage, cultivation, or postemergence treatment may be required for control of emerged weeds.

Refer to the SCEPTER label and Roundup label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS, PRECAUTIONARY STATEMENTS AND WORKER PROTECTION INFORMATION. Refer to the Roundup label for additional information regarding application rates, restrictions, weeds controlled and other information.

When tank-mixing herbicides, always follow the most restrictive label.

Use of SCEPTER herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

®Registered Trademark of American Cyanamid Company

¹ Roundup is a trademark of Monsanto Agricultural Products Company.

² Roundup Ready is a trademark of Monsanto Agricultural Products Company.

PE-XXXX 7/7/95 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1995

PM 25 241-289
EPA Reg. No. 241-289

3074

FOR USE IN SOYBEANS

SCEPTER PLUS ROUNDUP¹ TANK MIX FOR WEED CONTROL IN ROUNDUP READY² SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER LEAFLET LABEL BEFORE USING. SEE THE SCEPTER LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application. The use of SCEPTER herbicide not consistent with this labeling may result in injury to crops. Keep container closed to avoid spills and contamination. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

SCEPTER provides control of broadleaf and grass weeds when applied postemergence. SCEPTER can be applied to either conventional or no-till soybeans. Roundup may be tank-mixed with SCEPTER for enhanced control of weeds common to soybean production. When applied postemergence, tank mixes containing Roundup herbicide can only be applied to Roundup Ready soybeans. Refer to the SCEPTER label and the Roundup label for specific use recommendations, rates and weeds controlled.

MIXING INSTRUCTIONS


Fill the spray tank one-fourth to one-half full with clean water. When tank-mixing SCEPTER with Roundup, while agitating, add the required amount of SCEPTER to the spray tank first and make sure it is thoroughly mixed before adding Roundup; then fill the remainder of the tank with clean water. Maintain agitation while spraying to ensure a uniform spray mixture. For postemergence applications, add the nonionic surfactant as the last ingredient in the tank. To avoid injury to sensitive crops, spray equipment used for SCEPTER herbicide applications must be drained and thoroughly cleaned with water before being used to apply other products.

USE RATE

Refer to the SCEPTER and Roundup labels for recommended use rates.

SCEPTER HERBICIDE USE AREA

See SCEPTER leaflet label for use area and a description of Use Regions.



(See other side)

When tank-mixing herbicides, always follow the most restrictive label.

Use of SCEPTER herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

®Registered Trademark of American Cyanamid Company

¹ Roundup is a trademark of Monsanto Agricultural Products Company.

² Roundup Ready is a trademark of Monsanto Agricultural Products Company.

PE-XXXX 7/7/95 SD



American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1995