Page 192

SCEPTER® herbicide

Supplemental Labeling

EPA Reg. No. 241-289

FOR USE IN SOYBEANS THIS LABEL EXPIRES DECEMBER 31, 1998

REVISED TOBACCO ROTATIONAL INTERVAL

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER LABEL BEFORE USING. SEE THE SCEPTER LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

ROTATIONAL CROP RESTRICTIONS

Tobacco may be planted 9½ months following an application of SCEPTER at up to 2/3 pint per acre and no more than a total of 0.125 pounds of imazaquin applied per acre.

Use of SCEPTER herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER and, therefore, rotational crop injury is always possible.

ACCEPTED
with COMMENTS
In EPA Letter Dated
DEC - 4 1996

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the posticide registered under EPA Reg. No.



Registered Trademark of American Cyanamid Company PE- 10/28/96 SD

SCEPTER®

Supplemental Labeling

herbicide

EPA Reg. No. 241-289

FOR USE IN SOYBEANS THIS LABEL EXPIRES APRIL 30, 1998

%

ECO LM

SCEPTER followed by Classic for Sicklepod Control in Soybeans

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER LABEL BEFORE USING. SEE THE SCEPTER LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

An early preplant, preplant incorporated, or preemergence application of SCEPTER may be made at the 2/3 pint per acre rate. For heavy sicklepod infestations Classic may be applied postemergence at ½ to ¾ ounces per acre following a soil application of SCEPTER. Apply the higher rate of Classic if sicklepod seedlings are taller than 2 inches. A cultivation 14 days after Classic application may be required to control sicklepod escapes. SCEPTER followed by Classic sequential program may be used only in Use Region 1 (except Oklahoma), Virginia and Kentucky.

ROTATIONAL CROP RESTRICTIONS

Soybeans may be planted anytime. Field Corn may be planted 9½ months after the last herbicide application. Barley, edible beans, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application. Refer to rotational crop restrictions listed in the Classic label. Always follow the more restrictive label.

Consult the SCEPTER label for spraying and application information, weeds controlled, and use rates. When tank mixing, always follow the more restrictive label.

Use of SCEPTER herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER and, therefore, rotational crop injury is always possible.



ACCEPTED

SHIP COMMENTS
IN EPA Lener Dated

DEC - 4 1996

Under the Federal Insecticide, Fundicide, and Redenticide Act as amended, for the pesticide registered under EPA Reg. No.

 Registered Trademark of American Cyanamid Company
 Registered trademark of E. I. du Pont de Nemours and Company PE-12309 Rev. 1 10/17/96 SD