


- PM 25 241-294

10/14



BEST AVAILABLE COPY

CONTAIN

herbicide

FIRST AID

IF ON SKIN: Wash with plenty of soap and water.
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably, mouth-to-mouth. Get medical attention.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Harmful if absorbed through skin or inhaled. Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of CONTAIN should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers. DO NOT mix, store or apply CONTAIN or spray solutions of CONTAIN in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. CONTAIN herbicide should be used only in accordance with recommendations on the label. Keep containers closed to avoid spills and contamination. See inside label for complete Directions for Use.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT store below 32°F. DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. See Disclaimer inside Label.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION!

Harmful if absorbed through skin or inhaled. Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

IF ON SKIN: Wash with plenty of soap and water.
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

In case of emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3103.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of CONTAIN should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT mix, store or apply CONTAIN or spray solutions of CONTAIN in unlined steel (except stainless steel) containers or spray tanks.

IMPORTANT

DO NOT use on food or feed crops. DO NOT treat irrigation ditches or water used for crop irrigation or for domestic purposes. Keep from contact with fertilizers, insecticides, fungicides and seeds. DO NOT apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots extend, or in locations where the chemical may be washed or mixed into contact with their roots. DO NOT use on lawns, walks, driveways, or similar areas. DO NOT side trim desirable vegetation with this product.

uct. Prevent drift of spray to desirable plants. DO NOT USE in California.

Clean application equipment after using this product by thoroughly flushing with water.

GENERAL INFORMATION

CONTAIN herbicide is an aqueous solution to be mixed in water and applied as a spray for the non-selective control of most annual and perennial grasses and broadleaf weeds on noncropland areas.

CONTAIN may be applied either preemergence or postemergence to the weeds; however, postemergence application is the method of choice in most situations, particularly for control of perennials. For maximum activity, weeds should be growing vigorously at the time of postemergence applications. The preemergence activity of CONTAIN will provide residual control of many weed species following a postemergence application.

CONTAIN is readily absorbed through foliage and roots and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. Treated plants stop growing soon after spray application. Chlorosis appears first in the newest leaves, and necrosis spreads from this point. In perennials, the herbicide is translocated into and kills underground storage organs, thus preventing regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two weeks after application. Complete kill of plants may not occur for several weeks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CONTAIN herbicide should be used only in accordance with recommendations on this label. Keep containers closed to avoid spills and contamination.

A postemergence use of CONTAIN is recommended for control of most annual and perennial grasses, broadleaf weeds, brush, and trees on NONCROPLAND areas such as fence rows, farm buildings, utility installations, non-irrigation ditch banks, unimproved roads, and other areas around the farmyard where vegetation is undesirable. DO NOT treat where the roots of desirable vegetation may extend into the treated zone as death or severe injury may result. As a general guide, roots often extend to within twice the drip line (i.e. twice the branch extremities); however, the roots of

BEST AVAILABLE COPY

30814

BEST AVAILABLE COPY

some plants may extend beyond this zone (See diagram below).



STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT store below 32°F. DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPRAYING INSTRUCTIONS

Uniformly apply CONTAIN with a properly calibrated sprayer in sufficient volume to insure adequate coverage. DO NOT ALLOW DRIFT ONTO OR INTO THE VICINITY OF DESIRABLE NONTARGET PLANTS.

HAND-HELD SPRAY EQUIPMENT

Thoroughly mix a 1 to 2 percent solution of CONTAIN in water. To determine the proper percent solution of CONTAIN to use, see the "WEEDS CONTROLLED" section of this label and the "PERCENT SOLUTION RATE GUIDE" below.

PERCENT SOLUTION RATE GUIDE

CONTAIN RATE PER ACRE	PERCENT SOLUTION TO MIX
2 - 6 pints	1 %
6 - 8 pints	1.5 %
8 - 12 pints	2 %

For best results, uniformly cover the foliage of the vegetation to be controlled with the spray solution. DO NOT over apply and cause run-off on the treated foliage. To mix the spray solution, add the volume of CONTAIN indicated in this table below to the desired amount of water.

SPRAY SOLUTION MIXING GUIDE

SOLUTION VOLUME	AMOUNT OF CONTAIN TO USE (fluid ounces)		
	1 %	1.5 %	2 %
1 gallon	1-1/3 oz	2 oz	2-2/3 oz
5 gallons	6-2/3 oz	10 oz	13 oz
10 gallons	13-1/3 oz	20 oz	26 oz
25 gallons	1 quart	1-1/2 quarts	2 quarts

2 tablespoons = 1 fluid ounce

It is not necessary to add additional surfactants or wetting agents.

IMPORTANT: DO NOT side-trim or treat where roots of desirable vegetation may encroach within the treated zone (ie. within two times the drip line). Clean application equipment after using this product by thoroughly flushing with water. DO NOT exceed recommended dosage rate per acre.

WEEDS CONTROLLED

CONTAIN herbicide will provide postemergence control with residual control of the following target vegetation species at the rates listed. CONTAIN herbicide should be used only in accordance with the recommendation on this label and the leaflet label.

GRASSES

Apply 4-8 pints per acre (1% solution)*

Annual bluegrass (*Poa annua*)
Broadleaf signalgrass (*Brachiaria platyphylla*)
Canada bluegrass (*Poa compressa*)
Downy brome (*Bromus tectorum*)
Fescue (*Festuca* spp.)
Foxtail (*Setaria* spp.)
Italian ryegrass (*Lolium multiflorum*)
Johnsongrass (*Sorghum halepense*)
Kentucky bluegrass (*Poa pratensis*)
Lovegrass (*Eragrostis* spp.)
Orchardgrass (*Dactylis glomerata*)
Paragrass (*Brachiaria mutica*)
Quackgrass (*Agropyron repens*)
Saltgrass (*Distichlis stricta*)
Sandbur (*Cenchrus* spp.)
Smooth brome (*Bromus inermis*)
Vaseygrass (*Paspalum urvillei*)
Wild oats (*Avena fatua*)
Witchgrass (*Panicum capillare*)

Apply 6-8 pints per acre (1.5% solution)*

Beardgrass (*Andropogon* spp.)
Cheat (*Bromus secalinus*)
Crabgrass (*Digitaria* spp.)
Fall panicum (*Panicum dichotomiflorum*)
Goosegrass (*Eleusine indica*)
Prairie threeawn (*Aristida oligantha*)
Reed canary grass (*Phalaris arundinacea*)
Torpedograss (*Panicum repens*)
Wild barley (*Hordeum* spp.)

Apply 8-12 pints per acre (2% solution)*

Bahagrass (*Paspalum notatum*)
Bermudagrass (*Cynodon dactylon*)
Big bluestem (*Andropogon gerardi*)
Cattail (*Typha* spp.)
Oatgrass (*Paspalum distatum*)
Feathertop (*Pennisetum villosum*)
Guineagrass (*Panicum maximum*)
Phragmites (*Phragmites australis*)
Prairie cordgrass (*Spartina pectinata*)
Sand dropseed (*Sporobolus cryptandrus*)
Timothy (*Phleum pratense*)
Wrestem muhly (*Muhlenbergia frondosa*)

BROADLEAF WEEDS

Apply 4-8 pints per acre (1% solution)*

Burdock (*Arctium* spp.)
Camphorweed (*Heterotheca subaxillaris*)
Carpetweed (*Mullugo verticillata*)
Carolina geranium (*Geranium carolinianum*)
Clover (*Trifolium* spp.)
Common chickweed (*Stellaria media*)
Common ragweed (*Ambrosia artemisiifolia*)
Dandelion (*Taraxacum officinale*)
Dogfennel (*Eupatorium capillifolium*)
Filaree (*Frodium* spp.)
Fleabane (*Erigeron* spp.)
Hoary vervain (*Verbena stricta*)
Horseweed (*Coryza canadensis*)
Indian mustard (*Brassica juncea*)
Lambsquarters (*Chenopodium album*)
Lespedeza (*Lespedeza* spp.)
Marestail (*Montia perfoliata*)
Mullen (*Verbascum* spp.)
Nettleleaf goosefoot (*Chenopodium murale*)
Oxeye daisy (*Chrysanthemum leucanthemum*)
Peppercorn (*Lepidium* spp.)
Pigweed (*Amaranthus* spp.)
Plantain (*Plantago* spp.)
Smartweed (*Polygonum* spp.)
Sotol (*Ipomoea* spp.)
Sunflower (*Helianthus* spp.)
Sweet clover (*Melilotus* spp.)
Tansymustard (*Descurainia pinnata*)
Western ragweed (*Ambrosia psilostachya*)
Wild carrot (*Daucus carota*)
Wild lettuce (*Lactuca* spp.)
Wild parsnip (*Pastinaca sativa*)
Wild turnip (*Brassica campestris*)
Woollyleaf bursage (*Ambrosia grayi*)
Yellow woodsorrel (*Oxalis stricta*)

Apply 6-8 pints per acre (1.5% solution)*

Bull thistle (*Cirsium vulgare*)
Cocklebur (*Xanthium strumarium*)
Dock (*Rumex* spp.)
Goldenrod (*Solidago* spp.)
Pokeweed (*Phytolacca americana*)
Purslane (*Portulaca* spp.)
Yellow starthistle (*Centaurea solstitialis*)

BROADLEAF WEEDS

Apply 8-12 pints per acre (2% solution)*

Canada thistle (*Cirsium arvense*)
 Giant ragweed (*Ambrosia trifida*)
 Little mallow (*Malva parviflora*)
 Milkweed (*Asclepias* spp.)
 Penstemon (*Penstemon lunifolius*)
 Silver cholla (*Solanum elaeagnifolium*)
 Sowthistle (*Sonchus* spp.)
 Texas thistle (*Cirsium texanum*)

VINES AND BRAMBLES

Apply 4-6 pints per acre (1% solution)

Field bindweed (*Convolvulus arvensis*)
 Hedge bindweed (*Calyptegia sepium*)
 Wild buckwheat (*Polygonum convolvulus*)

Apply 6-8 pints per acre (1.5% solution)**

Greenbrier (*Smilax* spp.)
 Honeysuckle (*Lonicera* spp.)
 Morningglory (*Ipomoea* spp.)
 Poison ivy (*Rhus radicans*)
 Redvine (*Brunnicha ornithosia*)
 Wild rose (*Rosa* spp.)
 Including:
 Multiflora rose (*Rosa multiflora*)
 Macartney rose (*Rosa bracteata*)

Apply 8-12 pints per acre (2% solution)*

Blackberry (*Rubus* spp.)
 Dewberry (*Rubus* spp.)
 Kudzu (*Pueraria lobata*)
 Trumpet creeper (*Campsis radicans*)
 Virginia creeper (*Parthenocissus quinquefolia*)
 Wild grape (*Vitis* spp.)

BRUSH AND TREE SPECIES

Apply 8-12 pints per acre (2% solution)*

Ash (*Fraxinus* spp.)
 Boxelder (*Acer negundo*)
 Cherry (*Prunus* spp.)
 Chinese tallow-tree (*Sapium sebiferum*)
 Dogwood (*Cornus* spp.)
 Hawthorn (*Crataegus* spp.)
 Hickory (*Carya* spp.)
 Maple (*Acer* spp.)
 Mulberry (*Morus* spp.)
 Oak (*Quercus* spp.)
 Persimmon (*Diospyros virginiana*)
 Poplar (*Populus* spp.)
 Privet (*Ligustrum vulgare*)
 Red Maple (*A. rubrum*)
 Salcedo (*Tamarix ramosissima*)
 Sassafras (*Sassafras albidum*)
 Sumac (*Rhus* spp.)
 Sweetgum (*Liquidambar styraciflua*)
 Walnut (*Juglans* spp.)
 Yellow poplar (*Liriodendron tulipifera*)

* The higher rates should be used where heavy or well established infestations occur

** The degree of control is species dependent. Some *Rubus* species may not be controlled

ACCEPTED

MAR 2 1993

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 241-294

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. As such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and


is reasonably fit for the use thereon described when used in accordance with the directions for use, subject to the risks referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF CONTAIN. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product



American Cyanamid Company
 Agricultural Products Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470 ©1992

BEST AVAILABLE COPY




BEST AVAILABLE COPY

60414

ACCEPTED
MAR 2 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-294

CONTAIN™

NON-SELECTIVE
HERBICIDE
FOR NONCROP
FARMSTEAD USE



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION!**

Harmful if absorbed through skin or inhaled. Causes eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

BEST AVAILABLE COPY

70214

In case of emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of CONTAIN should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT mix, store or apply CONTAIN or spray solutions of CONTAIN in unlined steel (except stainless steel) containers or spray tanks.

IMPORTANT

DO NOT use on food or feed crops. DO NOT treat irrigation ditches or water used for crop irrigation or for domestic purposes. Keep from contact with fertilizers, insecticides, fungicides and seeds. DO NOT apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, or similar areas. DO NOT side trim desirable vegetation with this product. Prevent drift of spray to desirable plants. DO NOT USE in California.

Clean application equipment after using this product by thoroughly flushing with water.

GENERAL INFORMATION

CONTAIN herbicide is an aqueous solution to be mixed in water and applied as a spray for the non-selective control of most annual and perennial grasses and broadleaf weeds on noncrop and areas.

CONTAIN may be applied either preemergence or postemergence to the weeds; however, postemergence application is the method of choice in most situations, particularly for control of perennials. For maximum activity, weeds should be growing vigorously at the time of postemergence applications. The preemergence activity of CONTAIN will provide residual control of many weed species following a postemergence application.

CONTAIN is readily absorbed through foliage and roots and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. Treated plants stop growing soon after spray application. Chlorosis appears first in the newest leaves, and necrosis spreads from this point. In perennials, the herbicide is translocated into and kills underground storage organs, thus preventing regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two

BEST AVAILABLE COPY

weeks after application. Complete kill of plants may not occur for several weeks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CONTAIN herbicide should be used only in accordance with recommendations on this label. Keep containers closed to avoid spills and contamination.

A postemergence use of CONTAIN is recommended for control of most annual and perennial grasses, broad-leaf weeds, brush, and trees on NONCROPLAND areas such as fence rows, farm buildings, utility installations, non-irrigation ditch-banks, unimproved roads, and other areas around the farmyard where vegetation is undesirable. DO NOT treat where the roots of desirable vegetation may extend into the treated zone as

death or severe injury may result. As a general guide, roots often extend to within twice the drip line (i.e. twice the branch extremities); however, the roots of some plants may extend beyond this zone. (See diagram below).



STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT store below 32°F. DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in an approved sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPRAYING INSTRUCTIONS

Uniformly apply CONTAIN with a properly calibrated sprayer in sufficient volume to insure adequate coverage. DO NOT ALLOW DRIFT ONTO OR INTO THE VICINITY OF DESIRABLE NON-TARGET PLANTS.

BEST AVAILABLE COPY

90414

HAND-HELD SPRAY EQUIPMENT

Thoroughly mix a 1 to 2 percent solution of CONTAIN in water. To determine the proper percent solution of CONTAIN to use, see the "WEEDS CONTROLLED" section of this label and the "PERCENT SOLUTION RATE GUIDE" below.

PERCENT SOLUTION RATE GUIDE

CONTAIN RATE PER ACRE	PERCENT SOLUTION TO USE
2 - 6 pints	1 %
6 - 8 pints	1.5 %
8 - 12 pints	2 %

For best results, thoroughly cover the foliage of the vegetation to be controlled with the spray solution. DO NOT over apply and cause runoff from the treated foliage.

8

To mix the spray solution, add the volume of CONTAIN indicated in the table below to the desired amount of water.

SPRAY SOLUTION MIXING GUIDE

SOLUTION VOLUME	AMOUNT OF CONTAIN TO USE (fluid ounces)		
	1 %	1.5 %	2 %
1 gallon	2-1/3 oz.	3 oz.	4 oz.
5 gallons	11-2/3 oz.	15 oz.	20 oz.
10 gallons	23-1/3 oz.	30 oz.	40 oz.
25 gallons	1 quart	1-1/2 quarts	2 quarts

2 tablespoons = 1 fluid ounce

It is not necessary to add additional surfactants or wetting agents.

9

IMPORTANT: DO NOT side-trim or treat where roots of desirable vegetation may encroach within the treated zone (i.e. within two times the drip line). Clean application equipment after using this product by thoroughly flushing with water. DO NOT exceed recommended dosage rate per acre.

WEEDS CONTROLLED

CONTAIN herbicide will provide postemergence control with residual control of the following target vegetation species at the rates listed. CONTAIN herbicide should be used only in accordance with the recommendation on this label and the leaflet label.



10

BEST AVAILABLE COPY

100414

GRASSES

Apply 4-6 pints per acre (1% solution)*

Annual bluegrass (*Poa annua*)
 Broadleaf signalgrass (*Brachiaria platphylla*)
 Canada bluegrass (*Poa compressa*)
 Downy brome (*Bromus tectorum*)
 Fescue (*Festuca* spp.)
 Foxtail (*Setaria* spp.)
 Italian ryegrass (*Lolium multiflorum*)
 Johnsongrass (*Sorghum halepense*)
 Kentucky bluegrass (*Poa pratensis*)
 Lovegrass (*Eragrostis* spp.)
 Orchardgrass (*Dactylis glomerata*)
 Paragrass (*Brachiaria mutica*)
 Quackgrass (*Agropyron repens*)
 Saltgrass (*Distichlis stricta*)
 Sandbur (*Cenchrus* spp.)

BEST AVAILABLE COPY

Apply 4-6 pints per acre (1% solution)*

Smooth brome (*Bromus inermis*)
Vaseygrass (*Paspalum urvillei*)
Wild oats (*Avena fatua*)
Witchgrass (*Panicum capillare*)

Apply 6-8 pints per acre (1.5% solution)*

Beardgrass (*Andropogon* spp.)
Cheat (*Bromus tectorum*)
Crabgrass (*Digitaria* spp.)
Fall panicum (*Panicum dichotomiflorum*)
Goosegrass (*Elymus canadensis*)
Prairie threerawn (*Andropogon oligantha*)
Reed canary grass (*Phalaris arundinacea*)
Torpedograss (*Panicum repens*)
Wild barley (*Hordeum* spp.)

Apply 8-12 pints per acre (2% solution)*

Bahiagrass (*Paspalum notatum*)
Bernudagrass (*Cynodon dactylon*)
Big bluestem (*Andropogon gerardi*)
Cattail (*Typha* spp.)
Dallisgrass (*Paspalum dilatatum*)
Feathertop (*Pennisetum violaceum*)
Guinea grass (*Panicum maximum*)
Phragmites (*Phragmites australis*)
Prairie cordgrass (*Spartina pectinata*)
Sand dropseed (*Sporobolus cryptandrus*)
Timothy (*Phleum pratense*)
Wirestem muhly (*Muhlenbergia frondosa*)

BROADLEAF WEEDS

Apply 4-6 pints per acre (1% solution)*

Burdock (*Achium* spp.)
Camphorweed (*Heterotheca subaxillans*)
Carpetweed (*Mullugo verticillata*)
Carolina geranium (*Geranium carolinianum*)
Clover (*Trifolium* spp.)
Common chickweed (*Stellaria media*)
Common ragweed (*Ambrosia artemisiifolia*)
Dandelion (*Taraxacum officinale*)
Dogfennel (*Eupatorium capillifolium*)
Filaree (*Erodium* spp.)
Fleabane (*Erigeron* spp.)
Hairy vervain (*Verbena stricta*)
Horseweed (*Conyza canadensis*)
Indian mustard (*Brassica juncea*)
Lambsquarters (*Chenopodium album*)
Lespedeza (*Lespedeza* spp.)

BEST AVAILABLE COPY

1204/14

Apply 4-6 pints per acre (1% solution)*

Miners lettuce (*Montia perfoliata*)
 Mullein (*Verbascum* spp.)
 Nettleleaf goosefoot (*Chenopodium murale*)
 Oxeye daisy (*Chrysanthemum leucanthemum*)
 Pepperweed (*Lepidium* spp.)
 Pigweed (*Amaranthus* spp.)
 Plantain (*Plantago* spp.)
 Smartweed (*Polygonum* spp.)
 Sorrell (*Rumex* spp.)
 Sunflower (*Helianthus* spp.)
 Sweet clover (*Melilotus* spp.)
 Tansymustard (*Descurainia pinnata*)
 Western ragweed (*Ambrosia psilostachya*)
 Wild carrot (*Daucus carota*)
 Wild lettuce (*Lactuca* spp.)
 Wild parsnip (*Pastinaca sativa*)
 Wild turnip (*Brassica campestris*)
 Woollyleaf bursage (*Ambrosia grayi*)
 Yellow woodsorrel (*Oxalis stricta*)

15

Apply 6-8 pints per acre (1.5% solution)*

Bull thistle (*Cirsium vulgare*)
 Cocklebur (*Xanthium strumarium*)
 Dock (*Rumex* spp.)
 Goldenrod (*Solidago* spp.)
 Pokeweed (*Phytolacca americana*)
 Purslane (*Portulaca* spp.)
 Yellow starthistle (*Centaurea solstitialis*)

Apply 8-12 pints per acre (2% solution)*

Canada thistle (*Cirsium arvense*)
 Giant ragweed (*Ambrosia artemisiifolia*)
 Little mallow (*Malva parviflora*)
 Milkweed (*Asclepias* spp.)
 Pimrose (*Oenothera lutea*)
 Silverleaf nightshade (*Solanum elaeagnifolium*)
 Sowthistle (*Sonchus* spp.)
 Texas thistle (*Cirsium texense*)

16

VINES AND BRAMBLES

Apply 4-6 pints per acre (1% solution)

Field bindweed (*Convolvulus arvensis*)
 Hedge bindweed (*Calystegia sepium*)
 Wild buckwheat (*Polygonum convolvulus*)

Apply 6-8 pints per acre (1.5% solution)*

Greenbrier (*Smilax* spp.)
 Honeysuckle (*Lonicera* spp.)
 Morning glory (*Ipomoea* spp.)
 Poison ivy (*Rhus radicans*)
 Redvine (*Brunnichia cirtiosa*)
 Wild rose (*Rosa* spp.)

Including:

Multiflora rose (*Rosa multiflora*)
 Macartney rose (*Rosa bracteata*)

17

Apply 8-12 pints per acre (2% solution)*

Blackberry (*Rubus* spp.)
 Dewberry (*Rubus* spp.)
 Kudzu (*Pueraria lobata*)
 Trumpet creeper (*Campsis radicans*)
 Virginia creeper (*Parthenocissus vitacea*)
 Wild grape (*Vitis* spp.)

BRUSH AND TREE SPECIES

Apply 8-12 pints per acre (2% solution)*

Ash (*Fraxinus* spp.)
 Boxelder (*Acer negundo*)
 Cherry (*Prunus* spp.)
 Chinese tallow tree (*Sapium*)
 Dogwood (*Cornus florida*)
 Hawthorn (*Crataegus* spp.)
 Hickory (*Carya* spp.)
 Maple (*Acer* spp.)
 Mulberry (*Morus* spp.)

18

BEST AVAILABLE COPY

Apply 8-12 pints per acre (2% solution)*

Blackberry (*Rubus* spp.)
 Dewberry (*Hubus* spp.)
 Kudzu (*Pueraria lobata*)
 Trumpet creeper (*Campsis radicans*)
 Virginia creeper (*Parthenocissus quinquefolia*)
 Wild grape (*Vitis* spp.)

BRUSH AND TREE SPECIES

Apply 8-12 pints per acre (2% solution)*

Ash (*Fraxinus* spp.)
 Boxelder (*Acer negundo*)
 Cherry (*Prunus* spp.)
 Chinese tallow-tree (*Sapium sebiferum*)
 Dogwood (*Cornus* spp.)
 Hawthorn (*Crataegus* spp.)
 Hickory (*Carya* spp.)
 Maple (*Acer* spp.)
 Mulberry (*Morus* spp.)

18

Apply 8-12 pints per acre (2% solution)*

Oak (*Quercus* spp.)
 Persimmon (*Diospyros virginiana*)
 Poplar (*Populus* spp.)
 Privet (*Ligustrum vulgare*)
 Red Maple (*Acer rubrum*)
 Saltcedar (*Tamox ramosissima*)
 Sassafras (*Sassafras albidum*)
 Sumac (*Rhus* spp.)
 Sweetgum (*Liquidambar styraciflua*)
 Willow (*Salix* spp.)
 Yellow poplar (*Citrodendron tulipifera*)

*The higher rates should be used where heavy or well established infestations occur

*The degree of control is species dependent. Some *Rubus* species may not be controlled

19

DISCLAIMER

The label instructions for the use of this product are of experts based on research and field use. It is believed to be reliable and should be followed as it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other consequences may result because of such factors as conditions, presence of other materials, or the use of the product contrary to label instructions. All use is at the control of American Cyanamid Company. All risks are assumed by the user.

American Cyanamid Company shall not be responsible for damages resulting from use of this product in a manner not set forth on this label. User assumes all risks associated with use of this product in any manner not specifically described on the label.

American Cyanamid Company warrants only the product as contained herein conforms to the chemical description and is reasonably fit for the use thereon described in accordance with the directions for use, subject to the

20

BEST AVAILABLE COPY

140414

DISCLAIMER


The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred

to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF CONTAIN. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.



BEST AVAILABLE COPY