


PM 25 241-297
114

EPA Reg. No. 241-297

**FOR USE IN SOYBEANS
USE AREA ADDITIONS**



OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SQUADRON LEAFLET LABEL. SEE SQUADRON LEAFLET LABEL FOR MIXING, SPRAYING, APPLICATION INFORMATION, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

SQUADRON herbicide may be applied in the following Kansas counties:

Jewell, Mitchell, Lincoln, Ellsworth, McPherson, Harvey, Sedgewick, and Sumner.

Note: SQUADRON may also be applied in Kansas, east of U.S. 81 as stated on the SQUADRON leaflet label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.



applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Replanting: If replanting is necessary in field previously treated with SQUADRON, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of SQUADRON.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with liquid fertilizer. Prior to mixing SQUADRON or SQUADRON tank mixtures in liquid fertilizer, refer to Liquid Fertilizer Compatibility section Determinations of this label. While agitating, add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

USE RATES

Apply SQUADRON at a rate of 3.0 pints per acre. See the SQUADRON leaflet label for PROWL and Command use rates in tank-mix combinations.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

APPLICATIONS WITH LIQUID FERTILIZERS

SQUADRON can be applied to the soil in liquid fertilizers, alone or in combination with PROWL or Command. Follow all SQUADRON label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. A spray pressure of 20 to 40 psi is recommended.

When applying SQUADRON in minimum or no-till soybeans, use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply when wind velocity is greater than 10 mph.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the SQUADRON/liquid fertilizer mixture.

**LIQUID FERTILIZER COMPATIBILITY
DETERMINATIONS**

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of SQUADRON alone or with PROWL or Command in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint of fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant* to one jar.
3. (a) When using SQUADRON alone, add to each jar the correct amount of SQUADRON as specified in the following table below.

(b) When using SQUADRON plus PROWL or Command, first add the specified quantity of SQUADRON, and then add the correct amount of PROWL or Command.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
5. Determine compatibility
 - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
 - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.
 - (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
 - (d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

*Complex-Kalo Laboratories Incorporated, Kansas City, MO; Sponto 168 D-Witco Chemical Company, Houston, TX; Unite-Hopkins Agricultural Chemical Company, Madison, WI; or other comparable adjuvants.

4/4

Teaspoons of Specified Herbicide to be
Added to 1 Pint of Liquid Fertilizer Solution

Gallons of Liquid Fertilizer to Be Applied per Acre	PROWL	SQUADRON	Command
20	2 1/2	5	1
30	1 3/4	3 1/2	3/4
40	1 1/4	2 1/2	1/2

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plus², etc.) or imazethapyr (e.g. PURSUIT[®], PURSUIT PLUS[®]) the same year as SQUADRON or injury to follow crops may occur.


DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label should be in the possession of the user at the time of herbicide application

[®]Registered trademark of American Cyanamid Company

²Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E.I. du Pont de Nemours and Company.

PM 75 241-297
1/2



EPA Reg. No. 241-297

FOR USE IN SOYBEANS

SQUADRON APPLIED IN LIQUID FERTILIZER

SQUADRON PLUS PROWL[®] APPLIED IN LIQUID FERTILIZER

SQUADRON PLUS COMMAND¹ APPLIED IN LIQUID FERTILIZER

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SQUADRON, PROWL, AND COMMAND LABELS BEFORE USING. SEE THE SQUADRON LABEL FOR MIXING, SPRAYING AND APPLICATION INFORMATION, USE AREA, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION


SQUADRON alone, SQUADRON plus PROWL, and preplant incorporated applications of SQUADRON plus Command can be applied in liquid fertilizers.

There should be an interval of 90 days between SQUADRON application and soybean harvest. DO NOT feed plants or straw to livestock.

NOTE: DO NOT apply SQUADRON postemergence to soybeans as crop injury may occur.

Crops other than soybeans, such as cotton, corn, grain sorghum, rice and vegetables, may be injured by spray drift or other indirect contact with SQUADRON.

¹Command is a trademark of FMC Corporation



SQUADRON RATE PER ACRE	Fertilizer Rate lbs/acre					
	200	250	300	350	400	450
<u>3 PINTS</u>			<u>PINTS PER TON</u>			
	30	24	20	17	15	13 1/3

For those rates not listed in this table, calculate the pints of SQUADRON to be impregnated on a ton of dry bulk fertilizer using the following formula:


$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times 3 \text{ Pints of SQUADRON per acre (recommended rate)} = \text{Pints of SQUADRON per ton of fertilizer}$$

To impregnate SQUADRON on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of SQUADRON onto the fertilizer during mixing.

Apply the SQUADRON/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The SQUADRON/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

Refer to Preplant Incorporated Applications section on the SQUADRON leaflet label for incorporation directions.


PE-21045 4/4/89

 **CYANAMID**
 American Cyanamid Company
 Agricultural Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470

EPA Reg. No. 241-297

FOR USE IN SOYBEANS

USE AREA ADDITIONS



OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SQUADRON LEAFLET LABEL. SEE SQUADRON LEAFLET LABEL FOR MIXING, SPRAYING, APPLICATION INFORMATION, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

SQUADRON herbicide may be applied in the following Kansas counties:

Jewell, Mitchell, Lincoln, Ellsworth, McPherson, Harvey, Sedgewick, and Sumner.

Note: SQUADRON may also be applied in Kansas, east of U.S. 81 as stated on the SQUADRON leaflet label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.



applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Replanting: If replanting is necessary in field previously treated with SQUADRON, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of SQUADRON.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with liquid fertilizer. Prior to mixing SQUADRON or SQUADRON tank mixtures in liquid fertilizer, refer to Liquid Fertilizer Compatibility section Determinations of this label. While agitating, add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

USE RATES

Apply SQUADRON at a rate of 3.0 pints per acre. See the SQUADRON leaflet label for PROWL and Command use rates in tank-mix combinations.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

APPLICATIONS WITH LIQUID FERTILIZERS

SQUADRON can be applied to the soil in liquid fertilizers, alone or in combination with PROWL or Command. Follow all SQUADRON label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. A spray pressure of 20 to 40 psi is recommended.

When applying SQUADRON in minimum or no-till soybeans, use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply when wind velocity is greater than 10 mph.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the SQUADRON/liquid fertilizer mixture.

LIQUID FERTILIZER COMPATIBILITY
DETERMINATIONS

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of SQUADRON alone or with PROWL or Command in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint of fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant* to one jar.
3. (a) When using SQUADRON alone, add to each jar the correct amount of SQUADRON as specified in the following table below.

(b) When using SQUADRON plus PROWL or Command, first add the specified quantity of SQUADRON, and then add the correct amount of PROWL or Command.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
5. Determine compatibility
 - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
 - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.
 - (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
 - (d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

*Complex-Kalo Laboratories Incorporated, Kansas City, MO; Sponto 168 D-Witco Chemical Company, Houston, TX; Unite-Hopkins Agricultural Chemical Company, Madison, WI; or other comparable adjuvants.

Teaspoons of Specified Herbicide to be
Added to 1 Pint of Liquid Fertilizer Solution

Gallons of Liquid Fertilizer to Be Applied per Acre	PROWL	SQUADRON	Command
20	2 1/2	5	1
30	1 3/4	3 1/2	3/4
40	1 1/4	2 1/2	1/2


DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plus², etc.) or imazethapyr (e.g. PURSUIT[®], PURSUIT PLUS[®]) the same year as SQUADRON or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label should be in the possession of the user at the time of herbicide application.

[®]Registered trademark of American Cyanamid Company

²Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E.I. du Pont de Nemours and Company.



EPA Reg. No. 241-297

FOR USE IN SOYBEANS

SQUADRON APPLIED IN LIQUID FERTILIZER

SQUADRON PLUS PROWL[®] APPLIED IN LIQUID FERTILIZER

SQUADRON PLUS COMMAND¹ APPLIED IN LIQUID FERTILIZER

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SQUADRON, PROWL, AND COMMAND LABELS BEFORE USING. SEE THE SQUADRON LABEL FOR MIXING, SPRAYING AND APPLICATION INFORMATION, USE AREA, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

SQUADRON alone, SQUADRON plus PROWL, and preplant incorporated applications of SQUADRON plus Command can be applied in liquid fertilizers.

There should be an interval of 90 days between SQUADRON application and soybean harvest. DO NOT feed plants or straw to livestock.

NOTE: DO NOT apply SQUADRON postemergence to soybeans as crop injury may occur.

Crops other than soybeans, such as cotton, corn, grain sorghum, rice and vegetables, may be injured by spray drift or other indirect contact with SQUADRON.

¹Command is a trademark of FMC Corporation



SQUADRON RATE PER ACRE	Fertilizer Rate lbs/acre					
	200	250	300	350	400	450
<u>3 PINTS</u>	PINTS PER TON					
	30	24	20	17	15	13 1/3

For those rates not listed in this table, calculate the pints of SQUADRON to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000}{\text{Pounds of dry fertilizer per acre}} \times 3 \text{ Pints of SQUADRON per acre (recommended rate)} = \text{Pints of SQUADRON per ton of fertilizer}$$

To impregnate SQUADRON on bulk fertilizer, use a closed rotary-drum mixer or other commonly used dry bulk fertilizer blender equipped with suitable spray equipment. Spray nozzles must be placed to provide uniform coverage of SQUADRON onto the fertilizer during mixing.

Apply the SQUADRON/dry bulk fertilizer mixture with an accurately calibrated dry fertilizer spreader. The SQUADRON/dry bulk fertilizer mixture must be spread uniformly on the soil surface. Uneven spreading can cause poor weed control and crop injury.

Refer to Preplant Incorporated Applications section on the SQUADRON leaflet label for incorporation directions.


PE-21045 4/4/89



EPA Reg. No. 241-297

FOR USE IN SOYBEANS

USE AREA ADDITIONS



OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SQUADRON LEAFLET LABEL. SEE SQUADRON LEAFLET LABEL FOR MIXING, SPRAYING, APPLICATION INFORMATION, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

SQUADRON herbicide may be applied in the following Kansas counties:

Jewell, Mitchell, Lincoln, Ellsworth, McPherson, Harvey, Sedgewick, and Sumner.

Note: SQUADRON may also be applied in Kansas, east of U.S. 81 as stated on the SQUADRON leaflet label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.



ACCEPTED

MAR - 1 1953

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 241-297

EPA Reg. No. 241-297

FOR USE IN SOYBEANS

SQUADRON APPLIED IN LIQUID FERTILIZER

SQUADRON PLUS PROWL¹ APPLIED IN LIQUID FERTILIZER

SQUADRON PLUS COMMAND¹ APPLIED IN LIQUID FERTILIZER

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SQUADRON, PROWL, AND COMMAND LABELS BEFORE USING. SEE THE SQUADRON LABEL FOR MIXING, SPRAYING AND APPLICATION INFORMATION, USE AREA, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

SQUADRON alone, SQUADRON plus PROWL, and preplant incorporated applications of SQUADRON plus Command can be applied in liquid fertilizers.

There should be an interval of 90 days between SQUADRON application and soybean harvest. DO NOT feed plants or straw to livestock.

NOTE: DO NOT apply SQUADRON postemergence to soybeans as crop injury may occur.

Crops other than soybeans, such as cotton, corn, grain sorghum, rice and vegetables, may be injured by spray drift or other indirect contact with SQUADRON.

¹Command is a trademark of FMC Corporation



To avoid injury to sensitive crops spray equipment used for SQUADRON applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Replanting: If replanting is necessary in field previously treated with SQUADRON, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of SQUADRON.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with liquid fertilizer. Prior to mixing SQUADRON or SQUADRON tank mixtures in liquid fertilizer, refer to Liquid Fertilizer Compatibility section Determinations of this label. While agitating, add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

USE RATES

Apply SQUADRON at a rate of 3.0 pints per acre. See the SQUADRON leaflet label for PROWL and Command use rates in tank-mix combinations.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

APPLICATIONS WITH LIQUID FERTILIZERS

SQUADRON can be applied to the soil in liquid fertilizers, alone or in combination with PROWL or Command. Follow all SQUADRON label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. A spray pressure of 20 to 40 psi is recommended.

When applying SQUADRON in minimum or no-till soybeans, use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply when wind velocity is greater than 10 mph.

All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the SQUADRON/liquid fertilizer mixture.

40FJ

**LIQUID FERTILIZER COMPATIBILITY
DETERMINATIONS**

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of SQUADRON alone or with PROWL or Command in the specific liquid fertilizer to be used according to the following directions:

1. Add 1 pint of fertilizer to each of 2 one-quart jars.
2. Add 1/2 teaspoon of adjuvant* to one jar.
3. (a) When using SQUADRON alone, add to each jar the correct amount of SQUADRON as specified in the following table below.

(b) When using SQUADRON plus PROWL or Command, first add the specified quantity of SQUADRON, and then add the correct amount of PROWL or Command.
4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or other precipitates.
5. Determine compatibility
 - (a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.
 - (b) If the mixture with adjuvant does not separate, but the one without adjuvant separates, use the adjuvant mixture in your spray tank. Add the adjuvant to the liquid fertilizer as directed on the manufacturer's label.
 - (c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.
 - (d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

*Complex-Kalo Laboratories Incorporated, Kansas City, MO; Sponto 168 D-Witco Chemical Company, Houston, TX; Unite-Hopkins Agricultural Chemical Company, Madison, WI; or other comparable adjuvants.

Teaspoons of Specified Herbicide to be
Added to 1 Pint of Liquid Fertilizer Solution

Gallons of Liquid Fertilizer to Be Applied per Acre	PROWL	SQUADRON	Command
20	2 1/2	5	1
30	1 3/4	3 1/2	3/4
40	1 1/4	2 1/2	1/2

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plus², etc.) or imazethapyr (e.g. PURSUIT[®], PURSUIT PLUS[®]) the same year as SQUADRON or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label should be in the possession of the user at the time of herbicide application.

[®]Registered trademark of American Cyanamid Company

²Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E.I. du Pont de Nemours and Company.