

NOT REVISIONED
In Accordance with PR Notice 82-2,
Based on Draft Labeling Dated 3/1/89

EPA Reg. No. 241-297
FOR USE IN SOYBEANS

PM 25 241-297
1/2

SQUADRON APPLIED IN LIQUID FERTILIZER
SQUADRON PLUS PROWL® APPLIED IN LIQUID FERTILIZER
SQUADRON PLUS COMMAND¹ APPLIED IN LIQUID FERTILIZER

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

SQUADRON alone, SQUADRON plus PROWL, and pre-plant incorporated applications of SQUADRON plus Command can be applied in liquid fertilizers.

There should be an interval of 90 days between SQUADRON application and soybean harvest. DO NOT feed plants or straw to livestock.

NOTE: DO NOT apply SQUADRON postemergence to soybeans as crop injury may occur. Crops other than soybeans, such as cotton, corn, grain sorghum, rice and vegetables, may be injured by spray drift or other indirect contact with SQUADRON.

To avoid injury to sensitive crops, spray equipment used for SQUADRON applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Replanting: If replanting is necessary in field previously treated with SQUADRON, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of SQUADRON.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with liquid fertilizer. Prior to mixing SQUADRON or SQUADRON tank mixtures in liquid fertilizer, refer to Liquid Fertilizer Compatibility Determinations section of this label. While agitating, add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

¹Command is a trademark of EMC Corporation
*Registered Trademarks of American Cyanamid Company

USE RATES

Apply SQUADRON at a rate of 3.0 pints per acre. See the SQUADRON leaflet label for PROWL and Command use rates in tank-mix combinations.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops. Avoid overlaps when spraying.

APPLICATIONS WITH LIQUID FERTILIZERS

SQUADRON can be applied to the soil in liquid fertilizers, alone or in combination with PROWL or Command. Follow all SQUADRON label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. A spray pressure of 20 to 40 psi is recommended.

When applying SQUADRON in minimum or no-till soybeans, use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply when wind velocity is greater than 10 mph. All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the SQUADRON/liquid fertilizer mixture.

**LIQUID FERTILIZER
COMPATIBILITY DETERMINATIONS**

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of SQUADRON alone or with PROWL or Command in the specific liquid fertilizer to be used according to the following directions:

BEST AVAILABLE COPY

correct amount of SQUADRON as specified in the table below.

(b) When using SQUADRON plus PROWL or Command, first add the specified quantity of SQUADRON, then add the correct amount of PROWL or Command.

4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or their precipitates.

5. Determine compatibility

(a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.

(c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.

(d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

*Compex-Kalo Laboratories Inc., Kansas City, MO; Sponto 168 D-Witco Chemical Co., Houston, TX; Unite-Hopkins Agricultural Chemical Co., Madison, WI; or other comparable adjuvants.

Teaspoons of Specified Herbicide to be Added to 1 Pint of Liquid Fertilizer Solution

Gallons of Liquid Fertilizer to Be Applied per Acre	PROWL	SQUADRON	Command
20	2/3	2	1/3
30	1/2	1 1/2	1/4
40	1/3	1	1/6

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plus², etc.) or imazethapyr (e.g. PURSUIT[®], PURSUIT[®] PLUS) the same year as SQUADRON or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label should be in the possession of the user at the time of herbicide application.

²Classic, Canopy, Preview, Gemini and Lorox Plus are trademarks of E. I. du Pont de Nemours and Co.

*Registered Trademarks of American Cyanamid Company

PL 21914 3/21/89

 **CYANAMID**
American Cyanamid Company
New York, N.Y. 10017
The following is a list of Cyanamid's
offices in the United States:

PM 25 241-297
1/1

EPA Reg. No. 241-297

REVIEWED
with PR Notice 82-2
Draft Labeling Dated 3/1/81

FOR USE IN SOYBEANS

USE AREA ADDITIONS

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SQUADRON LEAFLET LABEL. SEE SQUADRON LEAFLET LABEL FOR MIXING, SPRAYING, APPLICATION INFORMATION, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

SQUADRON herbicide may be applied in the following Kansas counties:

Jewell, Mitchell, Lincoln, Ellsworth, Republic, Cloud, Ottawa, Saline, McPherson, Harvey, Sedgewick, and Sumner.

Note: SQUADRON may also be applied in Kansas, east of U.S. 81 as stated on the SQUADRON leaflet label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.



NOT REVIEWED
In Accordance with FR Notice 82-74
Based on Draft Labeling Dated 3/1/89

EPA Reg. No. 241-297
FOR USE IN SOYBEANS

SQUADRON APPLIED IN LIQUID FERTILIZER
SQUADRON PLUS PROWL® APPLIED IN LIQUID FERTILIZER
SQUADRON PLUS COMMAND¹ APPLIED IN LIQUID FERTILIZER

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system.

GENERAL INFORMATION

SQUADRON alone, SQUADRON plus PROWL, and pre-plant incorporated applications of SQUADRON plus Command can be applied in liquid fertilizers.

There should be an interval of 90 days between SQUADRON application and soybean harvest. DO NOT feed plants or straw to livestock.

NOTE: DO NOT apply SQUADRON postemergence to soybeans as crop injury may occur. Crops other than soybeans, such as cotton, corn, grain sorghum, rice and vegetables, may be injured by spray drift or other indirect contact with SQUADRON.

To avoid injury to sensitive crops, spray equipment used for SQUADRON applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops.

Replanting: If replanting is necessary in field previously treated with SQUADRON, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. Do not apply a second treatment of SQUADRON.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with liquid fertilizer. Prior to mixing SQUADRON or SQUADRON tank mixtures in liquid fertilizer, refer to Liquid Fertilizer Compatibility Determinations section of this label. While agitating, add the required amount of product and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

¹Command is a trademark of FMC Corporation

®Registered Trademarks of American Cyanamid Company

USE RATES

Apply SQUADRON at a rate of 3.0 pints per acre. See the SQUADRON leaflet label for PROWL and Command use rates in tank-mix combinations.

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops. Avoid overlaps when spraying.

APPLICATIONS WITH LIQUID FERTILIZERS

SQUADRON can be applied to the soil in liquid fertilizers, alone or in combination with PROWL or Command. Follow all SQUADRON label recommendations regarding incorporation, timing of application, special instructions and precautions. Apply treatments in 20 or more gallons of liquid fertilizer per acre with ground equipment. A spray pressure of 20 to 40 psi is recommended.

When applying SQUADRON in minimum or no-till soybeans, use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply when wind velocity is greater than 10 mph. All individual state regulations relating to fluid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company selling the SQUADRON/liquid fertilizer mixture.

**LIQUID FERTILIZER
COMPATIBILITY DETERMINATIONS**

If a liquid fertilizer and herbicide(s) mixture separates in the spray tank, clogged equipment and uneven application can result, which can cause poor weed control and crop injury. Always predetermine the compatibility of SQUADRON alone or with PROWL or Command in the specific liquid fertilizer to be used according to the following directions:

correct amount of SQUADRON as specified in the table below.

(b) When using SQUADRON plus PROWL or Command, first add the specified quantity of SQUADRON, then add the correct amount of PROWL or Command.

4. Close both jars and shake thoroughly for 10 seconds. Let them stand for 30 minutes and then observe the results. Look for signs of separation, an oily layer or globules, sludge, flakes or their precipitates.

5. Determine compatibility

(a) If the mixture without adjuvant does not separate, use this mixture in your spray tank.

(c) If either mixture separates, but mixes readily with shaking, the mixture can be used providing good agitation is maintained in the spray tank.

(d) If separation of the mixture occurs, and agitation and/or adjuvant does not correct this problem, DO NOT use the herbicide(s) in that specific liquid fertilizer.

*Complex-Kato Laboratories Inc., Kansas City, MO; Sponto 168 D-Witco Chemical Co., Houston, TX; Unite-Hopkins Agricultural Chemical Co., Madison, WI; or other comparable adjuvants.

Teaspoons of Specified Herbicide to be Added to 1 Pint of Liquid Fertilizer Solution

Gallons of Liquid Fertilizer to Be Applied per Acre	PROWL	SQUADRON	Command
20	2/3	2	1/3
30	1/2	1 1/2	1/4
40	1/3	1	1/6

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plus², etc.) or imazethapyr (e.g. PURSUIT[®], PURSUIT[®] PLUS) the same year as SQUADRON or injury to follow crops may occur.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label should be in the possession of the user at the time of herbicide application.

BEST AVAILABLE COPY

Classic, Canopy, Preview, Gemini and Lorox Plus are
trademarks of I.B.F. Co. Pont De Nemours and Co.
Registered Trademarks of American Cyanamid Company

RE-21044 3/21/89



CYANAMID

American Cyanamid Company
Agricultural Division
Crop Protection Chemical Department
Wayne, NJ 07470

EPA Reg. No. 241-297

FOR USE IN SOYBEANS

USE AREA ADDITIONS

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS ON THE SQUADRON LEAFLET LABEL. SEE SQUADRON LEAFLET LABEL FOR MIXING, SPRAYING, APPLICATION INFORMATION, WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS.

SQUADRON herbicide may be applied in the following Kansas counties:

Jewell, Mitchell, Lincoln, Ellsworth, Republic, Cloud, Ottawa, Saline, McPherson, Harvey, Sedgewick, and Sumner.


Note: SQUADRON may also be applied in Kansas, east of U.S. 81 as stated on the SQUADRON leaflet label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.

 **CYANAMID**
American Cyanamid Company
Agricultural Division
Crop Protection Chemical Department
Wayne, NJ 07470

3/21/89

REVIEWED
Accepted with PR Notice 82-
sd OH Draft Labeling Dated

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200