EPA Reg. No. 241-297

REDUCED RATE OF SQUADRON PLUS PROWL® HERBICIDE FOR USE IN SOYBEANS

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, LOUISIANA, MISSISSIPPI, NORTH CAROLINA, OKLAHOMA, SOUTH CAROLINA, TENNESSEE, TEXAS AND THE MISSOURI BOOTHEEL (BUTLER, CAPE GIRARDEAU, DUNKLIN, MISSISSIPPI, NEW MADRID, PEMISCOT, SCOTT AND STODDARD COUNTIES).

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SQUADRON AND PROWL LABELS BEFORE USING. SEE THE SQUADRON LABEL FOR USE AREA, MIXING, SPRAYING AND APPLICATION INFORMATION, AND ROTATIONAL CROP RESTRICTIONS. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SQUADRON may be applied preplant incorporated or preemergence as stated on the SQUADRON leaflet label.

DO NOT apply this product through any type of irrigation system.


USE RATES

From 7 days prior to planting soybeans up to 2 days after planting, SQUADRON may be applied at a rate of 2 1/4 pints per acre on coarse and medium textured soils. PROWL must be tank mixed with the 2 1/4 pint rate of SQUADRON, Add PROWL at a rate of 7/8 pint per acre.

DO NOT apply SQUADRON at rates below 3 pints per acre on fine (silty clay loams, silty clays, clay loams, clays) textured soils.

DO NOT apply SQUADRON at rates below 2 1/4 pints per acre on coarse (sands, loamy sands, sandy loams) and medium (sandy clay loams, sandy clays, loams, silt loams, silts) textured soils.

*Registered Trademark of American Cyanamid Company



-1-

Barnyardgrass (Echinochloa crus-galli) Johnsongrass, seedling (Sorghum halepense) Crabgrass (Digitaria spp.) **Panicum** Browntop (Panicum fásciculatum) (Dactyloctenium aegyptium) Crowfootgrass (Panicum dichotomiflorum) Fall Foxtail Texas (Panicum texanum) Giant (Setaria faberi) (Cenchrus incertus) (Setaria viridis) Sandbur, Field Green (Brachiaria platyphylla) Robust (Setaria sp.) Signalgrass, Broadleaf (Panicum capillare) Witchgrass (Setaria glauca) Yellow

Broadleaf weeds

Goosegrass

Smooth

Pigweed (continued) (Mollugo verticillata) Carpetweed Waterhemp, tall (Amaranthus tuberculatus) (Xanthium strumarium) Cocklebur, Common* Lambsquarters, Common (Chencoodium album) Purslane (Portulaça oleracea) Pigweed* Pusley, Florida (Richardia scabra) Sida, Prickly (Teaweed) Palmer (Amaranthus palmeri) (Sida spinosa) (Amaranthus retroflexus) Smartweed Redroot

Ladysthumb

Spiny (Amaranthus spinosus) Pennsylvania
SOUADRON plus PROWL will aid in the control and reduce competition from:

(Amaranthus hybridus)

(Eleusine indica)

Morningglory**	Spurge		
Palm Leaf	(Ipomoea wrightii)	Annual	(Euphorbia sp.)
Pitted	(Ipomoea lacunosa)	Prostrate	(Euphorbia humistrata)
Smallflower	(Ĵacquemontia tamnifolia)	Spotted	(Euphorbia maculata)

^{*}A sequential postemergence application of SCEPTER® or SCEPTER® O.T.® may be required to control heavy infestations of cocklebur or pigweed. See the SCEPTER or SCEPTER O.T. label for use directions.

SEQUENTIAL APPLICATIONS OF SCEPTER OR SCEPTER O.T. ARE NOT PERMITTED IN THE MISSOURI BOOTHEEL.

Use of SQUADRON herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plur², etc.), imazaquin (e.g. ALA-SCEPT®, SCEPTER®, TRI-SCEPT®, etc. [refer to Sequential Program section of the GQUADRON leaflet label for exceptions]), or imazethapyr (e.g. PURSUIT® or PURSUIT® PLUS) the same year as SQUADRON or injury to follow crops may occur.

This label must be in the possession of the user at the time of herbicide application.

*Classic, Cirropy, Preview, Geouss and Lorot. Plus are syndemetry of E.L. &s Post & Nestours and Company *Regimend Tradeback of American Cysnamid Company

PE-21052 3/1/90





(Polygonum persicaria)
(Polygonum pensylvanicum)

[&]quot;Commercial control of listed morningglory species will be achieved with a sequential application of SCEPTER O.T. See the SCEPTER O.T. label for use directions.