

**ACCEPTED**

28

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-297

**EPA Reg. No. 241-297****FOR USE ON SOYBEANS****FOR DISTRIBUTION AND USE ONLY IN INDIANA, OHIO, AND KENTUCKY****FOR CONTROL OF EASTERN BLACK NIGHTSHADE AND GIANT RAGWEED**

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SQUADRON LABEL. SEE THE SQUADRON LABEL FOR USE AREA, MIXING, SPRAYING AND APPLICATION INFORMATION, AND ROTATIONAL CROP RESTRICTIONS. WHEN TANK MIXING, ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of herbicide application. DO NOT apply this product through any type of irrigation system.

SQUADRON applied preemergence as directed on the SQUADRON leaflet label will control the following weeds in addition to the weeds labeled for control on the SQUADRON label.

Eastern black nightshade  
Giant ragweed

*(Solanum ptycanthum)*  
*(Ambrosia trifida)*

Large seeded weeds such as Eastern black nightshade and giant ragweed germinating deep in the soil or with subsequent flushes may require a cultivation or postemergence herbicide application for season long control.

Preemergence applications of SQUADRON will aid in the control and reduce competition from:

Velvetleaf

*(Abutilon theophrasti)*

Additional PROWL® herbicide may be required for certain soils. For rates, refer to the SQUADRON plus Additional PROWL herbicide Tank Mixture section of the SQUADRON Label.

Rainfall (approx. 1 inch) is required to activate surface applied herbicides such as SQUADRON before weed emergence. If adequate moisture is not received within seven days after application, a thorough shallow tillage or cultivation (preferably with a rotary hoe) may improve weed control.

To kill existing vegetation, SQUADRON may be tank mixed with paraquat, Roundup<sup>1</sup>, or 2,4-D low volatile ester. DO NOT apply 2,4-D low volatile ester within 30 days of planting. Consult labels for rates, use instructions, precautionary statements and restrictions. Always follow the most restrictive label. Maintain constant agitation of spray mixture.

Use of SQUADRON herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

<sup>1</sup>Roundup is a trademark of Monsanto Agricultural Products Company.

<sup>2</sup>Registered Trademark of American Cyanamid Company

PE-21067 2/21/81 DL

**CYANAMID**

American Cyanamid Company  
Agricultural Division  
Crop Protection Chemicals Department  
Wayne, NJ 07478 ©1981