


P125  
241-303  
pg 4 of 8

**IMAGE® 1.5 LC  
herbicide**



**A Selective Herbicide for Use in Established:**

- |                |                    |
|----------------|--------------------|
| Bermudagrass   | St. Augustinegrass |
| Centipedegrass | Zoysiagrass        |

**ACTIVE INGREDIENT:**

Ammonium salt of imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)*.....	17.3%
INERT INGREDIENTS.....	82.7%
TOTAL.....	100.0%

\*Equivalent to 16.4% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid.

(1 gallon contains 1.5 pounds of active ingredient as the free acid)

EPA Reg. No. 241-303

EPA Est. No. 5905-AR-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION!**

**PRECAUCION!**

**PRECAUCION AL USUARIC:** Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada amplimente.

See Additional Precautionary Statements Inside.

In case of emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

AMERICAN CYANAMID COMPANY  
AGRICULTURAL DIVISION  
VPC VENTURES  
WAYNE, NJ 07470 ©1990

Net Contents:

•Registered Trademark of American Cyanamid Company

6/90

2078

**STATEMENT OF PRACTICAL TREATMENT**

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**

**CAUTION!**

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**ENVIRONMENTAL HAZARDS**

DO NOT apply directly to water or wetlands (swamps, bogs, marshes and potholes).

DO NOT contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Observe all cautions and limitations on this label and on the labels of products used in combination with IMAGE. The use of IMAGE not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** KEEP FROM FREEZING.  
DO NOT store below 32°F.  
DO NOT contaminate water, food, or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.


**DISCLAIMER**

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

Any damages arising from breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY.



**GENERAL INFORMATION**

IMAGE is a unique, versatile herbicide that will control or aid in the control of many important weeds in warm-season turfgrasses. IMAGE may be used at weed control rates on established bermudagrasses, centipedegrasses, St. Augustinegrasses and zoysiagrasses. IMAGE will effectively control susceptible weeds when applied during the recommended application timings (pre and post). Postemergent control of weeds may require several weeks.

**HERBICIDE COMBINATIONS**


IMAGE may be tank-mixed with registered preemergence and broadleaf postemergence herbicides and with methylarsonate herbicides, such as MSMA, to control weeds not controlled by IMAGE alone. DO NOT tank-mix IMAGE with other postemergence herbicides labeled for grass weed control in turf. IMAGE may be applied prior to or sequential to the use of all currently labeled preemergence and postemergence herbicides.

When IMAGE is used in tank-mixture with another herbicide, refer to each label for rates, methods of application, proper timing, weeds controlled, limitations and precautions. ALWAYS USE in accordance with the more restrictive label limitations and precautions. IMAGE cannot be mixed with any product containing a label prohibition against such mixing.

**MIXING INSTRUCTIONS**

Fill the spray tank one-half to three-quarters full with clean water. While agitating, add the required amount of IMAGE and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank-mixing IMAGE with recommended herbicides, add the other herbicide to the spray tank first and make sure it is thoroughly mixed before adding IMAGE.



Soy?

### ADDITION OF SURFACTANTS

A nonionic surfactant at the rate of 2 pints per 100 gallons (0.25% v/v) should be added to the spray mixture. Add the nonionic surfactant to the spray mixture after herbicide(s) have mixed. An antifoaming agent may be added to the tank, if needed.

### SPRAYING INSTRUCTIONS

Uniformly apply IMAGE sprays with a properly calibrated sprayer in sufficient volume to insure adequate coverage (40-200 gallons/acre). AVOID drift onto vegetables, flowers, ornamental shrubs and other desirable plants or injury may result. AVOID overlaps when spraying. A spray indicator dye may be added to prevent overlaps. DO NOT apply when wind velocity is greater than 10 mph.

### SPECIAL PRECAUTIONS

USE ONLY on well established non-stressed turfgrass with a dense and uniform stand. TEMPORARY YELLOWING may occur.

DO NOT use on golf or bowling greens.

DO NOT use on unlabeled turfgrass species, including dichondra.

DO NOT mow St. Augustinegrasses until 48 hours after application.

DO NOT use on tall fescue or mixed stands of tall fescue and bermudagrass when tall fescue is a desirable turfgrass. IMAGE may injure tall fescue.

DO NOT reseed or winter overseed or sprig turfgrass for 1-1/2 months after treatment.


DO NOT apply to newly planted, sprigged, or sodded turfgrass.

DO NOT use on winter overseeded turfgrasses such as ryegrass, bentgrass or Poa trivialis when those species are desirable.

DO NOT use on ornamentals as injury may result.

DO NOT graze or feed clippings of treated turfgrasses.

DO NOT apply aerially.



6078

IMAGE USE RATES

IMAGE should be applied at 1 to 1 1/3 qts. per acre (0.75 to 1 fl. oz. per 1000 sq. ft.). The high rate should be used where heavy weed infestations occur. Irrigation or rainfall of 0.5 inch is suggested within 1-7 days after application to wash IMAGE into the shoot/root zone where greatest weed control activity occurs. For spot spraying, mix 2 fl. oz. of IMAGE per 3 gallons of water and spray to wet.

SPRAY SOLUTION MIXING GUIDE

Spray Volume (gal/A)	Quarts of IMAGE to Mix Per 100 Gallons of Water	
	0.75 oz/1000	1.0 oz/1000
20	5	6-3/4
43	2-1/4	3-1/8
100	1	1-1/4
172	1/2	3/4

CONTROL OF SUMMER WEEDS

For the postemergence control of weeds such as the nutsedges, dollarweed, kyllinga, field sandbur, and ryegrass, applications should be timed following spring transition of the turf and prior to the onset of winter dormancy. This application timing would be the period of active turf and summer weed growth. Applications should not be made just prior to or during transition or during periods of very slow turf growth or severe discoloration could occur.

WEED SPECIES CONTROLLED

GRASSES CONTROLLED:

Field Sandbur (Cenchrus incertus)

SEDGES CONTROLLED:

- Globe Sedge (Cyperus globosus)
- Green Kyllinga (Kyllinga brevifolia)
- Rice Flatsedge (Cyperus iria)
- Purple Nutsedge (Cyperus rotundus)
- Yellow Nutsedge (Cyperus esculentus)

IMAGE will aid in the control of and reduce competition from:

Dollarweed (Hydrocotyle umbellata)

Notes:

- 1) For control of kyllinga and more rapid and improved control of the sedges the addition of MSMA (2 lb. ai/A) is suggested in methylarsonate tolerant turfgrasses.
- 2) A repeat application may be required for extended postemergence control of weeds such as dollarweed, kyllinga, the nutsedges, and perennial ryegrass.
- 3) Tolerant turfgrasses detoxify IMAGE. Best tolerance occurs under totally dormant or optimal growing conditions. When the turfgrasses are growing under marginal conditions, such as low temperatures, drought stress, compaction, etc., temporary yellowing or browning can occur and recovery may take 2-4 weeks.

CONTROL OF WINTER WEEDS

For the control of weeds such as wild garlic, wild onion, henbit, chickweed, lawn burweed, etc., applications should be timed to follow the first killing frost. IMAGE may be applied prior to or soon after emergence of the weeds listed below with the exception of wild onion and wild garlic which must be emerged at application time for control.

WEEDS SPECIES CONTROLLED

GRASSES CONTROLLED:

Perennial Ryegrass - overseeded (Lolium perenne)

BROADLEAVES CONTROLLED:

- Black Medic (Medicago lupulina)
- Buttercup (Ranunculus parviflorus)
- Chickweed
  - Common (Stellaria media)
  - Mouseear (Cerastium vulgatum)
- Cutleaf Evening Primrose (Oenothera lanceolata)
- Geranium
  - Carolina (Geranium carolinium)
  - Dovetail (Geranium molle)
- Hairy bittercress (Cardamine hirsuta)
- Henbit (Lamium amplexicaule)
- Knawel (Scelranthus annuus)
- Lawn burweed (Soliva pterosperma)
- Parsley-piert (Alchemilla arvensis)
- Purple deadnettle (Lamium purpureum)
- Red sorrel (Rumex acetosella)
- White clover (Trifolium repens)
- Wild garlic (Allium vineale)
- Wild onion (Allium canadense)

