

1105 241-303 104

IMAGE[®]

PULL HERE TO OPEN
PRESS TO RESEAL

1.5 LC herbicide

A Selective Herbicide for Use in Established:
Bermudagrass Centipedegrass St. Augustinegrass
Selected Landscape Ornamentals Zoysiagrass

ACTIVE INGREDIENT

INERT INGREDIENTS

TOTAL

EPA Reg. No. 241-303

EPA Est. No. 5905 AR 1

**KEEP OUT OF REACH OF CHILDREN
CAUTION! / PRECAUCION!**


PRECAUCION AL USUARIO



Net Contents 1 gallon (3.78 liters)

Registered Trademark of American Cyanamid Company

12 91



BEST AVAILABLE COPY

IMAGE

1.5 LC herbicide

A Selective Herbicide for Use in Established:

Bermudagrass Centipedegrass St. Augustinegrass
Selected Landscape Ornamentals Zoysiagrass

ACTIVE INGREDIENT: Ammonium salt of imazaquin
(2,4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazo-2,1-f-3-quinolinecarboxylic acid) - 17.3%

INERT INGREDIENTS 82.7%


TOTAL 100.0%

Contains 1.5 pounds of active ingredient as the free acid.
EPA Reg. No. 241-303 EPA Est. No. 5905-19-1

**KEEP OUT OF REACH
OF CHILDREN
CAUTION!
¡PRECAUCION!**

PRECAUCION AL USUARIO: Si usted no lee inglés
use este producto hasta que la etiqueta le haya
sido explicada ampliamente

In case of emergency endangering life or property
involving this Product, call collect, day or night,
Area Code 201-835-3100



Net Contents: 1 gallon (3.78 liters)
Registered Trademark of American Cyanamid Company

12/91

STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

Observe all cautions and limitations on this label and on the labels of products used in combination with IMAGE. The use of IMAGE not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

See Leaflet for Complete Directions For use.

STORAGE AND DISPOSAL

PROHIBITIONS: KEEP FROM FREEZING. DO NOT store below 32°F. DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning, if burned, stay out of smoke.

See Leaflet for Disposal.

STATEMENT OF PRACTICAL TREATMENT

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION!

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

Observe all cautions and limitations on this label and on the labels of products used in combination with IMAGE. The use of IMAGE not consistent with this label may result in plant injury. Keep container closed to avoid spills and contamination.

STORAGE AND DISPOSAL

PROHIBITIONS: KEEP FROM FREEZING. DO NOT store below 32°F. DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, insect vermin, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond control of American Cyanamid Company. All such risks shall be assumed by the user.


American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above.

Any damages arising from breach of this warranty shall be limited to direct damages and shall not include consequential or commercial damages such as loss of profits or values or any other special or indirect damages.

American Cyanamid Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY.

GENERAL INFORMATION

IMAGE is a unique, versatile herbicide that will control or aid in the control of many important weeds.



in warm-season turfgrasses and selected landscape ornamentals. IMAGE may be used at weed control rates on established bermudagrasses, centipedegrasses, St. Augustinegrasses, zoysicgrasses and listed ornamentals. IMAGE will effectively control susceptible weeds when applied during the recommended application timings (pre and post). Postemergence control of weeds may require several weeks.

HERBICIDE COMBINATIONS (Warm-season turfgrasses)

IMAGE may be tank-mixed with registered preemergence and broadleaf postemergence herbicides and with methylarsenate herbicides, such as MSMA, to control weeds not controlled by IMAGE alone. DO NOT tank-mix IMAGE with other postemergence herbicides labeled for grass weed control in turf. IMAGE may be applied prior to or sequentially to the use of all currently labeled preemergence and postemergence herbicides.

(Landscape ornamentals)

IMAGE may be tank mixed with pendimethalin or oryzalin to control weeds not controlled by IMAGE alone.

When IMAGE is used in tank mixture with another herbicide, refer to each label for rates, methods of application, proper timing, weeds controlled, limitations and precautions. ALWAYS USE in accordance with the more restrictive label limitations and precautions. IMAGE cannot be mixed with any product containing a label prohibition against such mixing.

MIXING INSTRUCTIONS

Fill the spray tank one-half to three-quarters full with clean water. While agitating, add the required amount of IMAGE and then fill the remainder of the tank with water. Maintain agitation while spraying to insure a uniform spray mixture.

When tank-mixing IMAGE with recommended herbicides, add the other herbicide to the spray tank first and make sure it is thoroughly mixed before adding IMAGE.

ADDITION OF SURFACTANTS

A nonionic surfactant at the rate of 2 pints per 100 gallons (0.25% v/v) should be added to the spray mixture. Add the nonionic surfactant to the spray mixture after herbicide(s) have mixed. An antifoaming agent may be added to the tank, if needed.

SPRAYING INSTRUCTIONS

Uniformly apply IMAGE sprays with a properly calibrated sprayer in sufficient volume to insure adequate coverage (40-200 gallons/acre). AVOID drift onto vegetables, flowers, and other desirable plants or injury may result. AVOID overlaps when spraying. A spray indicator dye may be added to prevent overlaps. DO NOT apply when wind velocity is greater than 10 mph.

SPECIAL PRECAUTIONS (Warm-season turfgrasses)

USE ONLY on well established non-stressed turf grass with a dense and uniform stand. TEMPORARY YELLOWING may occur.

DO NOT use on golf or bowling greens.

DO NOT use on unlabeled turfgrass species, including dichondra.

DO NOT mow St. Augustinegrasses until 48 hours after application.

DO NOT use on tall fescue or mixed stands of tall fescue and bermudagrass when tall fescue is a desirable turfgrass. IMAGE may injure tall fescue.

DO NOT reseed or winter overseed or spring turf grass for 1-12 months after treatment.

DO NOT apply to newly planted, sprigged, or sodded turfgrasses.

DO NOT use on winter overseeded turfgrasses such as ryegrass, bentgrass or *Poa trivialis* when those species are desirable.

DO NOT graze or feed clippings of treated turfgrasses.

DO NOT apply aerially.

SPECIAL PRECAUTIONS (Landscape ornamentals)

USE ONLY on well-established non-stressed ornamentals or injury may result.

DO NOT apply to unlabeled ornamentals or annual bedding plants.

DO NOT use on container grown ornamental nursery stock.

DO NOT apply in the rooting area of unlabeled ornamentals or annual bedding plants or plant injury may result.

DO NOT apply in vegetable gardens.

DO NOT apply to soil where annual bedding plants may be planted the following year or plant injury may result.

IMAGE USE RATES

IMAGE should be applied at 1 to 1 1/3 qts. per acre (0.75 to 1 fl. oz. per 1000 sq. ft.). The high rate should be used where heavy weed infestations occur. Irrigation or rainfall of 0.5 inch is suggested within 1-7 days after application to wash IMAGE into the shoot root zone where greatest weed control activity occurs.

ACCEPTED

APR 17 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 344-505

SPRAY SOLUTION MIXING GUIDE

Spray Volume (gal/A)	Quarts of IMAGE to Mix Per 100 Gallons of Water	
	0.75 oz/1000 sq. ft.	1.0 oz/1000 sq. ft.
20	5	6-3/4
43	2-1/4	3-1/8
100	1	1-1/4
172	1/2	3/4

SPOT SPRAYING: For hand-held or back-pack sprayer, mix 2 fl. oz. of IMAGE per 3 gallons of water and spray to wet. For hose end sprayer (i.e. Ortho Spray-ette 4) add 3/4 fl. oz. of IMAGE to hose end container, fill with water to two gallon mark and apply over 1000 sq. ft. (the length in ft. times the width in ft. = 1000 sq. ft. in area).

TURFGRASSES

CONTROL OF SUMMER WEEDS

For the postemergence control of weeds such as the nutsedges, dollarweed, kyllinga, field sandbur, and ryegrass, applications should be timed following spring transition of the turf and prior to the onset of winter dormancy. This application timing would be the period of active turf and summer weed growth. Applications should not be made just prior to or during transition or during periods of very slow turf growth or severe discoloration could occur.

WEED SPECIES CONTROLLED

GRASSES CONTROLLED:

Field Sandbur (*Cenchrus incertus*)
Perennial Ryegrass - overseeded
(*Lolium perenne*)

SEDGES CONTROLLED:

Globe Sedge (*Cyperus globosus*)
Green Kyllinga (*Kyllinga brevifolia*)
Rice Flatsedge (*Cyperus iria*)

BEST AVAILABLE COPY

BEST AVAILABLE COPY