


SCEPTER 70DG herbicide may be applied in Minnesota, North Dakota, South Dakota, northern Michigan and northern Wisconsin.

SCEPTER 70DG can be applied in the states or parts of states shaded in the following map:




It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.

(R) Registered Trademark of American Cyanamid Company



CEPTED

1968

U.S. Federal Insecticide
Fungicide and Rodenticide Act
for the pesticide

EPA Reg. No. 241-306

FOR USE IN SOYBEANS

POSTEMERGENCE APPLICATIONS FOR COCKLEBUR CONTROL

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE
SCEPTER 70DG LEAFLET LABEL BEFORE USING

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner
inconsistent with its labeling.

When applied as directed, SCEPTER 70DG will provide control of
emerged common cocklebur (Xanthium strumarium) in soybeans.

SCEPTER 70DG RATES FOR POSTEMERGENCE COCKLEBUR CONTROL

WEED HEIGHT (inches)	SCEPTER (ounces/acre)	SURFACTANT OR COC*
up to 8	1.4	Yes
9 to 12	2.85	Yes

* Use a nonionic surfactant or crop oil concentrate.
The nonionic surfactant should contain at least 80%
active ingredient and should be applied at a rate
of 2 pints per 100 gallons of spray mixture. Apply
the crop oil concentrate (COC) at the rate stated
on the COC label.

DO NOT apply this product through any type of irrigation system.

This label should be in the possession of the user at the time of herbicide application.

(R) Registered Trademark of American Cyanamid Company

