

EPA Reg. No. 241-306

PL 25
241-306
Pg. 194

FOR USE IN SOYBEANS

**SCEPTER 70DG PREPLANT INCORPORATED OR
PREEMERGENCE APPLICATION FOLLOWED
BY CLASSIC¹ POSTEMERGENCE
FOR CONTROL OF SICKLEPOD**

For distribution and use only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee and Texas

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER 70DG AND CLASSIC LABELS BEFORE USING. SEE THE SCEPTER 70DG AND CLASSIC LABELS FOR SPRAYING AND APPLICATION INFORMATION.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product through any type of irrigation system.


ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

Apply SCEPTER 70DG at 2.8 ounces per acre preplant incorporated or preemergence in soybeans. A single postemergence application of Classic at a rate of 1/2 to 3/4 ounce per acre may be applied to control sicklepod.

Refer to the Classic label for weed height limitations. Use the high rate of Classic if sicklepod seedlings are taller than two inches. A cultivation approximately 14 days after Classic application may be required to control sicklepod escapes.

NOTE: If replanting is necessary in a field previously treated with SCEPTER 70DG, the field may be replanted to soybeans. Rework the soil no deeper than the treated zone. DO NOT apply a second treatment of SCEPTER 70DG.

¹Classic is a trademark of E.I. du Pont de Nemours and Company
*Registered trademark of American Cyanamid Company



soybeans.

1. Soybeans may be planted anytime.
2. Fifteen months after last herbicide application:

Barley	Grain sorghum	Rice
Edible beans	Oats	Tobacco
Field corn	Peanuts	Wheat
3. Sugarbeets may be planted 26 months after last herbicide application.
4. All other crops, including cotton, may be planted eighteen months after the last herbicide application unless specified otherwise in the PRECAUTIONS section of the Classic label.

Use of SCEPTER 70DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.


ONLY rotational crops harvested at maturity may be used for feed or food.


DO NOT graze or feed treated soybean forage, hay or straw to livestock.

This label must be in the possession of the user at the time of herbicide application.

PE-12138 3/13/80

 **CYANAMID**
American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07470 91988





FOR USE IN SOYBEANS

SCEPTER 70DG PREPLANT INCORPORATED OR PREEMERGENCE PLUS LINURON PREEMERGENCE TANK MIXTURE

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER 70DG AND LINURON LABELS BEFORE USING. SEE THE SCEPTER 70DG LABEL FOR USE AREA AND SPRAYING AND APPLICATION INFORMATION. SEE THE SCEPTER 70DG AND LINURON LABELS FOR WEED SPECIES CONTROLLED AND FOLLOW CROP RESTRICTIONS. ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DO NOT apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

NOTE: Linuron can cause soybean injury under some environmental conditions. DO NOT use this tank mix combination if soybean injury cannot be tolerated.

SCEPTER 70DG may be applied preemergence in tank mix combination with linuron (Lorox¹ or Linex²).

SCEPTER 70DG may also be applied preplant incorporated, followed by a preemergence application of linuron. The tank mix or sequential application of SCEPTER 70DG, and linuron will enhance control of common ragweed and lambsquarters in addition to the weeds controlled by a soil application of SCEPTER 70DG.

DO NOT apply this tank mixture postemergence.

DO NOT apply linuron preplant incorporated.

Apply SCEPTER 70DG at a rate of 2.8 ounces per acre. Refer to the SCEPTER 70DG label for registered tank mix options and rates. Refer to the linuron label for specific application rates, restrictions and precautionary instructions.

Only SCEPTER 70DG, PROWL, Lasso, or Dual may be applied in tank mix combination with linuron. If trifluralin is used then it must be incorporated prior to preemergence application of linuron.

¹Lorox, Classic, Canopy, Preview, Gemini, and Lorox Plus is a trademark of E.I. duPont de Nemours and Company

²Linex is a trademark of Griffin Ag. Products Co., Inc.

Lasso is a trademark of Monsanto Agricultural Products Company.

Dual is a trademark of Ciba-Geigy Corporation.

*Registered Trademark of American Cyanamid Company

PE-12183 2/16/90



American Cyanamid Company
Agricultural Division
Crop Protection Chemicals Department
Wayne, NJ 07090 2-1988

MIXING INSTRUCTIONS

Prior to mixing SCEPTER 70DG plus linuron tank mixtures, perform a jar mixing test to ensure compatibility of SCEPTER 70DG with the particular linuron formulation. When tank mixing SCEPTER 70DG with recommended herbicides, add the SCEPTER 70DG to the spray tank first and make sure it is thoroughly mixed before adding the other herbicide.

When tank mixing SCEPTER 70DG with PROWL[®] herbicide, Lasso[®], Dual[®], trifluralin or linuron, add the herbicides to the tank in the following order: (1) SCEPTER 70DG (2) PROWL, Lasso, Dual, or trifluralin, (3) linuron.

SPRAYING INSTRUCTIONS

Uniformly apply with properly calibrated ground equipment in 20 to 60 gallons of water or liquid fertilizer per acre. A spray pressure of 30 to 40 psi is recommended. Nozzle screens should be no smaller than 50 mesh.

DO NOT apply when wind velocity is greater than 10 mph.

Use of SCEPTER 70DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Preview, Gemini, Lorox Plus[®], etc.) or imazethapyr (e.g. PURSUIT[®] or FURSUIT[®] PLUS) the same year as SCEPTER 70DG or injury to future crops may occur.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

