

PH 25

241-306


12850

NOTIFICATION

SCEPTER[®] 70DG
EPA Reg. No. 241-306

Outer Package Printing

The enclosed wording is to be printed on the inside of the SCEPTER 70DG packaging and provides easy reference to this information. The SCEPTER 70DG rates referenced here are found on page 12 of the approved SCEPTER 70DG label. The SCEPTER 1.5ASU rates are given for comparison purposes and are found on page 12 of the approved SCEPTER label. The mixing instructions correspond to those on page 24 of the approved SCEPTER 70DG label. The dispersal time was noted as a reference to the user.



•Registered Trademark of American Cyanamid Company

12850



5 Acre Water-Soluble Packets

Each packet contains 14 ounces
and treats 5 acres, at 2.8 ounce rate
(for other rates, see RATE chart).

EASY TO USE:

- Begin filling spray tank, with agitation.
- Add ECO-PAK packets to water. Packets disperse in 1-2 minutes.
- Add other herbicides or additives.
- Finish filling tank and mix thoroughly.

ALL COPY AND LINE ART PRINT BLACK



RATE and CONVERSION CHART

ECO-PAK	SOIL APPLIED	POST APPLIED
Rate per acre	2.8 ounces	1.4 ounces
Acres per packet	5	10

SCEPTER 1.5 ASU

Rate per acre	2/3 pints	1/3 pints
Acres per gallon	12	24

NOTIFICATION

SCEPTER[®] 70DG
EPA Reg. No. 241-306

Printing on the Water Soluble Bag

The enclosed wording is to be printed on the water soluble bag for identification purposes. This information is taken from page 1 of the approved SCEPTER 70DG label.

[®]Registered Trademark of American Cyanamid Company

SCEPTER® 70DG
herbicide
ECO-PAK™

Contains: Imazaquin
EPA Reg. No. 241-306

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION!/¡PRECAUCION!

See additional Precautionary
Statements and Directions
For Use on outer container.



ALL COPY AND LINE ART PRINT BLACK

This Area Left Blank Intentionally

SCEPTER
70 DG herbicide

ECO-PAK
WATER-SOLUBLE PACKETS
FOR USE IN SOYBEANS

Each ECO-PAK Treats 5 Acres at the 2.8 ounce Rate.

ACTIVE INGREDIENT:

Imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-
5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)

INERT INGREDIENTS

TOTAL

EPA Reg. No. 241-306

SCEPTER 700G is a water soluble bag.

EPA Est.

**KEEP OUT OF REACH OF CHILDREN
CAUTION! ¡PRECAUCIÓN!**

PRECAUCIÓN AL USUARIO:

Si usted no lee inglés no use este producto hasta
que la etiqueta le haya sido explicada ampliamente.

In case of an emergency endangering life or property
involving this product, call collect, day or night,
Area Code 201-835-3100.

See Additional Precautionary Statements Inside
See Directions For Use Inside

Net Weight: 2 x 14 ounce ECO-PAK PACKETS
© Registered Trademark of American Cyanamid Company
™ Trademark of American Cyanamid Company

DO NOT remove packets from container
until immediate use.

STATEMENT OF PRACTICAL TREATMENT:

If on skin: Wash with plenty of soap and water.
If in eyes: Flush eyes with plenty of water and
physician: if irritation persists.

CYANAMID

FRONT LABEL

PRINTER INFORMATION:

SCEPTER 700G LOGO PRINTS 4 COLOR PROCESS, ECO-PAK LOGO AND 2
LINES OF COPY DIRECTLY BENEATH ECO-PAK LOGO PRINT BLACK. THE
1/8" HORIZONTAL RULE OVERPRINTS BLACK.
THE PANEL INDICATED BY THE HOLDING RULE PRINTS PMS 382, THE
EQUIVALENT OF GCM1 22. ALL COPY AND LINE ART PRINTS BLACK. THE
HOLDING RULE DOES NOT PRINT.

TABLE OF CONTENTS

	Page		Page
INGREDIENTS	Cover	HERBICIDAL APPLICATIONS	9
STATEMENT OF PRACTICAL TREATMENT	Cover	WEEDS CONTROLLED (Combination Treatments)	10
PRECAUTIONARY STATEMENTS	3	SCEPTER 700G Plus PROWL [®] Tank Mixture	11
STORAGE AND DISPOSAL	3	SCEPTER 700G Plus Trifluralin	11
DISCLAIMER	3	SCEPTER 700G Plus Lasso ¹	12
GENERAL INFORMATION	4	SCEPTER 700G Plus Dual ²	13
USE AREA	5	SCEPTER 700G Plus PROWL or Trifluralin Plus Command ³ Tank Mixture	13
WEEDS CONTROLLED (Soil-Applied Treatments)	6	SCEPTER 700G Plus Metribuzin Tank Mixture	14
WEEDS CONTROLLED (Postemergence Treatments)	8	Tank Mixtures with 2,4-DB	14
DIRECTIONS FOR CONVENTIONAL, MINIMUM, AND NO-TILL APPLICATIONS	8	Tank Mixtures with Bronco ⁴	15
APPLICATION RATE	8	Tank Mixtures with Gramoxone Extra ⁵	15
SOIL APPLICATIONS	9	Tank Mixtures with Roundup ⁶	15
PREPLANT INCORPORATED APPLICATIONS	9	MIXING INSTRUCTIONS	15
PREEMERGENCE APPLICATIONS (Including no-till)	9	SPRAYING INSTRUCTIONS	16
POSTEMERGENCE APPLICATIONS	9	ROTATIONAL CROP RESTRICTIONS	17
		WEED SCIENTIFIC NAMES	19

¹Lasso is a trademark of Monsanto Agricultural Products Company.

²Dual is a trademark of Ciba-Geigy Corporation.

³Command is a trademark of FMC Corporation.

⁴Roundup and Bronco are trademarks of Monsanto Agricultural Products Company.

⁵Gramoxone Extra is a trademark of ICI Americas, Inc.

⁶Registered Trademark of American Cyanamid Company.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION!

Causes eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if absorbed through skin. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with SCEPTER 700G. DO NOT use SCEPTER 700G other than in accordance with the instructions set forth on this label. The use of SCEPTER 700G not consistent with this label may result in injury to crops. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER DISPOSAL: DO NOT reuse outer packaging. Dispose of outer packaging in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burial. If burned, stay out of smoke.

DISCLAIMER

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of American Cyanamid Company. All such risks shall be assumed by the user.

American Cyanamid Company shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

American Cyanamid Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein.

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

described when used in accordance with the directions for use, subject to the risks referred to above. CYANAMID DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AMERICAN CYANAMID'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF SCEPTER 700G. In no case shall Cyanamid or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

USES WITH OTHER PRODUCTS (TANK MIXES)

If this product is used in combination with any other product except as specifically recommended in writing by American Cyanamid Company then American Cyanamid Company shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically recommended. If used in a combination recommended by American Cyanamid Company, the liability of American Cyanamid Company shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the American Cyanamid Company product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

GENERAL INFORMATION

SCEPTER 700G is a uniquely flexible broad-spectrum herbicide which may be applied preplant incorporated, preemergence or postemergence in soybeans.

When SCEPTER 700G is applied to the soil, susceptible weeds emerge, growth stops, and then the weeds either die or are not competitive with the crop.

When SCEPTER 700G is applied postemergence, absorption occurs through both the foliage and roots. Susceptible weeds stop growing and either die or are not competitive with the crop.

A timely cultivation may aid in the control of certain weeds or improve general weed control when adequate moisture is not received after application. Cultivation should be shallow. After postemergence treatments, wait at least 10 days before cultivating.

The mode of weed killing activity involves uptake of SCEPTER 700G by weed roots and/or foliage and rapid translocation to growing points. Therefore, adequate soil moisture is important for optimum SCEPTER 700G activity. The amount of rainfall or irrigation required following application depends on existing soil moisture, soil texture and organic matter content. Sufficient water to moisten the soil to a depth of 2 inches is normally adequate. If adequate moisture is not received within 7 days after treatment, then a cultivation or postemergence herbicide application may be needed to improve weed control. When adequate moisture is received after dry conditions, SCEPTER 700G will provide residual control of susceptible germinating weeds; activity on established weeds will

depend on the weed species and the location of its root system in the soil.

A soil treatment of SCEPTER 700G followed by a postemergence application of SCEPTER 700G will control certain problem weeds. Use sequential treatments in recommended states only.

Occasionally, internode shortening of soybean plants may be observed with SCEPTER 700G applications. This has no effect on soybean yields.

There must be an interval of at least 90 days between the last SCEPTER 700G application and soybean harvest. DO NOT graze or feed treated soybean forage, hay or straw to livestock.

PRECAUTIONS:

DO NOT use on crops other than soybeans. Crops other than soybeans, such as cotton, corn and vegetables, may be injured by spray drift or other indirect contact with SCEPTER 700G.

To avoid injury to sensitive crops from spray drift, follow all use directions and precautions in **SPRAYING INSTRUCTIONS** on page 16.

To avoid injury to sensitive crops, spray equipment used for SCEPTER 700G applications must be drained and thoroughly cleaned with water before being used to apply other products to these crops. See pages 17 and 18 for **ROTATIONAL CROP RESTRICTIONS**.


DO NOT use SCEPTER 700G other than in accordance with the instructions set forth on this label.

Use of SCEPTER 700G herbicide in accordance with label directions is expected to result in normal growth of rota-

tional crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, crop injury, is always possible.

USE AREA

SCEPTER 700G can be applied only in the states or parts of states shaded in the following map:



USE REGION 1 includes eastern Oklahoma, Arkansas, the Missouri Bootheel, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, and eastern Texas.

USE REGION 2 includes eastern Kansas (east of U.S. 81; the counties of Cloud, Ellsworth, Harvey, Jewell, Lincoln, Mitchell, McPherson, Ottawa, Republic, Saline, Sedgewick, and Sumner), southeastern Nebraska (east of U.S. 81, south of U.S. 34), Missouri, Illinois (south of S.R. 116 west of Peoria; south of U.S. 24 east of Peoria),

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK

Indiana, Ohio, Michigan, Kentucky, Virginia, West Virginia, Pennsylvania, Maryland, Delaware, and New Jersey. USE REGION 3 includes Nebraska (east of U.S. 81, north of U.S. 34), South Dakota (east of U.S. 81), Illinois (north of S.R. 116 west of Peoria; north of U.S. 24 east of Peoria), Wisconsin, Iowa, and southern Minnesota.

NOTE: See USE RATES and ROTATIONAL CROP RESTRICTIONS sections for recommendations applying to each Use Region.

WEEDS CONTROLLED SCEPTER 70DG Alone

When applied as directed, soil-applied treatments of SCEPTER 70DG will control the following weeds:

WEED	LEVEL OF CONTROL
BROADLEAF WEEDS	
Beggarweed, Florida	Control ¹
Bristly Starbur	Control
Burcucumber	Control ²
Cocklebur, Common	Control
Copperleaf, Hophornbeam	Control ⁴
Jimsonweed	Control
Lambsquarters, Common	Control
Mallow, Venice	Control
Mexicanweed	Control ¹

WEED	LEVEL OF CONTROL
BROADLEAF WEEDS	
Morningglory	Control ^{3,4}
Entireleaf	Control ^{3,4}
Ivyleaf	Control
Palmleaf	Control
Pink	Control
Smallflower	Control
Tail	Control ^{3,4}
Mustard species	Control
Nightshade, Eastern Black	Control
Nutsedge, Yellow	Suppression ²
Pigweed	Control
Palmer	Control
Redroot	Control
Smooth	Control
Spiny	Control
Waterhemp, Tail	Control
Poinsettia, Wild	Control
Puncturevine	Control
Rusley, Florida	Control
Ragweed, Common	Control
Giant	Control ³
Redweed	Control ²
Sesbania, Hemp	Control ⁴
Sicklepod	Control ⁴
Sida, Prickly (leafweed)	Control

6

WEED	LEVEL OF CONTROL
BROADLEAF WEEDS	
Smartweed, Ladysthumb	Control
Pennsylvania	Control
Spurge, Spotted	Suppression
Sunflower, Common	Control
Texasweed	Control ¹
Velvetleaf	Control ²
GRASSES	
Barnyardgrass	Suppression
Corn, Volunteer	Suppression
Foxtail	Control
Giant	Control
Green	Control
Yellow	Control
Goosegrass	Suppression
Johnsongrass, seedling	Control
Shallbarn	Suppression
Signalgrass, Broadleaf	Suppression

¹ In USE REGION 1 ONLY. A 2.8 ounces per acre soil application of SCEPTER 70DG will provide suppression of these weeds. SCEPTER 70DG must be applied postemergence at 2.8 ounces per acre following a soil application of SCEPTER 70DG at 2.8 ounces per acre in a planned sequential program to control these weeds. Apply SCEPTER 70DG before weeds exceed the 1 to 2 true leaf stage.

² To obtain consistent control or suppression of these weeds under a wide range of environmental conditions, a preplant incorporated application is required.

³ A 2.8 ounces per acre soil application of SCEPTER 70DG will provide suppression of these weeds. For control of these morningglory species, the sequential program in footnote #4 should be used.

⁴ USE REGION 1 and USE REGION 2. SCEPTER 70DG herbicide must be applied postemergence at 1 pint per acre following a soil application of SCEPTER 70DG for control of these weeds. The total amount of SCEPTER 70DG applied should not exceed 1 pint per acre in the Region 2 or 2 pints per acre in Use Region 1 in one season when following 2.8 ounces per acre of SCEPTER 70DG. Refer to the SCEPTER 70DG label for application instructions. In Use Region 2, refer to the Rotational Crop Restrictions section of this label for special instructions on planting follow crops when using this program.

⁵ Some broadleaf weeds such as giant ragweed germinating deep in the soil or with subsequent flushes may require a cultivation or postemergence herbicide application for season long control.

⁶ Only light to moderate infestations of sicklepod are controlled. Later sicklepod flushes may require a cultivation or a postemergence herbicide application. For heavy sicklepod infestations, the sequential program in footnote #1 may be used or Classic may be applied postemergence at 1/2 to 3/4 ounces per acre following a soil application of SCEPTER 70DG at 2.8 ounces per acre. Apply the higher rate of Classic if sicklepod seedlings are taller than 2 inches. A cultivation 14 days after Classic application may be required to control sicklepod escapes. Refer to the Rotational Crop Restrictions section of this label for instructions on planting follow crops when using this program. The Classic sequential program may be used only in Use Region 1 (except Oklahoma).

⁷ A preplant incorporated application is recommended for velvetleaf control. A postemergence application of SCEPTER 70DG will aid in the control and reduce competition from velvetleaf.

NOTE: The total amount of SCEPTER 70DG applied should not exceed 5.6 ounces per acre in one year.

7

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

WEEDS CONTROLLED BY POSTEMERGENCE APPLICATIONS

When applied as directed, postemergence treatments of SCEPTER 700G will control the following weeds:

WEEDS CONTROLLED*	MAXIMUM WEED HEIGHT CONTROLLED (inches)	
	1.4 ounces/ acre**	2.8 ounces/ acre**
Cocklebur, Common	8	12
Pigweed		
Palmer	N	12
Radroot	4	12
Smooth	4	12
Spiny	N	12
Tall waterhemp	N	12
Poinsettia, Wild	N	8
Sunflower, Common	4	8
Volunteer Corn	8	12

N = Not Controlled

Apply when weeds are actively growing. DO NOT apply SCEPTER 700G postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes.

USE REGION 1 and USE REGION 2. A sequential postemergence treatment of SCEPTER 700G at 1.4 ounces per acre (Use Region 1 and Use Region 2) or 2.8 ounces per acre (Use Region 1 only) is permitted following a soil application of SCEPTER 700G for additional control of the

above weeds. In Use Region 2, refer to the Rotational Crop Restrictions section of this label for special instructions on planting follow crops when using this program.

*Use a nonionic surfactant or crop oil concentrate (see page 9).

**Each soluble bag contains 14 ounces of SCEPTER 700G and will treat 10 acres at the 1.4 ounces per acre rate or 5 acres at the 2.8 ounces per acre rate.

DIRECTIONS FOR CONVENTIONAL, MINIMUM, AND NO-TILL APPLICATIONS

APPLICATION RATE

Apply SCEPTER 700G at a broadcast rate of 2.8 ounces per acre for preplant incorporated and preemergence (including minimum and no-till systems) applications. At this broadcast rate, one 14 ounce soluble bag of SCEPTER 700G will treat 5 acres of soybeans. Apply SCEPTER 700G at a broadcast rate of 1.4 (10 acres per soluble bag) or 2.8 (5 acres per soluble bag) ounces per acre for postemergence applications.

NOTE: For preplant incorporated and preemergence applications on sands and loamy sands, 2.15 ounces per acre (8.5 acres per soluble bag) can be used without a substantial reduction in the control of only cocklebur and pigweed.

NOTE: In Use Region 3 as defined by the Use Area section of this label, apply SCEPTER 700G prior to July 1.

APPLICATION INSTRUCTIONS

SOIL APPLICATIONS

SCEPTER 700G may be applied in conventional, minimum, or no-till as a preplant incorporated or preemergence application up to 45 days (30 days in Use Region 1) before planting. Preemergence applications may also be applied during or after planting but before the crop emerges.

Adequate moisture is required for activation of SCEPTER.

PREPLANT INCORPORATED APPLICATIONS

Apply SCEPTER 700G before planting and incorporate uniformly into the top 1 to 2 inches of soil.

SCEPTER 700G may be applied immediately before planting or up to 30 days prior to planting in Use Region 1. For Use Region 2 and Use Region 3, SCEPTER 700G may be applied 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or rolling cultivator.

PREEMERGENCE APPLICATIONS (including no-till)

Apply SCEPTER 700G up to 45 days (30 days in Use Region 1) before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate SCEPTER 700G in the soil. If sufficient rainfall or irrigation to activate SCEPTER 700G is not received within 7 days after application, a thorough shallow tillage, cultivation, or

postemergence treatment may be required for control of emerged weeds.

POSTEMERGENCE APPLICATIONS

SCEPTER 700G applied postemergence not only controls the emerged weed species listed previously, but can also provide residual control of susceptible weeds that may emerge after application. Application should be made to small soybeans so more herbicide reaches the soil surface. The level of residual control will depend on weed density and canopy closure which affect the amount of SCEPTER 700G reaching the soil surface and environmental conditions, such as rainfall, at the time of and following application.

To maximize weed control following a postemergence SCEPTER 700G application, wait at least 10 days before cultivating. This timely cultivation will enhance residual weed control, especially under dry conditions.

Use a nonionic surfactant or crop oil concentrate. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

There must be an interval of at least 90 days between the last SCEPTER 700G application and soybean harvest.

HERBICIDE COMBINATIONS

A soil treat rate of SCEPTER 700G is active against many grass species, but for maximum grass control SCEPTER 700G should be tank mixed with PROWL® herbicide, trifluralin, Lasso or Dual. SCEPTER 700G may also be applied as a sequential preemergence or postemer-

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

gence treatment following application of one of the above grass herbicides.

For no-till uses, SCEPTER 700G may be tank mixed with Gramoxone Extra, Roundup, or 2,4-DB to kill existing vegetation. Gramoxone Extra, Roundup, or 2,4-DB should be deleted from the tank mixture if vegetation is absent at the time of application. SCEPTER 700G may also be tank mixed with Bronco, a product which gives both preemergence control of annual grasses and postemergence control of existing vegetation. Use a minimum of 10 gallons of water per acre. Use higher gallonage for fields with dense vegetation or heavy crop residues. Use ground equipment only. Plant soybeans at least one inch deep and adjust planters to ensure adequate seed coverage.

SCEPTER 700G may be followed by a postemergence application of Fuslade[®] or Poast[®]. DO NOT tank mix SCEPTER 700G with postemergence grass herbicides.

When used in combination with another herbicide, SCEPTER 700G should be used only in accordance with recommendations on this label. Always follow the more restrictive label limitations and precautions. SCEPTER 700G cannot be mixed with any product containing a label prohibition against such mixing.

Soil Texture

Although the rate of SCEPTER 700G does not vary with soil texture, the recommended rates of other herbicides used with this product generally do vary with soil texture. Combination rate tables in this label refer to three soil texture groups: COARSE, MEDIUM, and FINE. The following table lists soil textures included in each of these three groups:

COARSE	MEDIUM	FINE
sand loamy sands sandy loams	sandy clay loams* sandy clays loams silt loams silt	silty clay loams* silty clays clay loams clays

* Sometimes considered transitional soils.

[†] Fuslade is a trademark of ICI Americas, Inc.

[‡] Poast is a trademark of BASF Wyandotte Corporation.

[§] Registered trademark of American Cyanamid Company.

ADDITIONAL WEEDS CONTROLLED BY COMBINATION TREATMENTS

When applied as directed, SCEPTER 700G preplant incorporated or preemergence combination treatments with PROWL, Trifluralin, Lasso or Dual will control, as indicated, weeds listed in the following table in addition to those controlled by SCEPTER 700G alone.

WEED	PROWL TriFluralin [®] Lasso Dual			
GRASSES				
BARNYARDGRASS	X	X	X	X
CRABGRASS, SMOOTH	X	X	X	X
CRABGRASS, LARGE	X	X	X	X
CROWFOOT GRASS	X	X		
GOOSEGRASS	X	X	X	X
PANICUM, FALL	X	X [†]	X	X

WEED	PROWL Trifluralin [§] Lasso Dual			
PANICUM, TEXAS	X	X		
SANDBUR, FIELD	X	X		
SHATTERCANE	X [‡]	X [†]		
SIGNALGRASS	X	X	X	X
WITCHGRASS	X	X	X	X
BROADLEAF WEEDS				
CARPETWEED	X	X	X	X
GALINSOGA			X	X
KOCHIA	X	X		
PURSLANE	X	X	X	
SPURGE, ANNUAL	X			

[†]Preplant incorporated treatments only.

[‡]See Note on page 11.

[§]See Note on page 12.

SCEPTER 700G Plus PROWL Tank Mixture

Apply SCEPTER 700G plus PROWL preplant incorporated or preemergence using rates in the table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. For preplant incorporated applications, incorporate into the soil within 7 days after application, and plant soybeans within 45 days after application (30 days in Use Region 1). For preemergence applications, apply from up to 45 days prior to planting (30 days in Use

Region 1) to a maximum of 2 days after planting. DO NOT apply if soybeans have emerged.

Broadcast Rate of SCEPTER 700G Plus PROWL

Less Than 3% Organic Matter			
Soil Texture	SCEPTER 700G [†] ounces/acre	PROWL 4EC [‡] pints/acre	PROWL 3.3EC [§] pints/acre
Coarse	2.0	1.0 to 1.5	1.2 to 1.8
Medium	2.0	1.5 to 2.0	1.8 to 2.4
Fine	2.0	1.5 to 2.0	1.8 to 2.4
3% or More Organic Matter			
Coarse	2.0	1.5	1.0
Medium	2.0	1.5 to 2.0	1.8 to 2.4
Fine	2.0	2.0 to 2.5	2.4 to 3.0

NOTE: Use the higher rate of PROWL if heavy weed populations are anticipated or if shattercane will be present.

Observe all precautions and limitations on PROWL label.

[†] One soluble bag contains 14 ounces of SCEPTER 700G and will treat 5 acres at the 2.0 ounces per acre rate.

SCEPTER 700G Plus Trifluralin Tank Mixture

Apply SCEPTER 700G plus trifluralin preplant incorporated using rates in the table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. Incorporate

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK

porate into the soil within 24 hours after application, and plant soybeans within 45 days after application (30 days in Use Region 1). Use equipment that provides uniform 2 inch incorporation.

Broadcast Rate of SCEPTER 700G Plus Trifluralin*		
Soil Texture	SCEPTER 700G** ounces/acre	Trifluralin* pints/acre
Coarse	2.8	1
Medium	2.8	1.5
Fine	2.8	2

NOTE: If fall panicum or shattercane will be present, refer to trifluralin label for higher rates to control these weeds.

Observe all precautions and limitations on the trifluralin label.

*Based on 4 pounds active ingredient per gallon.

**One soluble bag contains 14 ounces of SCEPTER 700G and will treat 5 acres at the 2.8 ounces per acre rate.

SCEPTER 700G Plus Lasso Tank Mixture

Apply SCEPTER 700G plus Lasso preplant incorporated or preemergence using rates in the table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. For preplant incorporated applications, apply and incorporate within 7 days prior to planting. For preemergence applications, apply up to 7 days prior, during or immediately after planting.

Broadcast Rate of SCEPTER 700G Plus Lasso

Soil Texture	Less than 3% Organic Matter		
	SCEPTER 700G* (ounces/acre)	Lasso (quarts/acre)	
		Preplant Incorporated	Preemergence
Coarse	2.8	2.0 to 2.5	2.0
Medium	2.8	2.5 to 3.0	2.0 to 2.5
Fine	2.8	2.5 to 3.0	2.0 to 3.0**
	3% or More Organic Matter		
	SCEPTER 700G*	Lasso (quarts/acre)	
		Preplant Incorporated	Preemergence
Coarse	2.8	2.0 to 2.5	2.0
Medium	2.8	2.5 to 3.0	2.0 to 2.5
Fine	2.8	2.5 to 3.0	2.5 to 3.0

*One soluble bag contains 14 ounces of SCEPTER 700G and will treat 5 acres at the 2.8 ounces per acre rate.

**On silty clay and heavy clay soils of the Mississippi Delta, use 2 to 2-1/2 quarts of Lasso.

NOTE: For preplant incorporated applications, use the higher rate of Lasso if heavy weed populations are anticipated.

Use the higher rate of Lasso (1) when heavy grass infestations are anticipated, (2) when fall panicum or crabgrass will be present or (3) when previous crop residues are extremely dense. Apply up to 7 days before, during or immediately after planting.

Observe all precautions and limitations on the Lasso label.

SCEPTER 700G Plus Dual Tank Mixture

Apply SCEPTER 700G plus Dual preplant incorporated or preemergence using rates in table below. Follow use directions under APPLICATION INSTRUCTIONS in this label. For preplant incorporated applications, apply up to 29 days prior to planting. Incorporate within 7 days after application. For preemergence applications, apply up to 29 days prior, during or immediately after planting.

Broadcast Rate of SCEPTER 700G Plus Dual

Soil Texture	Less than 3% Organic Matter	
	SCEPTER 700G* ounces/acre	DUAL pints/acre
Coarse	2.8	1.25 to 1.5
Medium	2.8	1.5 to 2.0
Fine	2.8	2.0 to 2.5
	3% or More Organic Matter	
	SCEPTER 700G*	DUAL
Coarse	2.8	1.5
Medium	2.8	2.0
Fine	2.8	2.0 to 2.5**

*One soluble bag contains 14 ounces of SCEPTER 700G and will treat 5 acres at the 2.8 ounces per acre rate.

**Use the higher rate of Dual if heavy weed populations are anticipated.

NOTE: Observe all precautions and limitations on Dual label.

SCEPTER 700G Plus PROWL or Trifluralin Plus Command Tank Mixture

Observe all precautionary statements and limitations in the PROWL or trifluralin and Command labels before using. The SCEPTER 700G plus PROWL or trifluralin plus Command 4EC preplant incorporated tank mixture may be used when heavy infestations of velvetleaf are anticipated. Uniformly apply the tank mixture with properly calibrated ground equipment in 10 to 40 gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application.

Apply SCEPTER 700G plus PROWL or trifluralin plus Command 4EC to the soil surface and uniformly incorporate. Refer to the Command 4EC label for incorporation instructions. DO NOT apply aerially or through irrigation equipment. DO NOT apply this tank mixture to overly moist or wet soils.

SPECIAL PRECAUTION

Command 4EC is a volatile compound. On-site movement of spray drift or vapors of Command can cause foliar whitening or yellowing of some plants. Prior to using Command, read and strictly follow all precautions and application instructions of the Command label. The use of an agriculturally approved drift reducing agent is required at finished spray volumes of 10 to 15 gallons per acre.

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

BROADCAST RATE PER ACRE OF SCEPTER 700G PLUS PROWL OR TRIFLURALIN PLUS COMMAND 4EC

Apply SCEPTER 700G at 2.8 ounces per acre*. Apply Command at 1/4 to 1/2 pint per acre. Refer to HERBICIDE COMBINATIONS section for PROWL (page 11) or trifluralin (page 11) use rates.

ROTATIONAL CROP RESTRICTIONS: Follow all rotational crop restrictions on the SCEPTER 700G, PROWL or trifluralin, and Command label.

ALWAYS FOLLOW THE MOST RESTRICTIVE LABEL.

*One soluble bag contains 14 ounces of SCEPTER 700G and will treat 5 acres at the 2.8 ounces per acre rate.

SCEPTER 700G Plus Metribuzin Tank Mixtures

Before using, observe all precautionary statements in the SCEPTER 700G, Sencor®, Lexone®, PROWL, Lasso, Dual, and trifluralin leaflet labels. SCEPTER 700G may be applied in tank mix combination with Sencor or Lexone plus one of the following grass herbicides: PROWL, Lasso, Dual, or trifluralin; either preplant incorporated or preemergence (excluding trifluralin). SCEPTER 700G may also be applied in tank mix with any one of the above grass herbicides as a preplant incorporated application followed by a preemergence application of Sencor or Lexone. For preemergence applications, apply during or immediately after planting. These treatments will control hemp sesbania and hophornbeam copperleaf in addition to the weeds controlled by a soil application of SCEPTER 700G plus the grass herbicide.

Apply Sencor or Lexone at the broadcast rate per acre specified in the table below. In tank mix combination with 2.8 ounces per acre* of SCEPTER 700G plus one of the above grass herbicides.

Sencor or Lexone Rate**		
	4.0L	75%DF
Coarse Soils	1/2 pint	1/3 lb.
Medium Soils	1/2 pint	1/3 lb.
Fine Soils	2/3 pint	1/2 lb.

*One soluble bag contains 14 ounces of SCEPTER 700G and will treat 5 acres at the 2.8 ounces per acre rate.

**Do not use Sencor or Lexone on coarse soils of less than 2% organic matter. Use 1/2 lb. or 2/3 pint of Sencor or Lexone on medium soils of greater than 4% organic matter.

*Sencor is a trademark of Mobay Company

*Lexone is a trademark of E I du Pont de Nemours and Company

Tank Mixtures with 2,4-DB

2,4-DB may be used with SCEPTER 700G alone or in combination with other SCEPTER 700G tank mixes for the control of some annual broadleaf weeds. Refer to the 2,4-DB label for specific use recommendations, rates and weeds controlled.

Observe all cautions and limitations on the 2,4-DB label.

Tank Mixtures With Bronco

Use the following rates of Bronco for SCEPTER 700G plus Bronco tank mixtures:

Soil Texture	Bronco (quarts/acre)
Coarse	3 to 5
Medium	4 to 5
Fine	4 to 5

Use the higher rate of Bronco (1) when previous crop residues are extremely dense, (2) when emerged weeds are over 6 inches tall, (3) when heavy infestations of weeds are expected, and (4) when soils contain more than 3% organic matter. Apply up to 7 days before, during or immediately after planting.

Observe all precautions and limitations on Bronco label.

Tank Mixtures With Gramoxone Extra

Apply 1-1/2 to 2-1/2 pints of Gramoxone Extra per acre in SCEPTER 700G tank mixtures. Use the 2-1/2 pint rate if weeds are 4 to 6 inches tall. Weeds over 6 inches may not be controlled with this treatment. Apply up to 14 days before, during or immediately after planting.

When Gramoxone Extra is included in a tank mixture, add a nonionic spreader surfactant at a rate of 8 fl. oz. per 100 gallons of spray mixture as the last ingredient in the tank.

Gramoxone Extra will control most annual emerged weeds and suppress many emerged perennials. Refer to the Gramoxone Extra label for specific use recommendations and weeds controlled.

Observe all precautions and limitations on Gramoxone Extra label.

Tank Mixtures With Roundup

Roundup may be used with SCEPTER 700G alone or in combination with other tank mixes for the control of certain emerged grasses and broadleaf weeds. Refer to the Roundup label for specific use recommendations, rates and weeds controlled.

Observe all cautions and limitations on the Roundup label.

MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. While agitating add the required amount of SCEPTER 700G (including the water soluble packaging) and then fill the remainder of the tank with water. For postemergence applications, add the nonionic surfactant or the crop oil concentrate as last ingredient in the tank. An antifoaming agent may be added to the tank if needed. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank mixing SCEPTER 700G with recommended herbicides, add SCEPTER 700G (including the water soluble packaging) to the spray tank first and make sure it is thoroughly mixed before adding the other herbicide.

When tank mixing Roundup or Gramoxone Extra with SCEPTER 700G and PROWL, Lasso or Dual, add herbicides to the tank in the following order: (1) SCEPTER 700G (2) PROWL, Lasso or Dual, (3) Roundup or Gramoxone Extra, and (4) nonionic surfactant.

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK

SPRAYING INSTRUCTIONS

NOTE: DO NOT make applications when spray may be carried by wind to sensitive crops.

Avoid overlaps when spraying.

GROUND APPLICATIONS:

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. Use sprayers equipped with nozzles that provide accurate and uniform application. To minimize drift, use a maximum spray pressure of 40 psi.

To ensure thorough coverage, use a minimum of 20 gallons of water per acre when applying SCEPTER 700G as a postemergence treatment. Use higher gallonage for fields with dense vegetation or heavy crop residues.

DO NOT apply with ground equipment when wind velocity is greater than 10 mph.

For band treatment, apply the broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Rate per acre} = \text{Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per acre} = \text{Band VOLUME per acre}$$

AERIAL APPLICATIONS:

Uniformly apply with properly calibrated aerial equipment in 5 or more gallons of water per acre.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS:

- Use nozzles which produce a coarse spray.
- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed toward the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph.

Applicator is responsible for any loss or damage which results from spraying SCEPTER in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regard to spraying.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted after applying SCEPTER at recommended rates in soybeans:

CROP	USE REGION 1	USE REGION 2 (excl. Michigan**)	USE REGION 3
Soybeans	No restrictions	No restrictions	No restrictions
Wheat	4 months	4 months***	See NOTE
Rice	Spring following SCEPTER 700G application	Spring following SCEPTER 700G application***	—
Barley	11 months	11 months***	See NOTE
Field Corn	11 months*	11 months*****	See NOTE
Edible Beans	11 months	11 months***	11 months
Grain Sorghum	11 months	11 months***	11 months
Oats	11 months	11 months***	See NOTE
Peanuts	11 months	11 months***	11 months
Tobacco	11 months	11 months***	11 months
Sugar Beets	26 months	26 months	26 months

NOTE: For USE REGION 3 as defined in the USE AREA section of this label, field corn, wheat, barley, or oats may be planted as a rotational crop in the fall or in the spring of the year following SCEPTER 700G application ONLY if SCEPTER 700G was applied as a postemergence application at 1.4 ounces per acre and at least 10 inches of rainfall or irrigation has been received from the date of application through October of the same year.

*For USE REGION 1 as defined in the USE AREA section of this label, field corn may be planted in the spring of the year following SCEPTER 700G application unless extreme drought conditions develop (less than 15 inches of rainfall or irrigation is received within 6 months following the date of last application). A minimum of 10 inches of rainfall is needed for a postemergence application of SCEPTER 700G only if the total amount of SCEPTER 700G applied in one season does not exceed 1.4 ounces per acre.

**For USE REGION 2 (except Michigan) as defined in the USE AREA section of this label, field corn may be planted in the spring of the year following SCEPTER 700G application unless extreme drought

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

conditions develop (less than 15 inches of rainfall or irrigation is received from two weeks prior to the date of last application through November 15 of the same year). A minimum of 18 inches of rainfall is needed for a postemergence application of SCEPTER 700G only. The total amount of SCEPTER 700G applied in one season does not exceed 1.4 ounces per acre.

If the minimum rainfall requirements are not met, only field corn hybrids which are classified as tolerant or resistant to SCEPTER 700G and/or other midazolinone herbicides may be planted the spring of the year following SCEPTER 700G application. Contact your chemical dealer, seed supplier or American Cyanamid to obtain a list of SCEPTER 700G tolerant hybrids.

DO NOT plant field corn, oats, or barley in Michigan in the fall or spring of the year following SCEPTER 700G application.

ROTATIONAL CROP RESTRICTIONS for sequential SCEPTER 700G or SCEPTER OT application following a soil application of SCEPTER 700G (USE REGION 2 only):

For USE REGION 2 as defined in the USE AREA section of this label, soybeans may be planted anytime. Barley, edible beans, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application.

Only field corn hybrids which are classified as tolerant or resistant to SCEPTER 700G and/or other midazolinone herbicides may be planted the spring of the year following a sequential SCEPTER 700G application. Contact your chemical dealer, seed supplier, or American Cyanamid to obtain a list of SCEPTER 700G tolerant field corn hybrids. Other field corn varieties may be planted 15 months after the last herbicide application.

ROTATIONAL CROP RESTRICTIONS for sequential Classic application following SCEPTER 700G soil treatments (USE REGION 1 only):

Soybeans may be planted anytime. Barley, edible beans, field corn, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after

the last herbicide application. Cotton may be planted 18 months after the last herbicide application. Refer to rotational crop restrictions listed in the Classic label. Always follow the more restrictive label.

Only rotational crops harvested at maturity may be used for feed or food.

DO NOT plant other rotational crops prior to 18 months after last SCEPTER 700G application.

Use of SCEPTER 700G herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, crop injury, is always possible.

Except where stated on a Supplemental Label or in this label, DO NOT apply products containing chlorimuron ethyl (e.g. Classic, Canopy, Gemini, Loric Plus, Preview[®], etc.) or imazamox (SCEPTER, SQUADRON[®], TRI-SCEPT, SCEPTER-O.T.[®]) the same year as SCEPTER. DO NOT apply products containing imazethapyr (e.g. PURSUIT[®], PURSUIT Plus or PASSPORT[®]) the same year as SCEPTER 700G.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than 2 inches.

DO NOT graze or feed treated soybean forage, hay or straw to livestock.

[®]Classic, Canopy, Gemini, Loric Plus and Preview are trademarks of E I du Pont de Nemours and Company.

[®]Registered Trademarks of American Cyanamid Company.

WEED SCIENTIFIC NAMES

BROADLEAF WEEDS	
Beggarweed, Florida	(<i>Desmodium illinoense</i>)
Bristly Starbur	(<i>Acanthospermum hispidum</i>)
Burcucumber	(<i>Sicyos angulatus</i>)
Carpetweed	(<i>Mollis verticillata</i>)
Cockspur, Common	(<i>Xanthium strumarium</i>)
Copperleaf, Hophornbeam	(<i>Acalypha ostryaefolia</i>)
Galinsoga	(<i>Galinsoga spp.</i>)
Jimsonweed	(<i>Datura stramonium</i>)
Kochia	(<i>Kochia scoparia</i>)
Lambsquarters, Common	(<i>Chenopodium album</i>)
Mimosa, Venice	(<i>Hibiscus trionum</i>)
Mexicanweed	(<i>Caperonia coccinea</i>)
Morningglory	
Entireleaf	(<i>Ipomoea hederacea</i> var. <i>intergracula</i>)
Wyleaf	(<i>Ipomoea hederacea</i>)
Painted	(<i>Ipomoea wrightii</i>)
Sm-Mallow	(<i>Ipomoea lacunosa</i>)
Tall	(<i>Jacquemontia tamnifolia</i>)
	(<i>Ipomoea purpurea</i>)

BROADLEAF WEEDS

Mustard species	(<i>Brassica</i> spp.)
Nightshade, Eastern Black	(<i>Solanum phycanthum</i>)
Nutsedge, Yellow	(<i>Cyperus esculentus</i>)
Pigweed	
Palmer	(<i>Amaranthus palmeri</i>)
Redroot	(<i>Amaranthus retrofractus</i>)
Smooth	(<i>Amaranthus hybridus</i>)
Spiny	(<i>Amaranthus spinosus</i>)
Waterhemp, Tall	(<i>Amaranthus tuberculatus</i>)
Poinsettia, Wild	(<i>Euphorbia heterophylla</i>)
Puncturevine	(<i>Tribulus terrestris</i>)
Purslane	(<i>Portulaca oleracea</i>)
Pustle, Florida	(<i>Richardia scabra</i>)
Ragweed, Common Giant	(<i>Ambrosia artemisiifolia</i>)
	(<i>Ambrosia trifida</i>)
Redweed	(<i>Melochia coccinea</i>)
Sesbania, Hemp	(<i>Sesbania exaltata</i>)
Sicklepod	(<i>Cassia obtusifolia</i>)
Sida, Prickly (tearweed)	(<i>Sida spinosa</i>)
Smartweed, Ladystumb Pennsylvania	(<i>Polygonum persicaria</i>)
	(<i>Polygonum pennsylvanicum</i>)

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK

BROADLEAF WEEDS

Spurge
Annual (Euphorbia spp.)
Spotted (Euphorbia maculata)

Sunflower, Common (Helianthus annuus)

Texasweed (Caperonia palustris)

Velvetleaf (Abutilon theophrasti)

GRASSES

Barnyardgrass (Echinochloa crus-galli)

Corn, Volunteer (Zea mays)

CRABGRASS

Smooth (Digitaria ischaemum)

Large (Digitaria sanguinalis)

Crowfoot Grass (Dactyloctenium aegyptium)

GRASSES

Foxtail
Giant (Setaria faberii)
Green (Setaria viridis)
Yellow (Setaria glauca)

Goosegrass (Eleusine indica)

Johnsongrass, seedling (Sorghum halepense)

Panicum
Fall (Panicum dichotomiflorum)
Texas (Panicum texanum)

Sandbar, Field (Cenchrus incertus)

Shattercane (Sorghum bicolor)

Signalgrass, Broadleaf (Brachiaria platyphylla)

Witchgrass (Panicum capillare)

NOTES

20

21

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK

NOTES

NOTES

22

23

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

A:\7dgws14.chp 305.7632
Date: 10/19/92, 10/27

SCEPTER 70DG ECO-PAK 14 OUNCE
INBOOK, HERO, LIDDING STOCK AND PACKET BAG

CLIENT REVIEW - 2
TLA

NOTES

NOTES

24

25

PRINTER INFORMATION:
ALL COPY AND LINE ART PRINT BLACK.

10/27/92



INBOOK™ is protected by the following Patents:
U.S.A. 4744161, 4680080, 4592572, 4675082, Canada 422793, and others.

25

SCEPTER

70 DG herbicide

FOR USE IN SOYBEANS **ECO-PAK™**

ACTIVE INGREDIENT:

Imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid) 70.0%

INERT INGREDIENTS 30.0%

TOTAL 100.0%

SCEPTER 70DG is a water soluble bag.

EPA Reg. No. 241-308

EPA Est. No. 241-PR-2

KEEP OUT OF REACH OF CHILDREN

CAUTION!
¡PRECAUCION!

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT
If on skin: Wash with plenty of soap and water.
If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

DO NOT remove packets from container except for immediate use.



Net wt.: 2 x 14 oz. ECO-PAK PACKETS

© Registered Trademark of American Cyanamid Company
™ Trademark of American Cyanamid Company

10/92

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS

CAUTION

Causes eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if absorbed through skin. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with SCEPTER 70DG. DO NOT use SCEPTER 70DG other than in accordance with the instructions set forth on this label. The use of SCEPTER 70DG not consistent with this label may result in injury to crops. Keep container closed to avoid spills and contamination.

DO NOT apply this product through any type of irrigation system.

See Leaflet For Complete Directions For Use.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

CONTAINER DISPOSAL: DO NOT reuse outer packaging. Dispose of outer packaging in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

See Leaflet For Disclaimer.

In case of an emergency endangering life or property involving this product, call collect, day or night, Area Code 201-835-3100.

TEAR ALONG PERFORATION TO REMOVE

ATTACHED LABEL

BACK PANEL PRINTER INFORMATION:
STRIP IN SCEPTER 70DG 4-COLOR LOGO ART TO SIZE AND POSITION INDICATED. ALL REMAINING COPY AND LINE ART PRINTS BLACK.

SCEPTER[®]

70 DG herbicide

ECO-PAK[™]
WATER-SOLUBLE PACKETS



a:\alt021ghcdr

Date: 10/17/92, 10/20, 10/27

SCEPTER 70 DG ECO-PAK 14 OZ.HERO

Version A

305.7632

TLA