

MAY 24 1993

Dr. Zareen Ahmed
American Cyanamid Co.
P.O. Box 400
Princeton, NJ 08543

Dear Dr. Ahmed:

Subject: Supplemental Label- Tank Mix with Pursuit
SCEPTER® 70 DG HERBICIDE
EPA Reg. No. 241-306
Your letter dated May 18, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comment listed below. A stamped copy is enclosed for your records.

Sincerely yours,

Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

RD:RTaylor:eja:5/24/93:3056800

EPA Reg. No. 241-306

ACCEPTED
with COMMENTS
in EPA Letter Dated:

MAY 24 1993

FOR USE IN SOYBEANS

**SCEPTER 70DG PLUS PURSUIT® HERBICIDE
FOR SUPPRESSION OF VOLUNTEER CORN**

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

**FOR USE IN SCEPTER USE REGION 2 AND USE REGION 3
(EXCLUDING IOWA, MINNESOTA AND WISCONSIN)**

241-306

THIS LABEL EXPIRES DECEMBER 31, 1993

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER 70DG AND PURSUIT LABELS BEFORE USING. SEE THE PURSUIT LABEL FOR HERBICIDE APPLICATION INFORMATION AND WEEDS CONTROLLED. WHEN TANK MIXING, ALWAYS FOLLOW THE MORE RESTRICTIVE LABEL.

GENERAL INFORMATION

Apply SCEPTER 70DG at 0.53 ounces per acre in combination with PURSUIT at 4.0 fluid ounces per acre postemergence for suppression of volunteer corn. At this broadcast rate, one 14 ounce soluble bag of SCEPTER 70DG will treat 26.4 acres.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

Follow all MIXING, SPRAYING and APPLICATION instructions in the PURSUIT leaflet label.

USE RATE AND APPLICATION INSTRUCTIONS

Apply SCEPTER 70DG at 0.53 ounces per acre in combination with PURSUIT at 4.0 fluid ounces per acre postemergence. At this broadcast rate, one 14 ounce soluble bag of SCEPTER 70DG will treat 26.4 acres. Apply before the corn is ten inches tall.

USE AREA

This tank-mix application of SCEPTER 70DG and PURSUIT may be used in Region 2 and Region 3 (excluding Iowa, Minnesota and Wisconsin) as described in the USE AREA section of the SCEPTER 70DG label.

Refer to the SCEPTER 70DG leaflet label for a listing of those states included in USE REGION 2 and the approved use area of USE REGION 3.

WEEDS SUPPRESSED - VOLUNTEER CORN

This tank-mixture will provide suppression of volunteer corn in soybeans in addition to those weeds controlled by a postemergence application of PURSUIT. Refer to the PURSUIT label for a listing of weeds controlled by a postemergence application of PURSUIT.

A postemergence application of SCEPTER 70DG will **NOT** suppress those field corn hybrids which possess tolerance or resistance to SCEPTER 70DG or other imidazolinone herbicides (i.e. PURSUIT).

ROTATIONAL CROP RESTRICTIONS.....

SCEPTER 70DG USE REGION 2: Follow all rotational crop restrictions on the PURSUIT and SCEPTER 70DG leaflet labels for that area designated as SCEPTER 70DG USE REGION 2 on the SCEPTER 70DG leaflet label. Always following the most restrictive label.

FOR THIS SUPPLEMENTAL LABEL ONLY: THE ENTIRE STATE OF ILLINOIS MAY FOLLOW THE ROTATIONAL CROP RESTRICTION GUIDELINES FOR SCEPTER USE REGION 2.

(See other side)

1. Anytime:
Soybeans
2. Four months after SCEPTER 70DG plus PURSUIT application:
Wheat
3. Eleven months after SCEPTER 70DG plus PURSUIT application:
Field corn (resistant)*
4. Eighteen months after SCEPTER 70DG plus PURSUIT application:

Alfalfa	Popcorn
Cotton	Safflower
Field corn	Sorghum
Lettuce	Sweet corn
Oats	
5. Twenty six months after SCEPTER 70DG plus PURSUIT application:
All crops not listed elsewhere in this ROTATIONAL CROP GUIDELINE.
6. Forty months after SCEPTER 70DG plus PURSUIT application:
Sugar beets

corn hybrids. Other field corn hybrids may be planted 18 months after application of this tank-mixture.

NOTE: Availability of imidazolinone resistant field corn seed cannot be assured.

Use of SCEPTER 70DG and PURSUIT herbicides in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of these products and, therefore, rotational crop injury is always possible.

EXCEPT as stated above, **DO NOT** use this tank mixture in the same year as an application of SCEPTER, SQUADRON®, TRI-SCEPT®, SCEPTER® O.T.®, PURSUIT, or PURSUIT PLUS®. Also, **DO NOT** use the above tank mixture in the same year as an application of a product containing chlorimuron ethyl (e.g. Classic¹, Canopy¹, Gemini¹, Lorox Plus¹, Preview¹, etc.)

*Registered Trademark of American Cyanamid Company
¹Trademark of E. I. duPont de Nemours & Company, Inc.
 PE- 4/5/93 SD

 **CYANAMID**
 American Cyanamid Company
 Agricultural Products Division
 Crop Protection Chemicals Department
 Wayne, NJ 07470 ©1993