

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 241-306

EPA Reg. No. 241-306

## FOR USE IN SOYBEANS

### SCEPTER 70DG AND ROUNDUP<sup>1</sup> FOR WEED CONTROL IN SOYBEANS AND SCEPTER 70DG USE IN ROUNDUP READY<sup>2</sup> SOYBEANS

OBSERVE ALL PRECAUTIONARY STATEMENTS IN THE SCEPTER 70DG LEAFLET LABEL BEFORE USING. SEE THE SCEPTER 70DG LEAFLET LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, USES WITH OTHER PRODUCTS AND ROTATIONAL CROP RESTRICTIONS.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of herbicide application. The use of SCEPTER 70DG herbicide not consistent with this labeling may result in injury to crops. Keep container closed to avoid spills and contamination. **DO NOT** apply this product through any type of irrigation system.

#### GENERAL INFORMATION

SCEPTER 70DG provides control of broadleaf and grass weeds when applied preplant, preemergence and postemergence. SCEPTER 70DG can be applied to either conventional or no-till soybeans. When applied as a burndown application, Roundup may be tank-mixed with SCEPTER 70DG for control of weeds common to no-till production, such as marehail.

SCEPTER 70DG may also be applied to Roundup Ready soybeans.

Refer to the SCEPTER 70DG label and the Roundup label for specific use recommendations, rates and weeds controlled.

##### No-till Soybeans (non-glyphosate resistant)

Apply SCEPTER 70DG plus Roundup for weed control in no-till soybeans. Roundup will enhance the control of weeds including marehail and prickly lettuce. Always include a nonionic surfactant in the spray solution. **DO NOT** apply Roundup postemergence to non-glyphosate resistant soybeans.

##### Roundup Ready Soybeans (glyphosate resistant)

SCEPTER 70DG may be used for weed control in Roundup Ready soybeans. Apply SCEPTER 70DG either preplant soil applied, preplant as a burndown treatment, or for postemergence weed control. SCEPTER 70DG may be applied either prior to Roundup, or following Roundup for general weed control and residual activity on broadleaf and grass weeds. For sequential treatments, a sufficient time period should occur between treatments to allow an appropriate assessment of weed control needs.

SCEPTER 70DG may be tank-mixed with other herbicides (not Roundup) on Roundup Ready soybeans when applied postemergence. Refer to the SCEPTER 70DG label for recommended herbicides. Refer to the other herbicide labels for weeds controlled and recommended application rates.

#### MIXING INSTRUCTIONS

Fill the spray tank one-fourth to one-half full with clean water. When tank-mixing SCEPTER 70DG with recommended herbicides, while agitating, add the required amount of SCEPTER 70DG (including the water soluble packaging) to the spray tank first and make sure it is thoroughly mixed before adding the other herbicide; then fill the remainder of the tank with clean water. Maintain agitation while spraying to ensure a uniform spray mixture. For postemergence applications, add the nonionic surfactant as the last ingredient in the tank. To avoid injury to sensitive crops, spray equipment used for

(See other side)

10/2

# SCEPTER 70DG HERBICIDE USE AREA

See SCEPTER 70DG leaflet label for use area and a description of Use Regions.

## APPLICATION INSTRUCTIONS

### SOIL APPLICATIONS

SCEPTER 70DG may be applied in conventional, minimum, or no-till soybeans as a preplant incorporated or preemergence application up to 45 days (30 days in Use Region 1) before planting. Preemergence applications may also be applied during or after planting but before the crop emerges.

Adequate moisture is required for activation of SCEPTER 70DG.

### PREPLANT INCORPORATED APPLICATIONS

Apply SCEPTER 70DG before planting and incorporate uniformly into the top 1 to 2 inches of soil.

SCEPTER 70DG may be applied immediately before planting or up to 30 days prior to planting in Use Region 1. For Use Region 2 and Use Region 3, SCEPTER 70DG may be applied 45 days prior to planting. Incorporate within 7 days after application.

If soybeans are planted on beds, apply and incorporate after bed formation using PTO-driven equipment or rolling cultivator.

### PREEMERGENCE APPLICATIONS (including no-till)

Apply SCEPTER 70DG up to 45 days (30 days in Use Region 1) before, during or after planting but before the crop emerges. As with other herbicides applied preemergence, rainfall or irrigation is necessary to activate SCEPTER 70DG in the soil. If sufficient rainfall or irrigation to activate SCEPTER 70DG is not received within 7 days after application, a thorough shallow tillage, cultivation, or postemergence treatment may be required for control of emerged weeds.

Refer to the SCEPTER 70DG label and Roundup label for additional information regarding WEEDS CONTROLLED, MIXING AND SPRAYING INSTRUCTIONS, PRECAUTIONARY STATEMENTS AND WORKER PROTECTION INFORMATION. Refer to the Roundup label for additional information regarding application rates, restrictions, weeds controlled and other information.

**When tank-mixing herbicides, always follow the most restrictive label.**

Use of SCEPTER 70DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

---

®Registered Trademark of American Cyanamid Company

<sup>1</sup> Roundup is a trademark of Monsanto Agricultural Products Company.

<sup>2</sup> Roundup Ready is a trademark of Monsanto Agricultural Products Company.

PE-XXXX 7/7/95 SD



American Cyanamid Company  
Agricultural Products Division  
Crop Protection Chemicals Department  
Wayne, NJ 07470 1995