

SCEPTER®
70 DG herbicide

Supplemental Labeling

EPA Reg. No. 241-306

FOR USE IN SOYBEANS
THIS LABEL EXPIRES DECEMBER 31, 1998

REVISED TOBACCO ROTATIONAL INTERVAL

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER 70 DG LABEL BEFORE USING. SEE THE SCEPTER 70 DG LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

ROTATIONAL CROP RESTRICTIONS

Tobacco may be planted 9½ months following an application of SCEPTER 70 DG at up to 2.8 ounces per acre and no more than a total of 0.125 pounds of imazaquin applied per acre.

Use of SCEPTER 70 DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER 70 DG and, therefore, rotational crop injury is always possible.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

DEC - 4 1996

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

241-306



SCEPTER®
70 DG herbicide

Supplemental
Labeling

EPA Reg. No. 241-306

FOR USE IN SOYBEANS
THIS LABEL EXPIRES DECEMBER 31, 1998

REVISED WHEAT ROTATIONAL INTERVAL

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER 70DG LABEL BEFORE USING. SEE THE SCEPTER 70DG LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION, WEEDS CONTROLLED, USE AREA, OTHER ROTATIONAL CROP RESTRICTIONS AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

ROTATIONAL CROP RESTRICTIONS

In Use Regions 1 and 2 east of Interstate Highway 35, wheat may be planted 3 months following a soil application of SCEPTER 70DG at a rate up to 2.8 ounces per acre or a postemergence application up to 1.4 ounces per acre.

Use of SCEPTER 70DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER 70DG and, therefore, rotational crop injury is always possible.

ACCEPTED
~~with COMMENTS~~
In EPA Letter Dated

DEC - 4 1996

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

241-306

SCEPTER[®]
70 DG herbicide

3 7 3
**Supplemental
Labeling**

EPA Reg. No. 241-306

FOR USE IN SOYBEANS
THIS LABEL EXPIRES APRIL 30, 1998

SCEPTER 70DG followed by Classic¹ for Sicklepod Control in Soybeans

OBSERVE ALL PRECAUTIONARY STATEMENTS AND LIMITATIONS IN THE SCEPTER 70DG LABEL BEFORE USING. SEE THE SCEPTER 70DG LABEL FOR PRECAUTIONS, WORKER PROTECTION STANDARD REQUIREMENTS, APPLICATION INFORMATION AND USES WITH OTHER PRODUCTS.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product through any type of irrigation system. This label must be in the possession of the user at the time of herbicide application.

An early preplant, preplant incorporated, or preemergence application of SCEPTER 70DG may be made at the 2.8 ounces per acre rate. For heavy sicklepod infestations Classic may be applied postemergence at 1/2 to 3/4 ounces per acre following a soil application of SCEPTER 70DG. Apply the higher rate of Classic if sicklepod seedlings are taller than 2 inches. A cultivation 14 days after Classic application may be required to control sicklepod escapes. SCEPTER 70DG followed by Classic sequential program may be used only in Use Region 1 (except Oklahoma), Virginia and Kentucky.

ROTATIONAL CROP RESTRICTIONS

Soybeans may be planted anytime. Field Corn may be planted 9 1/2 months after the last herbicide application. Barley, edible beans, grain sorghum, oats, peanuts, rice, tobacco, and wheat may be planted 15 months after the last herbicide application. Cotton may be planted 18 months after the last herbicide application. Refer to rotational crop restrictions listed in the Classic label. Always follow the more restrictive label.

Consult the SCEPTER 70DG label for spraying and application information, weeds controlled, and use rates. When tank mixing, always follow the more restrictive label.

Use of SCEPTER 70DG herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risk associated with the use of SCEPTER 70DG and, therefore, rotational crop injury is always possible.

 **CYANAMID**
American Cyanamid Company
Agricultural Products Division
Crop Protection Chemicals Department
Wayne, NJ 07470 ©1996

ACCEPTED
~~WITH COMMENTS~~
In EPA Letter Dated

DEC - 4 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

[®] Registered Trademark of American Cyanamid Company

¹ Registered trademark of E. I. du Pont de Nemours and Company

241-306